

THÈSE DE DOCTORAT

Soutenue à Aix-Marseille Université
en cotutelle avec l'université Cheikh Anta Diop de Dakar
le 19 décembre 2024 par

Manthia DIALLO

Les filarioses humaines au Sénégal : dépistage et épidémiologie

Discipline

Biologie santé

Spécialité

Maladies infectieuses

École doctorale

ED 62 Sciences de la vie et la santé

Laboratoire/Partenaires de recherche

- Maladies Infectieuses Négligées (MINES)
- IHU Méditerranée infection /MEPHI

Composition du jury


Pr Christelle POMARES	Rapporteuse
Université de Côte d'Azur, Nice	
Pr Doudou SOW	Rapporteur
Université Gaston Berger de Saint-Louis, Sénégal	
Pr Mame Cheikh SECK	Examineur
Université Cheikh Anta Diop de Dakar, Sénégal	
Pr Laurence CAMOIN	Présidente du jury
Université d'Aix-Marseille	
Dr Oleg MEDIANNIKOV	Directeur de thèse
Université d'Aix-Marseille	
Dr Cheikh SOKHNA	Co-directeur de thèse
Institut de Recherche pour le Développement, France	

Affidavit

Je soussigné, Manthia DIALLO, déclare par la présente que le travail présenté dans ce manuscrit est mon propre travail, réalisé sous la direction scientifique de **Dr Oleg MEDIANNIKOV** et du **Dr Cheikh SOKHNA**, dans le respect des principes d'honnêteté, d'intégrité et de responsabilité inhérents à la mission de recherche. Les travaux de recherche et la rédaction de ce manuscrit ont été réalisés dans le respect à la fois de la charte nationale de déontologie des métiers de la recherche et de la charte d'Aix-Marseille Université relative à la lutte contre le plagiat.

Ce travail n'a pas été précédemment soumis en France ou à l'étranger dans une version identique ou similaire à un organisme examinateur.

Fait à Marseille, le 19 décembre 2024




Cette œuvre est mise à disposition selon les termes de la Licence Creative Commons Attribution - Pas d'Utilisation Commerciale - Pas de Modification 4.0 International.

Affidavit

I, undersigned, Manthia DIALLO, hereby declare that the work presented in this manuscript is my own work, carried out under the scientific supervision of **Dr Oleg MEDIANNIKOV** and **Dr Cheikh SOKHNA**, in accordance with the principles of honesty, integrity and responsibility inherent to the research mission. The research work and the writing of this manuscript have been carried out in compliance with both the French national charter for Research Integrity and the Aix-Marseille University charter on the fight against plagiarism.

This work has not been submitted previously either in this country or in another country in the same or in a similar version to any other examination body.

Place Marseille, date december, 19 , 2024



Cette œuvre est mise à disposition selon les termes de la Licence Creative Commons Attribution - Pas d'Utilisation Commerciale - Pas de Modification 4.0 International.

Liste des publications et participation aux conférences et congrès

I. Liste des publications réalisées dans le cadre de ma thèse (premier auteur) :

1. **Diallo Manthia**, Hubert Bassene, Georges Diatta, Florence Fenollar, Cheikh Sokhna, Oleg Mediannikov.
Human filariasis in Senegal: screening and epidemiology. (Review in progress)
2. **Diallo Manthia**, Bassene Hubert, Diatta Georges, Noel Magloire Manga, Ba Ousseynou, Fenollar Florence, Sokhna Cheikh, Mediannikov Oleg.
Elephantiasis and *Mansonella perstans* in Senegal (West Africa)
3. **Diallo Manthia**, Bassene Hubert, Diatta Georges, Masse Sambou, Fenollar Florence, Sokhna Cheikh, Mediannikov Oleg.
Role of Tabanidae in the transmission of filarial (In progress)

II. Autre publication :

Pascal Dembélé, **Manthia Diallo**, Papa Mouhamadou Gaye, Mady Cissoko, Mahamadou A Thera and Stéphane Ranque*
Epidemiology of *Mansonella perstans* in Mali

III. Participation aux conférences et congrès au cours de la période de thèse :

Manthia DIALLO, (2022), Journée scientifique de la Fondation Méditerranée Infection et Infectiopôle Sud, Marseille, (présentation sous forme de poster)

Manthia DIALLO, (2023), 1er Congrès de la Société Sénégalaise de Parasitologie, Mycologie et Entomologie (SOSEPAME). (Présentation orale).

Manthia DIALLO, (2024), Journée scientifique de la Fondation Méditerranée Infection et Infectiopôle Sud, Marseille, (présentation sous forme de poster)

Résumé

Les filarioses sont des infections parasitaires courantes, causées par des vers nématodes filaires dans l'organisme. Parmi les filarioses humaines, on distingue la filariose lymphatique, causée par les espèces *Wuchereria bancrofti*, *Brugia timori* et *Brugia malayi* ; l'onchocercose, ou cécité des rivières, causée par *Onchocerca volvulus* ; la loase, provoquée par le parasite *Loa loa* et la mansonellose, due aux espèces *Mansonella perstans*, *Mansonella streptocerca* et *Mansonella ozzardi*. Au Sénégal, les programmes de lutte ont permis d'interrompre la transmission des parasites dans la majorité des districts sanitaires notamment pour l'onchocercose et la filariose lymphatique. Cependant, la filariose lymphatique reste encore endémique dans les zones rurales du pays.

Ce travail de thèse porte sur les filarioses humaines au Sénégal. Il se concentre principalement sur le dépistage et l'épidémiologie des filaires impliquées dans la transmission des maladies filariennes dans le sud du Sénégal. Notre étude a pour objectif principal d'explorer l'étiologie et l'épidémiologie des filarioses dans les zones étudiées, et/est scindé en deux parties :

La première partie présente une revue de la littérature sur les filarioses en Afrique de l'Ouest, abordant les pathologies existantes, leur épidémiologie, leur pathogénicité, les méthodes de diagnostic, le traitement et les différentes mesures de prévention ciblant les vecteurs arthropodes.

Dans la seconde partie, la technique de la PCR en temps réel (RT-PCR) et le séquençage ont été utilisés pour détecter *M. perstans* à partir d'échantillons de sang capillaire, de cassettes TDR *Plasmodium falciparum* négatifs et de moustiques culicoïdes vecteurs. *M. perstans* a été détecté chez 2 patients sur 23 atteints d'éléphantiasis collectés en 2021, chez 12 sur 279 échantillons de sang de sujets fébriles de Mlomp (2018-2020) et chez 11 sur 3714 cassettes TDR du paludisme négatives. Des moustiques (n=1487) ont été capturés avec des pièges lumineux CDC, OVI et à partir de la Faune Matinale Résiduelle. En plus, des spécimens de culicoïdes (n=991) ont été capturés uniquement avec les pièges lumineux. Après analyses moléculaires basées l'ADN, aucun spécimen de moustique et de culicoïdes n'a été retrouvé infecté par *M. perstans*.

Mots-clés : Filarioses, *Wuchereria bancrofti*, *Brugia timori*, *Brugia malayi*, *Onchocerca volvulus*, *Loa loa*, *Mansonella perstans*, *Mansonella streptocerca*, *Mansonella ozzardi*, Arthropodes.

Abstract

Filariasis is a common parasitic disease caused by the presence of filarial nematode worms in the body. Human filariasis includes pathologies: lymphatic filariasis, caused by *Wuchereria bancrofti*, *Brugia timori*, and *Brugia malayi*; onchocerciasis or river blindness, caused by *Onchocerca volvulus*; loiasis, caused by the parasite *Loa loa*; and mansonellosis, caused by the species *Mansonella perstans*, *Mansonella streptocerca*, and *Mansonella ozzardi*. In Senegal, control programs have successfully interrupted the transmission of the parasite in most health districts for onchocerciasis and lymphatic filariasis. However, lymphatic filariasis remains endemic in rural areas of Senegal.

This thesis work focuses on human filariasis in Senegal, particularly screening and epidemiology of filariasis involved in the transmission of human filarial diseases in southern Senegal. Our study aims to investigate the etiology and epidemiology of filariasis in the studied areas and is divided into two parts. The first part presents a literature review on filariasis in West Africa, addressing existing pathologies, their epidemiology, pathogenicity, diagnostic methods, treatment, and various prevention measures targeting arthropod vectors. In the second part, the real-time PCR (RT-PCR) technique and sequencing were used to detect *M. perstans* from capillary blood samples, *Plasmodium falciparum*-negative RDT cassettes, and vector *Culicoides* mosquitoes. *M. perstans* was detected in 2 out of 23 patients with elephantiasis collected in 2021, in 12 out of 279 blood samples from febrile subjects in Mlomp (2018-2020), and in 11 out of 3714 negative malaria RDT cassettes. Mosquitoes (n=1487) were captured using CDC light traps, OVI traps, and from the morning residual fauna. Additionally, *Culicoides* specimens (n=991) were captured exclusively with light traps. Following DNA-based molecular analyses, no mosquito or *Culicoides* specimens were found to be infected with *M. perstans*.

Keywords: Filariasis, *Wuchereria bancrofti*, *Brugia timori*, *Brugia malayi*, *Onchocerca volvulus*, *Loa loa*, *Mansonella perstans*, *Mansonella streptocerca*, *Mansonella ozzardi*, Arthropods.

REMERCIEMENTS

Je tiens tout d'abord à remercier les membres du jury d'avoir accepté d'évaluer le travail effectué au cours de cette thèse.

Je tiens à adresser ma plus grande reconnaissance à mon directeur de thèse **Dr Oleg MEDIANNIKOV**, pour m'avoir offert la chance de travailler sur ce sujet et me permettant d'accueillir de nombreuses connaissances. Je lui en suis infiniment reconnaissante. Je le remercie pour la confiance et le soutien qu'il m'a toujours témoignés ainsi que pour ses conseils avertis.

A **Dr Cheikh SOKHNA**, ce travail est le témoignage de la confiance que vous avez placé en nous. Nous avons été séduits par votre simplicité, votre amour pour le travail bien fait, votre souci constant de la bonne formation des futurs cadres. Nous vous serons toujours reconnaissants pour tous les services rendus. Par ailleurs, nous vous prions de bien vouloir accepter nos excuses pour toutes les fois où nous n'avons pas été à hauteur de souhait.

A notre présidente du jury **Pr Laurence CAMOIN**, vous nous faites un grand honneur en acceptant de présider ce travail malgré vos multiples occupations. Vos qualités exceptionnelles d'enseignant et de chercheur. Veuillez trouver ici cher maître, l'expression de notre profonde gratitude.

Pr Doudou SOW à l'Université Gaston Berger de Saint-Louis pour m'avoir bien voulu juger mon travail de thèse. En acceptant d'être rapporteur de mon jury de thèse, vous me faites un honneur. Je vous adresse ma profonde gratitude.

Pr Mama Cheikh SECK de l'Université Cheikh Anta Diop de Dakar (UCAD), qui nous a fait l'honneur de participer au jury et accepter d'évaluer ce travail.

Pr Christelle POMARES de l'Université Cote d'Azur, CHU Nice, pour avoir accepté d'évaluer mon travail. Veuillez trouver ici l'expression de ma profonde reconnaissance.

Mes remerciements touchent également **Pr Florence FENOLLAR**, pour l'intérêt qu'elle a porté à mon travail. Je vous remercie également pour votre aide scientifique, vos conseils incontournables durant ce travail et pour toujours avoir trouvé le temps disponible pour répondre à mes questions.

Mes remerciements vont à l'endroit du **Dr Hubert BASSENE**, **Dr Georges DIATTA**, **Dr GAYE Papa Mouhamed** et sa famille, **Dr Masse SAMBOU**, **Dr Alissa HAMMOUD**, **Dr Joa Braïthe Mangombi-PAMBOU**, **Dr Ibrahima NDIAYE**, **Adama DIARRA** pour leur aide si précieuse durant toute ma thèse.

Je remercie tous les étudiants, personnel de l'IHU, en particulier ceux de mon équipe: **Dame, Mapenda, Pricile, Clement, Rim, Cheikh, Charles, Keita, Ignestin, Abouacar, Karifa**. Je garde de bons souvenirs avec chacun d'entre vous.

Je remercie les ingénieurs, les techniciens, étudiants et tout le personnel MINES de Dakar pour les bons moments passés ensemble particulièrement **Dr DOUCOURE, Dr BA Charles BOUGANALI, Malick TOURE, Malick DIOP, Kokou**.

Je remercie les ingénieurs, les techniciens et tout le personnel de l'IHU particulièrement **Mme Micheline PITACCOLO, Mme Katie MZE, Mme Marine SANTORIELLO, Mme CORONA et Mme Monna** et ses collègues qui ont d'une manière ou d'une autre contribué à la réalisation de cette thèse.

DEDICACES

A ma très chère mère **Penda NDIAYE**,

Tu es pour nous un exemple de bonté, de courage et d'amour en tant que maman. Toujours prête à tout sacrifier pour que nous, tes enfants, devenions meilleurs. Tu es pour moi une mère attentionnée une amie et confidente. Tu n'as pas cessé un seul jour de penser à notre avenir durant tout ce parcours. Autant de phrases aussi expressives soient-elles ne sauraient montrer le degré d'amour et d'affection que j'éprouve pour toi. Une femme battante, affectueuse qui a toujours su m'écouter, me conseiller, m'épauler, et me soutenir dans les moments les plus difficiles. Ce travail est le fruit de ton affection. Tu représentes pour moi la femme exemplaire soucieuse de l'avenir de ses enfants. Tes prières, ton affection maternelle, ton soutien tant psychologique que matériel ne m'ont jamais fait défaut. Ce travail est le fruit de ton sacrifice consentis pour moi. Que Dieu nous permette de te témoigner longtemps notre attachement et te prête longue vie et une bonne santé afin que tu puisses continuer à guider mes pas !

A mon défunt Père : **Elhadji Salif DIALLO**,

Je dédie ce travail à la mémoire de mon cher père Elhadji Salif DIALLO. J'aurais tant aimé que tu sois présent papa. Aucune dédicace ne saurait exprimer mon respect, mon amour éternel et ma considération pour les sacrifices que tu as consenti pour mon instruction et mon bien-être. Je te remercie pour tout le soutien et l'amour que tu m'as apportés depuis mon enfance et j'espère que ta bénédiction m'accompagnera toujours. Priez pour que nous puissions en faire de même pour nos enfants. Que Dieu le tout puissant t'Accueille dans le plus grand paradis.

A mes frères et sœurs :

Abdourahmane, Abdoulaye, Elimane, Moussa, Tacko, Marième et Oulaye.

Vous êtes très sympathiques et travailleurs. En souvenir d'une belle enfance vécue ensemble et la complicité et l'entente qui nous unissent. En témoignage de mon amour, de mon admiration et de ma grande affection ; en reconnaissance de votre soutien et patience à mon égard durant ces longues années d'étude, je vous prie de trouver dans ce travail l'expression de mon estime et de mon sincère attachement. Je vous exhorte à persévérer dans la patience et le courage et prie Dieu le tout puissant pour qu'il vous donne la piété, la santé, le bonheur et la prospérité. Ce travail est le vôtre.

A la mémoire de ma grande sœur, que j'ai beaucoup aimé et qui était ma meilleur amie, ma confidente **Mané DIALLO**, son âme repose en paix.

A **Mr Badji Ibrahima**, son amour qui illumine ma vie, son soutien, ses conseils, ses encouragements et son aide quotidienne.

Tu m'as toujours encouragé et soutenu durant tout le cycle. Reçois ici mon sincère remerciement cher bien aimé.

A mes oncles et tantes

Je vous suis très reconnaissante. Que DIEU vous bénisse !

Aux membres de ma belle famille

Vous avez fait beaucoup de bénédictions. Trouvez ici l'expression de ma profonde gratitude. Que Dieu vous accorde longue vie dans le bonheur, la santé et la prospérité.

A tous mes amis et collègues,

je vous remercie de votre esprit de bonne collégialité ! Particulièrement à **Mme DIOUF Coumba, Mme SY Oumou, Mme BALDE Safietou** et Mme **Anna julienne**.

Trouvez ici toute ma reconnaissance, pour tous ces souvenirs et ces moments de bonheur vécus grâce à vous.

Table des matières

AFFIDAVIT	I
AFFIDAVIT	II
LISTE DES PUBLICATIONS ET PARTICIPATION AUX CONFERENCES ET CONGRES	III
RESUME	IV
DEDICACES	VIII
INTRODUCTION.....	1
PARTIE 1: REVUE DE LA LITTERATURE	6
ARTICLE 1	7
FILARIASIS IN WEST AFRICA.....	7
PARTIE 2	62
ARTICLE 2	63
ELEPHANTIASIS AND <i>MANSONELLA PERSTANS</i> IN SENEGAL (WEST AFRICA).....	63
ANNEXES	110
PUBLICATION EN COLLABORATION	110
ARTICLE 3	111
EPIDEMIOLOGY OF <i>MANSONELLA PERSTANS</i> IN MALI	111
(SOUMIS DANS PARASITOLOGY & VECTOR-BORNE DISEASES).....	111
EPIDEMIOLOGY OF <i>MANSONELLA PERSTANS</i> IN MALI	112
CONCLUSIONS ET PERSPECTIVES	149
REFERENCES BIBLIOGRAPHIQUES	149

1 INTRODUCTION

2 Les filarioses sont un groupe de maladies infectieuses causées par des nématodes appelés
3 filaires, transmises à l'Homme par l'intermédiaire d'insectes vecteurs (arthropodes
4 hématophages femelles : moucheron, moustique, simulie ou taon) (Walther et Muller 2003;
5 Mendoza et al. 2009). Les filarioses sont largement répandues dans le monde, touchant
6 principalement les populations pauvres vivant dans les pays en développement,
7 particulièrement dans les zones à climat tropical et subtropical. Environ 1 milliard de personnes
8 sont exposées, et plus de 200 millions de personnes sont actuellement atteintes par au moins
9 une forme de filariose (Desoubieux, Chesnay, et Chandenier 2020). Ces maladies sont
10 associées à un faible niveau de développement, à de mauvaises conditions d'hygiène et à
11 certaines conditions écologiques (Pérez-Molina *et al.* 2010). Bien que les filarioses aient des
12 manifestations cliniques, celles-ci n'apparaissent généralement qu'une fois que les parasites ont
13 atteint leur stade adulte, et dépendent des espèces ainsi que de la localisation des adultes et des
14 microfilaires (Bouchery et al. 2013). En outre, ces parasites peuvent entraîner des incapacités
15 sévères ou des lésions permanentes. Les filarioses ont un impact socio-économique majeur,
16 provoquant une forte morbidité et des maladies graves qui entraînent une stigmatisation sociale,
17 une marginalisation et une perte d'emploi pour les personnes affectées (Lustigman, Grote, et
18 Ghedin 2017).

19 Plusieurs espèces de filaires sont connues pour leur intérêt médical, notamment
20 *Onchocerca volvulus*, *Loa loa*, *Wuchereria bancrofti*, *Brugia malayi*, *Brugia timori*,
21 *Mansonella perstans*, *Mansonella streptocerca* et *Mansonella ozzardi*. Les vers adultes ou
22 macrofilaires produisent des embryons ou microfilaires qui circulent dans le derme (pour *O.*
23 *volvulus* et *M. streptocerca*) ou dans le sang (M. Boussinesq 2007). Ces microfilaires sont
24 récupérées par les insectes vecteurs lors de leurs repas sanguins et se développent en larves L3
25 après plusieurs mues. Les larves L3 sont ensuite transmises à un autre individu lors des repas
26 sanguins suivants (Walther et Muller 2003; Alhassan *et al.* 2015). Les adultes peuvent se
27 localiser à différents endroits du corps humain en fonction de l'espèce impliquée. Dans
28 certaines régions d'Afrique, la filariose lymphatique, l'onchocercose, la mansonellose à *M.*
29 *perstans* et/ou la loase sont co-endémiques, et certains patients peuvent présenter des infections
30 multiples. Les filaires chez l'homme sont classées selon leur localisation dans le corps, leur
31 mode de contamination, leurs manifestations cliniques et leur répartition géographique. La
32 **filariose lymphatique (FL)**, communément appelée **Eléphantiasis**, fait partie des maladies
33 tropicales négligées (MTN) les plus anciennes. C'est l'une des maladies les plus dévastatrices et

défigurantes chez l'Homme avec des conséquences socio-économiques énormes matérialisées par la stigmatisation sociale, la baisse ou la perte totale de productivité et le coût élevé des traitements (Devi et Raju 2018). Selon l'OMS, 51,4 millions de personnes sont infectées (World Health Organization 2022) dans le monde. La FL est principalement causée par trois espèces de nématodes appartenant à la famille des *Filarioidea*, dont *Wuchereria bancrofti* (Cobbold, 1887), responsable de 90 % des cas dans le monde (Ottesen 2006; Katiyar et K. Singh 2011), principalement dans les régions tropicales et subtropicales et dans le Pacifique. Les deux autres espèces, *Brugia malayi* (Brug, 1927) et *Brugia timori* (Partono, 1977), sont endémiques en Asie. En Afrique, où la maladie est entièrement causée par *W. bancrofti*, plus de 45 millions de personnes sont touchées (Hotez et Kamath 2009). Au Sénégal, la FL est endémique avec une prévalence globale de 4,7 % chez les personnes âgées de plus de 45 ans (Faye et al. 2009). Bien que l'espèce *W. bancrofti* ait été morphologiquement décrite au Sénégal, elle n'a jamais été confirmée sur la base de l'ADN. L'infection survient lorsque les parasites filaires sont transmis à l'homme par des moustiques des genres *Culex*, *Aedes*, *Anopheles* et *Mansonia* (Manguin et al. 2010; Diwakar et al. 2018). La FL présente des symptômes chroniques et visibles, tels que l'hydrocèle, un lymphœdème secondaire des membres inférieurs (Ramaiah et Ottesen 2014), ou une éosinophilie pulmonaire tropicale qui ressemble à l'asthme (Tahita et al. 2024).

L'**onchocercose** est une filariose cutanée causée par le nématode *Onchocerca volvulus* (Leuckart, 1893), qui se développe dans le tissu sous-cutané de l'Homme. Ce parasite est transmis par la piqure de la femelle du genre *Simulium*. C'est une maladie invalidante, un obstacle majeur à l'établissement de la population. L'infection par *O. volvulus* est responsable de manifestations cutanées et oculaires graves (Adewole et Ayeni 2009; Katawa et al. 2015; Njim, Ngum, et Aminde 2015), et est également associée à l'épilepsie et au syndrome du hochement de tête (Colebunders et al. 2018; Vinkeles Melchers et al. 2018). C'est une maladie liée à la pauvreté et à la productivité des ménages (Ibe et al. 2015; Opoku et al. 2018) et constitue un réel problème de santé publique. L'onchocercose touche environ 21 millions de personnes, dont 99 % des cas signalés dans 31 pays d'Afrique subsaharienne (World Health Organization (WHO) 2020). En 2019, une évaluation entomologique réalisée dans les 8 districts concernés a révélé un taux d'infectivité des simules inférieur à 0,05 %. Ce résultat confirme l'interruption de la transmission de l'onchocercose dans ces zones. Ces données témoignent des avancées significatives dans le contrôle de la maladie (World Health Organization (WHO) 2022).

L'infestation par ***Loa loa*** (Cobbold, 1894) est une maladie parasitaire connue sous le nom de **ver oculaire africain**. La loase est endémique dans les zones tropicales d'Afrique de l'Ouest et d'Afrique centrale, notamment au Cameroun, au Congo, au Gabon et au Nigéria (Zouré et al. 2011).

Bien que la loase ne soit pas endémique au Sénégal, un cas a été détecté, avec la présence du ver adulte en migration dans l'œil, chez un patient originaire de la République centrafricaine (Diagne et al. 2018). Le filaire *Loa loa* est transmise à l'Homme par des tabanidés du genre *Chrysops* (Anderson 2000). La loase est bien souvent considérée comme bénigne et dans la plupart des cas asymptomatique, mais peut présenter de nombreux aspects cliniques (A. Aubry et al. 2022). Les manifestations cliniques les plus frappantes sont la migration du ver adulte sous la conjonctive bulbaire et les œdèmes migratoires transitoires, appelés tuméfaction de Calabar (Gardon et al. 1997; A. Aubry et al. 2022).

Bien que négligée, la **mansonellose** est une filariose humaine bénigne, mais reste une infection filarienne négligée malgré le fait que 600 millions de personnes sont exposées à un risque élevé en Afrique et dans certaines parties de l'Amérique centrale et du Sud (Simonsen, Onapa, et Asio 2011; Ta-Tang et al. 2021). À ce jour, la mansonellose est causée par trois espèces de nématodes filaires : *Mansonella perstans* (Manson, 1891), *M. ozzardi* (Manson, 1897) et *M. streptocerca* (Macfie & Corson, 1922), qui ne provoquent que des symptômes bénins et généralement non spécifiques (Medeiros et al. 2017; Ta-Tang et al. 2018; Ehrens, Hoerauf, et Hübner 2022). Récemment, des études génétiques ont montré la présence d'une autre variante génétique, *Mansonella sp. Deux*, infectant les humains en Afrique, notamment au Gabon (Mourembou et al. 2015; Sandri et al. 2021; Rodi et al. 2023). *M. perstans* est la plus fréquente en Afrique, avec une prévalence souvent élevée dans les zones endémiques. Elle infecte environ 114 millions de personnes en Afrique, principalement dans 33 pays d'Afrique subsaharienne. Cette espèce est responsable de l'infection au Sénégal, où la prévalence est de 14,5 % dans la région rurale de Kédougou (Bassene et al. 2015).

Des études antérieures ont montré qu'il existe de nombreuses méthodes de diagnostic pour détecter les filarioses, utilisées en première intention en cas de suspicion clinique. Ces recherches peuvent être effectuées :

- **Dans le sang** : par observation à l'état frais, par frottis mince ou goutte épaisse, ou à l'aide de techniques de concentration comme la leucoconcentration, la technique de Knott ou la filtration membranaire.

- **Dans une biopsie cutanée exsangue (BCE)** : soit directement à l'état frais, soit après coloration ou par examen histologique.

En outre, des techniques telles que la précipitation (électrosynérèse, coélectrosynérèse, immunoélectrophorèse), l'immunofluorescence indirecte et des techniques immunoenzymatiques de type ELISA sont employées pour rechercher les anticorps et antigènes circulants. Ces méthodes s'appliquent aux diagnostics des filarioses lymphatiques, de la loase, des mansonelloses (*M. perstans*, *M. ozzardi*) et de l'onchocercose (*O. volvulus*, *M. streptocerca*). Au Sénégal, la réaction en chaîne par polymérase (PCR), qui est une méthode moléculaire très spécifique et sensible, est également utilisée pour détecter ces filaires.

Selon l'OMS, plusieurs stratégies de contrôle ont été mises en place pour stopper la propagation de ces maladies. Ces stratégies reposent principalement sur la **chimiothérapie préventive** par l'administration à grande échelle de médicaments (Mass Drug Administration, MDA). Ces médicaments agissent soit sur les microfilaires, soit sur les adultes. Classiquement, les principaux antifilaricides disponibles sont la **diéthylcarbamazine (DEC)**, l'**ivermectine (IVM)** et l'**albendazole (ABL)**, utilisés seuls ou en combinaison. Le choix du traitement dépend de la confirmation du type de filaire.

Le Sénégal est endémique à l'onchocercose et à la FL. Cependant, après plusieurs campagnes de traitement de masse avec l'IVM, l'onchocercose a été totalement éliminée et la prévalence de la FL a été réduite de 95 % dans plusieurs districts sanitaires endémiques (World Health Organization (WHO). 2022). À noter que *M. perstans* montre une certaine résistance, voire une insensibilité, à certains médicaments filaricides, notamment la DEC et l'IVM (S. M. Asio, Simonsen, et Onapa 2009; Chelsea et William 2022).

Ce travail vise à étudier l'étiologie et l'épidémiologie des filarioses au Sénégal, en mettant particulièrement l'accent sur les nématodes parasites et leurs vecteurs dans le sud du pays. L'étude est scindée en plusieurs parties :

1. **Revue de la littérature** : Analyse des filarioses en Afrique de l'Ouest, incluant leurs généralités, leur épidémiologie, leur pathogénicité, leurs méthodes de diagnostic et les moyens de lutte employés dans les programmes de contrôle.
2. **Étiologie et dépistage** : Cette partie porte sur la détermination des causes des cas d'éléphantiasis, l'évaluation de l'utilité des bandelettes nitrocellulose des tests diagnostiques rapides (TDR) pour le dépistage de la filariose, et l'identification des

moustiques et culicoïdes comme vecteurs potentiels de *M. perstans* en milieu rural au Sénégal.

Nous avons ainsi réalisé la détection des parasites filaires dans des prélèvements sanguins de patients, à partir de TDR négatifs pour le paludisme, ainsi que sur des spécimens de moustiques et de culicoïdes. Pour cela, des analyses basées sur l'ADN, notamment la **PCR en temps réel (RT-PCR)** et le **séquençage Sanger**, ont été utilisées.

Ces filaires, bien que peu nombreuses parmi les espèces parasitant l'homme, représentent une diversité importante et jouent un rôle crucial dans les efforts de contrôle des maladies infectieuses.

160
161
162
163
164
165
166
167

168

169

170
171
172
173
174
175
176
177
178
179
180
181
182
183

PARTIE 1: REVUE DE LA LITTERATURE

Filariasis in West Africa

184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208

ARTICLE 1

Filariasis in West Africa

Manthia Diallo, Hubert Bassene, Goerges Diatta, Florence Fenollar, Cheikh Sokhna, Oleg
Mediannikov

209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233

Filariasis in West Africa

Manthia DIALLO^{1,3,5}, Hubert BASSENE², Cheikh SOKHNA^{2,3,4}, Georges DIATTA²,

Florence FENOLLAR^{3,4} Oleg MEDIANNIKOV^{1,3,6,*}

¹ MEPHI, Aix Marseille Univ, AP-HM, Marseille, France

² Aix Marseille Univ, IRD, MINES, Campus International IRD-UCAD, Dakar, Sénégal

³ IHU-Méditerranéen Infection, Marseille, France

⁴Aix Marseille Univ, SSA, RITMES, Marseille, France

⁵Department of Animal Biology, Faculty of Sciences and Techniques, Cheikh Anta Diop University
of Dakar, Senegal

⁶ IRD, Marseille, France

*Corresponding author: Oleg MEDIANNIKOV olegusss1@gmail.com

234

235 **Résumé**

236 Les maladies parasitaires liées à la filariose sont très fréquentes à travers le monde. Elles
237 sont provoquées par des nématodes, ou filaires, appartenant à la famille des Filaroidae. En
238 Afrique, la filariose se manifeste avec des tableaux cliniques variés, en fonction de l'espèce
239 parasitaire impliquée. Ces maladies constituent un problème de santé publique majeur,
240 entraînant des pertes socio-économiques significatives dans de nombreux pays tropicaux et
241 subtropicaux. En Afrique de l'Ouest, la filariose occupe une place prépondérante en termes de
242 santé publique. Selon leur impact sanitaire, les espèces filariennes sont classées en filarioses
243 pathogènes (filariose lymphatique, onchocercose, loase) et filarioses non pathogènes
244 (mansonellose). Toutes ces espèces partagent un cycle de vie similaire, nécessitant un hôte
245 intermédiaire indispensable à leur développement.

246 Cette revue se concentre sur les principales filarioses présentes en Afrique de l'Ouest,
247 notamment leur épidémiologie, leur pathogénicité, les méthodes diagnostiques employées, les
248 stratégies de prévention, ainsi que les traitements recommandés par l'OMS. La capacité à
249 détecter les parasites chez l'hôte est cruciale pour mettre en œuvre des stratégies de contrôle
250 efficaces. Cette revue examine les différentes méthodes de détection et de prévention des
251 infections filariennes, incluant l'utilisation de médicaments antifilariens efficaces ou
252 partiellement efficaces, qui agissent à différents stades du cycle parasitaire. Face au défi que
253 représente l'absence de médicaments à la fois sûrs et performants, il est essentiel d'envisager
254 des mesures approfondies et durables, permettant d'optimiser l'efficacité des traitements
255 antifilariens, utilisés seuls ou en association.

256 **Mots-clés** : Filariose, Filariose lymphatique, Onchocercose, Loase, Mansonellose,
257 Pathogénicité, Diagnostic, Prévention, Traitement, Afrique de l'Ouest.

258

259

260

261

262

263

Abstract

Parasitic diseases caused by filariasis are widespread across the globe. They result from infections by nematodes, or filariae, belonging to the Filaroidae family. In Africa, filariasis exhibits varied clinical presentations depending on the species of parasite involved. These diseases represent a major public health issue, leading to significant socio-economic losses in many tropical and subtropical countries. In West Africa, filariasis holds significant importance to public health. Based on their impact on public health, filarial species are classified into pathogenic filariasis (lymphatic filariasis, onchocerciasis, loiasis) and non-pathogenic filariasis (mansonellosis). These species share a similar life cycle, with an obligatory intermediate host playing a key role in their development.

This review focuses on the main filarial diseases present in West Africa, including their epidemiology, pathogenicity, diagnostic methods, prevention strategies, and treatments recommended by the WHO. The ability to detect parasites in the host is essential for implementing effective control measures. This review explores various methods of parasite detection and infection prevention, including the use of antifilarial drugs, both effective and less effective, which act at different stages of the parasite's life cycle. Given the challenge posed by the lack of safe and highly effective drugs, it is crucial to implement deeper and more sustainable measures, allowing antifilarial treatments to be optimized, either used individually or in combination, to enhance their efficacy.

Keywords: Filariasis, Lymphatic Filariasis, Onchocerciasis, Loiasis, Mansonellosis, Pathogenicity, Diagnosis, Prevention, Treatment, West Africa.

294

295 I. Generality of filariasis

296 Filariasis is a tropical parasitic disease prevalent worldwide. It is caused by nematodes
297 called filariae (roundworms), which belong to the Filarioidea families. These pathologies
298 present varied clinical manifestations depending on the species involved and are endemic in
299 several countries, particularly in West Africa. Filariasis is classified according to the location
300 of the macrofilariae (adult worms) in the body. These diseases show a great diversity of filariae,
301 although very few species parasitize humans. The infections can be categorized as lymphatic
302 (lymphatic filariasis), cutaneous or dermal (onchocerciasis and loiasis), or serous
303 (mansonellosis), depending on the tropism of the viviparous nematodes that cause them and
304 their location in the human body. The infections can be categorized as lymphatic (lymphatic
305 filariasis), cutaneous or dermal (onchocerciasis and loiasis), or serous (mansonellosis),
306 depending on the tropism of the viviparous nematodes that cause them and their location in the
307 human body (Bhalla, Dumas, et Preux 2013).

308 Transmission to humans occurs via the bite of infected mosquitoes, midges, blackflies,
309 or horseflies, which act as intermediate hosts for the infective larvae. Humans serve as the
310 definitive host. Filariasis constitutes a global public health problem, due to its impact on
311 population health and the enormous burden it imposes on endemic countries. Worldwide, more
312 than 200 million people are affected (Kwarteng, Ahuno, et Akoto 2016), and almost a billion
313 live in areas at risk of filariasis (Risch et al. 2021). Worldwide, more than 200 million people
314 are affected (Kwarteng, Ahuno, et Akoto 2016), and almost a billion live in areas at risk of
315 filariasis (Risch et al. 2021). These diseases are defined as conditions caused by invasive or
316 filarial tissue nematodes. The adult worms typically reside under the skin, in lymphatic vessels,
317 or in lymph nodes, while their microfilaria circulates in the blood, the dermis, or the lymphatic
318 pathways, depending on the species of filaria. They also result in enormous economic losses in
319 many African countries. Although many species of filariae exist, only a few infect humans.
320 These are divided into major filariasis (pathogenic) and minor filariasis (little or no pathogen)
321 (Table 1). All these species use humans as definitive hosts and are distributed globally,
322 particularly in West Africa, depending on their preferred location in the body.

323 Pathogenic filariasis includes lymphatic filariasis caused by *Wuchereria bancrofti*,
324 *Brugia malayi* and *Brugia timori*, cutaneous filariasis: onchocerciasis caused by *Onchocerca*
325 *volvulus*, loiasis caused by *Loa loa* and serous filariasis caused by *Mansonella perstans*,
326 *Mansonella streptocerca* and *Mansonella ozzardi* (Gentilini, Pinon, et Danis 1973).

327 They all have comparable reproductive cycles and parasitological characteristics (Frédéric
328 2018) according to their geographical distribution, modes of transmission, and clinical
329 manifestations. They all have comparable reproductive cycles and parasitological
330 characteristics (Frédéric 2018) according to their geographical distribution, modes of
331 transmission, and clinical manifestations.

332 These filarial species have a significant social and economic impact on affected populations.
333 They cause serious morbidity, leading to social stigma, exclusion from society, and loss of
334 employment for those affected, resulting in a heavy disease burden. These filarial species have
335 a significant social and economic impact on affected populations. They cause serious morbidity,
336 leading to social stigma, exclusion from society, and loss of employment for those affected,
337 resulting in a heavy disease burden (Lustigman, Grote, et Ghedin 2017).

Table 1. Characteristics of filarial species.

	Species	Locations: Adults / microfilariae	Vectors	Clinical manifestations	Geographical distribution	References
Highly pathogenic filariae	<i>Wuchereri bancroftii</i>	Lymphatic / Blood (Nocturnal)	<i>Anopheles</i> sp, <i>Aedes</i> sp, <i>Culex</i> sp	Lymphedema, hydrocele, elephantiasis	Tropical and intertropical regions	(Taylor, Hoerauf, et Bockarie 2010; Nutman 2013; World Health Organization (WHO) 2023)
	<i>Brugia timori</i>	Lymphatic / Aperiodic blood	<i>Mansonia</i> sp		Timor	
	<i>Brugia malawi</i>				Malaysia	
	<i>Onchocerca volvulus</i>	Cutano-dermal / Dermal Suc	<i>Simulie</i>	Lesions (lymphatic-ocular- cutaneous), pruritus, nodules	Sub-Saharan Africa, Central and South Africa, Yemen	(Coffeng et al. 2014)
	<i>Loa loa</i>	Cutano-dermal / Blood (Diurnal)	<i>Chrysops</i>	Squid edema and adult worm under conjunctiva	West Equatorial Africa	(Zouré et al. 2011)
Filariae with low pathogenicit y	<i>Mansonella perstans</i>	Serous / Aperiodic blood	<i>Culicoides</i> sp	Pruritus, headache, arthralgia, fever, oedema, rash, keratitis, lymphadenopathy, pulmonary symptoms	Sub-Saharan Africa	(Simonsen, Onapa, et Asio 2011a; Enrico Rino Bregani et al. 2006)
	<i>Mansonella streptocerca</i>	Subcutaneous / dermal			Sub-Saharan Africa	
	<i>Mansonella ozzardi</i>	Subcutaneous / Aperiodic blood			America, Caribbean	
	<i>Dirofilaria immitis</i>	Pulmonary artery and sometimes in the right ventricle of the heart	<i>Anopheles</i> sp, <i>Aedes</i> sp, <i>Culex</i> sp	Benign asymptomatic pulmonary nodules	It is more widespread in tropical and temperate regions, particularly in the south-eastern United States,	(Alsarraf et al. 2021; Simón et al. 2012; Pampiglione, Canestri Trotti, et Rivasi 1995)

		(Immature adult) / Blood			many South American countries, Asia, Australia and southern Europe.
	<i>Dirofilaria repens</i> (Subcutaneous and ocular heartworm disease)	subcutaneous connective tissue (Adult and pre-adult worms) / Blood		<ul style="list-style-type: none"> - Affects different ocular regions (orbital area, eyelids and subconnective and intravitreal tissues). - Severe cases, with vision disturbances and complications such as retinal detachment, glaucoma, vitreous humor opacity, lens. 	The Old World, with areas of high prevalence in Southern and Eastern Europe, Asia Minor, Central Asia and Sri Lanka.

II. Epidemiology of filariasis in West Africa

Known filarial infections that parasitize humans include *W. bancrofti*, *Brugia* sp., *O. volvulus*, *L. loa*, *M. perstans*, *M. streptocerca*, and *M. ozzardi*. In this review, we focus only on filarial species endemic to West African countries (Figure 1).




Figure 1: Countries endemic for lymphatic filariasis, onchocerciasis, loiasis, and mansonellosis. (Senegal is not endemic to loiasis)

1. Lymphatic filariasis

A. Description

Lymphatic filariasis (LF), also called elephantiasis, continues to be a major cause of morbidity and permanent disability in poor, rural, and marginalized populations, as well as in endemic areas (Hotez et al. 2007; Davis et al. 2019). It is caused by parasitic of the genera *Wuchereria* and *Brugia* sp transmitted by mosquitoes (Yaya I. Coulibaly et al., 2016; Dewi et al., 2015). This disease encompasses pathological manifestations resulting from parasitism of the human body by filariae. More than 90% of infections worldwide are caused by the filarial species *W. bancrofti*, which is widely present in tropical and subtropical zones in Africa

(Manguin et al. 2010; Small, Tisch, et Zimmerman 2014), Asia Pacific, and America. *B. malayi* accounts for most of the remaining cases (Taylor, Hoerauf, et Bockarie 2010; Bockarie et Deb 2010). The visible, painful, and disfiguring manifestations of the disease lymphedema (elephantiasis) and hydrocele negatively impact the lives and socio-economic well-being of those affected, as well as their families (Ouedraogo et al. 2016). These symptoms often appear later in life. Lymphatic filariasis also affects the lymphatic system, leading to psychosocial difficulties such as social disgrace, reduced marriage prospects, and sometimes the dislocation of couples or families (Wynd 2007; Molyneux et Taylor 2001). Many individuals feel socially excluded in their own communities due to the stigma associated with the disease (Coreil et Mayard 2009). Infection occurs when filariae responsible for the disease are transmitted to humans through the bites of blood-sucking female mosquitoes belonging to the genera *Aedes*, *Anopheles*, *Mansonia*, and *Culex* (World Health Organization (WHO) 2021) (This disease encompasses pathological manifestations resulting from parasitism of the human body by filariae. More than 90% of infections worldwide are caused by the filarial species *W. bancrofti*, which is widely present in tropical and subtropical zones in Africa (Manguin et al. 2010; Small, Tisch, et Zimmerman 2014), Asia Pacific, and America. *B. malayi* accounts for most of the remaining cases (Taylor, Hoerauf, et Bockarie 2010; Bockarie et Deb 2010). The visible, painful, and disfiguring manifestations of the disease lymphedema (elephantiasis) and hydrocele negatively impact the lives and socio-economic well-being of those affected, as well as their families (Ouedraogo et al. 2016). These symptoms often appear later in life. Lymphatic filariasis also affects the lymphatic system, leading to psychosocial difficulties such as social disgrace, reduced marriage prospects, and sometimes the dislocation of couples or families (Wynd 2007; Molyneux et Taylor 2001). Many individuals feel socially excluded in their own communities due to the stigma associated with the disease (Coreil et Mayard 2009). Infection occurs when filariae responsible for the disease are transmitted to humans through the bites of blood-sucking female mosquitoes belonging to the genera *Aedes*, *Anopheles*, *Mansonia*, and *Culex* (World Health Organization (WHO) 2021) (<https://iris.who.int/handle/10665/346577>). These mosquitoes are intermediate hosts of the parasite, while humans are the definitive host. which are the intermediate hosts of the parasite while humans are the definitive host.

B. Epidemiology

Lymphatic filariasis is one of the neglected tropical diseases (NTDs) and is recognized as the second most debilitating mosquito-borne disease after malaria (Manguin et al. 2010;

Bockarie et Deb 2010; Shenoy 2008). It is particularly impactful due to the physical disabilities and socio-economic consequences it causes. Lymphatic filariasis is one of the neglected tropical diseases (NTDs) and is recognized as the second most debilitating mosquito-borne disease after malaria (Manguin et al. 2010; Bockarie et Deb 2010; Shenoy 2008). It is particularly impactful due to the physical disabilities and socio-economic consequences it causes. In 2018, 893 million people in 49 countries were at risk of lymphatic filariasis and required preventive chemotherapy to halt the spread of infection (Nizeyimana 2021). By 2021, 51.4 million people were infected in 72 endemic countries (P. P. Aubry et Docteur Bernard-Alex Gaüzère 2023), with 36 million developing chronic manifestations such as hydrocele or secondary lymphedema (Ramaiah et Ottesen 2014; Cromwell et al. 2020). In 2018, 893 million people in 49 countries were at risk of lymphatic filariasis and required preventive chemotherapy to halt the spread of infection (Nizeyimana 2021). By 2021, 51.4 million people were infected in 72 endemic countries (P. P. Aubry et Docteur Bernard-Alex Gaüzère 2023), with 36 million developing chronic manifestations such as hydrocele or secondary lymphedema (Ramaiah et Ottesen 2014; Cromwell et al. 2020).

Globally, initial estimates for lymphatic filariasis indicated that 25 million men were affected by hydrocele, while more than 15 million people suffered from debilitating lymphedema (Adhikari et al. 2015). Globally, initial estimates for lymphatic filariasis indicated that 25 million men were affected by hydrocele, while more than 15 million people suffered from debilitating lymphedema (Adhikari et al. 2015). In Senegal, the presence of lymphatic filariasis has been known for decades. Early survey studies conducted in the Mbour department on the southern coast identified a focus of the disease (Diallo et al. 1977). However, since then, few studies have been conducted, and filarial infection remains a low national priority due to the prominence of diseases like malaria and bilharzia. The latest study in Senegal showed an overall prevalence of 4.7% among people aged over 45 years, with significant regional variations (Faye et al. 2009). Lymphatic filariasis is also endemic in other West African countries, such as Mali, particularly in the Sikasso district, where *W. bancrofti* is transmitted primarily by *Anopheles gambiae sensu lato* and *Anopheles funestus* (Brenques et al. 1968; Yaya Ibrahim Coulibaly et al. 2013). In a study of two neighboring villages in Mali, located in a *W. bancrofti* endemic area, the overall prevalence of circulating antigen was greater than 50% among 373 participants aged 14 to 65 years (Dolo et al. 2018). Lymphatic filariasis affects children as young as 6 years old, with an antigenemia prevalence of 0.24% detected using the FTS test for the parasite *W. bancrofti* (Kima et al. 2021). In Niger, low levels of *W. bancrofti*

transmission have been observed in previously non-endemic areas. Among participants aged 5 to 105 years, 29 individuals (0.7%) tested positive using the filariasis test strip (FTS) (Badia-Rius et al. 2023). In Côte d'Ivoire, pre-mass drug administration (MDA) surveys showed a prevalence of *W. bancrofti* infection ranging from 3.38% to 5.50% in border health districts, including Aboisso, Bloléquin, Odienné, and Ouangolodougou, based on FTS (Yokoly et al. 2020).

2. Onchocerciasis

A. Description

Onchocerciasis is an endemic disease caused by a nematode called *O. volvulus* (O'Hanlon et al. 2016; Vinkeles Melchers et al. 2021; Crawford et al. 2024), for which humans are the only reservoir. The vector breeds in fast-flowing, oxygen-rich rivers in affected areas, with disease transmission and prevalence typically extending along these river basins, hence the name river blindness (Herricks et al. 2017). It is a serious disease with slow progression, caused by the accumulation of filariae in the human body. These parasites are transmitted by a small black fly, the intermediate host, which belongs to the genus *Simulium*. The Damnosum complex is most often responsible for onchocerciasis in Africa. Vector species are mainly present in humid areas and reproduce in fast-flowing waterways (Robert et al. 2022).

B. Epidemiology

O. volvulus causes significant skin and eye diseases. The vector is a widely distributed dipteran of the genus *Simulium*, a black fly measuring 1–3 mm with a painful bite, and is most active in Africa (Brattig, Cheke, et Garms 2021). Black flies lay their eggs on plants or rocks in cool, fast-flowing, nutrient-rich, and highly oxygenated waters, such as waterfalls and rapids. The location of larval breeding grounds explains why the distribution of onchocerciasis is closely tied to hydrographic networks. Black flies typically have a lifespan of no more than one month. In 2017, onchocerciasis affected 20.9 million people worldwide, including 14.6 million cases of dermatological manifestations and 1.15 million cases of vision loss (James et al. 2018). In West Africa, female black flies live up to 9 days after their first blood meal, with 9% in forest areas and 47% in savannah areas (M. Boussinesq 2003). Approximately 160 million people are at risk of onchocerciasis, spread across 31 African countries (Porter et Hopkins 2015). In Burkina Faso, the disease decreased by 15% in 2002 after the implementation of the Onchocerciasis Control Program in West Africa. However, in 2011, a resurgence of the disease was observed, with prevalence ranging from 0.7% to 71%

(Sawadogo 2019). In Nigeria, before ivermectin treatment, prevalence ranged between 23.1% and 84.9%, with a median prevalence of 52.0%. After 15 to 17 years of treatment, prevalence dropped to 0% in all communities (Tekle et al., 2012). However, *O. volvulus* infections remained at 15.2% in skin samples (Ojurongbe et al., 2015). According to Herricks, Nigeria bears one of the highest burdens of onchocerciasis worldwide (Herricks et al. 2017). A recent study in Togo identified ocular diseases in children and young adults in villages, including papillitis, cataracts, chorioretinitis, conjunctivitis, and tropical limbo-conjunctivitis (LCET). Other findings included iridocyclitis, sclerosing keratitis, and blindness in one eye, despite ongoing MDA programs that reduced microfilariae levels (Komlan et al. 2018). Currently, 99% of cases are found in 31 sub-Saharan African countries (World Health Organization (WHO) 2020). In the villages along the Gambia River in Senegal, a study conducted by De Sole et al. in 1991 revealed high microfilarial loads, exceeding 10 microfilariae per skin (mf/s) among adults, indicating intense onchocerciasis transmission (De Sole et al. 1991). However, due to targeted control efforts and interventions, an entomological evaluation conducted in 2019 across the eight affected districts showed a blackfly infectivity rate below 0.05%. This result confirms the interruption of disease transmission in these areas, highlighting significant progress in the fight against onchocerciasis (World Health Organization, 2022).

3. Mansonellosis

A. Description

Mansonellosis is a tropical disease, often considered even more neglected than Neglected Tropical Diseases (NTDs). Although it is arguably the most common filarial infection worldwide, it is the least studied and classified as a neglected parasitic infection (Lima et al. 2016; Ta-Tang et al. 2018). Human infection is caused by filarial nematodes of the genus *Mansonella*, including *M. perstans*, *M. streptocerca* and *M. ozzardi* (Tatuene et al. 2014a). These species are widespread in West Africa and other parts of Africa (Ta-Tang et al. 2018). Outbreaks of *M. perstans* and *M. streptocerca* are limited and occur mainly in sub-Saharan Africa, with sporadic cases in South America (Tavares da Silva et al. 2017; Ta-Tang et al. 2018). *M. ozzardi*, on the other hand, is found exclusively in the western hemisphere, from southern Mexico to north-western Argentina (Lima et al. 2016; Ta-Tang et al. 2018). *M. streptocerca* is primarily a human parasite but has been reported to infect wild chimpanzees in rare cases. A fourth species, *Mansonella* sp. 'TWO', has recently been described from Gabon, although its clinical relevance remains uncertain (Rodi et al. 2023a). Biting midges of the genus

Culicoides are the main vectors of *M. perstans* and *M. streptocerca*. For *M. ozzardi*, transmission is mainly by black flies (Simulium) (Gaillard et al. 2020; Ta-Tang et al. 2021b). After a blood meal, intermediate hosts ingest microfilariae, which develop into adult worms residing in pleural or peritoneal cavities or in the subcutaneous dermal layer (Medeiros et al. 2017; Ferreira, Crainey, et Luz 2021). Adult Mansonella females are viviparous and release *M. perstans* and *M. ozzardi* microfilariae into the peripheral blood, while *M. streptocerca* larvae are found in the skin (Ta-Tang et al. 2021b; Medeiros et al. 2017).

B. Epidemiology

Mansonellosis affects an estimated 114 million people across Africa, mainly in 33 countries, (B. Downes et Jacobsen 2010; Simonsen, Onapa, et Asio 2011). Prevalence is often high in endemic areas, even among children (Simonsen, Onapa, et Asio 2011a; Bassene et al. 2015). In Senegal, the prevalence was reported at 14.5% in febrile and afebrile children in the Kedougou region. *M. perstans* infections were more frequent among individuals aged over 29 years (33.3%) (Bassene et al. 2015a). In Ghana, a 2014 study reported a high burden of *M. perstans* among individuals co-infected with Buruli ulcers, with a prevalence of 23% in affected patients versus 13% in controls (Phillips et al. 2014). The overall prevalence in the studied communities was 32%, with variation between 2% and 75%, depending on vector species (L. B. Debrah et al. 2017). In Nigeria, among 755 nomadic Fulani, 8.7% were infected with *M. perstans*. The infection rate was higher in men (9.7%) than in women (6.5%) (Anosike et al. 2005). In addition, the prevalence of *M. perstans* increases with age in volunteers aged between 18 and 65. *M. perstans* is endemic in the two districts of Kolokani and Kolondieba in Mali, with prevalences of 11.79%, 4.35% and 9.50% respectively in the villages of Tieneguebougou, Bougoudiana and Boundioba (Sangare et al. 2018)

4. Loiasis

A. Description

Loiasis caused by *Loa loa* and commonly known as the African eye worm, mainly affects the forested areas of Central Africa (Jacobsen et al. 2022; Zouré et al. 2011). The disease is transmitted to humans by the bites of tabanid flies of the genus *Chrysops*, in particular *Chrysops silacea* and *Chrysops dimidiata*, with humans being the only known reservoir (Knopp et al. 2012; Kelly-Hope et al. 2017). In most cases, loasis is asymptomatic or presents as a mild infection (Adam et al. 2020; A. Aubry et al. 2022). Parasites develop in the lungs at night to escape clearance by the spleen (Ramharter et al. 2023). The adults migrate to the subcutaneous

tissues, passing under the skin and the transparent mucous membrane of the eyes, and are visible in the subconjunctival space. *L. loa* parasites are round, whitish worms measuring 3 to 7 cm, but can exceed 10 cm (Mandal et al. 2012; Jacobsen et al. 2022). Transmission occurs all year round in endemic areas, with peaks during the rainy season (Demanou, Pion, et Boussinesq 2001; Akue et al. 2011).

B. Epidemiology

Loasis is endemic in the tropical forest regions of Central and West Africa, with an estimated 10 million people infected in 10 endemic countries (Zouré et al. 2011; Metzger et Mordmüller 2014). Areas of high prevalence include southern Nigeria, Cameroon, Gabon, Congo, Equatorial Guinea, northern Angola and parts of the Democratic Republic of Congo (Zouré et al., 2011). *Loa loa* filariasis is not endemic in Senegal or Niger, although sporadic cases have been reported. These include a student from central Africa in Senegal (Diagne et al. 2018) and a 25-year-old boy in Niger (Adam et al. 2020). In Nigeria, loasis has been reported in people aged 65 to 82 years, with a prevalence of 14.3% in men from venous blood samples (Ojurongbe et al. 2015). A 2016 study in southern Nigeria found a 6.3% prevalence of *L. loa* microfilariae in villages hypoendemic for onchocerciasis, with a mean of 326 mf/ml among positives (Emukah et al. 2018).

III. Pathogenicity and clinical signs

The clinical signs of filariasis only appear when the parasites have reached the adult stage. They depend on the species involved and the location of the adult worms and microfilariae (Bouchery et al. 2013).

a) Lymphatic filariasis

The causal agent of lymphatic filariasis is the roundworm *Wuchereria bancrofti* (Famakinde 2018). Mosquitoes of the genera *Aedes*, *Anopheles*, *Culex* and *Mansonia* ingest microfilariae during a blood meal when they bite humans (Erickson et al. 2013). During this process (Figure 2), an infected mosquito deposits third-stage (infective) larvae on the skin of the human host, where they enter through the bite wound.

Adult viviparous females are larger than males, measuring 80 to 100 mm in length and 0.24 to 0.30 mm in diameter, while males measure around 40 mm in length and 0.1 mm in diameter. Females produce microfilariae that circulate permanently in the lymph and periodically in the blood (either nocturnally or diurnally (P. P. Aubry et Docteur Bernard-Alex Gaüzère 2023)).

These microfilaria are mobile and pass into the blood at night. They live for 1 to 2 months before transforming into adult worms.

Adult filariae are mainly found in the lymphatic vessels or sinuses of the lymph nodes of the lower limbs and scrotum (Chakraborty et al. 2013). Macrofilariae (adult worms) have a circadian periodicity linked to the vector's lifestyle and can survive 5 to 8 years in their host (Taylor, Hoerauf, et Bockarie 2010; Khan et al. 2015). Microfilariae present in blood have specific characteristics that are easily identified using thick Giemsa-stained smears. In mosquito vectors, microfilariae undergo several transformations: coiled forms (stage 1), intermediate forms (stage 2) and infectious forms (stage 3).

Lymphatic filariasis is often asymptomatic in many individuals. It is generally contracted during childhood and causes invisible lesions of the lymphatic system. Clinical manifestations range from acute symptoms - such as adenolymphangitis, filarial fever and tropical pulmonary eosinophilia - to chronic symptoms, including hydrocele, lymphoedema and elephantiasis (Figure 3) in severe cases (Chandy et al. 2011).

According to Haddix et al, hydrocele and lymphoedema can lead to substantial reductions in productivity, which can last a lifetime (Haddix et Kestler 2000). Hydrocoele: this condition, caused by *W. bancrofti* in male patients, is characterised by the accumulation of fluid between the two layers of the tunica vaginalis in the scrotum. It is associated with the presence of adult parasites in the lymphatic vessels of the scrotum (Noroës et al. 2003; Babu et Nutman 2012).

Lymphatic filariasis can also lead to mental health problems and social and financial difficulties, contributing to both stigma and poverty.

In West Africa, particularly Senegal, elephantiasis is one of the most debilitating health problems associated with lymphatic filariasis. The condition causes permanent swelling of a lower limb but can also affect the arms and genitals. The affected limb is at risk of secondary streptococcal infections, which can worsen the disease (Shenoy 2008).




Figure 2. Life cycle of *Wuchereria bancrofti*. Created with BioRender.com.



A= Hydrocèle



B= lymphedema



C= Elephantiasis

Figure 3. Clinical signs of lymphatic filariasis (a and b) = cases from our study in Kolda and Sedhiou

b) Loiasis

Loiasis is a cutaneous-dermal parasite that occurs exclusively in Africa. The vector flies live in the forests of equatorial Africa and take their blood meal during the day. After infection (Figure 4), the filaria penetrates the subcutaneous tissues and rapidly mature. They migrate in the subcutaneous tissues for more than 20 years, with females producing thousands of

microfilariae that can circulate in the blood for 6 to 12 months (S. D. S. Pion et al. 2006; Díaz-Menéndez et al. 2011). The microfilariae produced by females circulate in peripheral blood with a diurnal periodicity (S. D. Pion et al. 2020) and probably reside in lung tissue at night, although this has not been formally demonstrated (M. Boussinesq 2006). Unlike other filaria, *Loa loa* does not harbour a *Wolbachia* endosymbiont (Büttner et al. 2003; McGarry et al. 2003), and it does not appear to have developed new metabolic capacities to compensate (Desjardins et al. 2013).

Clinical signs of the disease may appear years after infection, often after the age of 21 (Richardson et al. 2012). Infected individuals may remain asymptomatic for long periods, with symptoms often mild and transient. Common manifestations include allergic, temporary and migratory oedema localised to the limbs, known as ‘Calabar oedema’ (Jain et al. 2008), as well as subcutaneous crawling of the worm. Generalised symptoms may include Pruritus (itching), Urticaria (hives), Myalgia (muscle pain), Arthralgia (joint pain), Fatigue, Visible migration of adult worms under the skin.

In severe cases, serious complications such as encephalopathy may occur in heavily infested patients. Rarely, subconjunctival manifestations (worms visible in the eye) are observed in affected individuals (Meda et al. 2009; Varenne et al. 2016).




Figure 4. Life cycle of *Loa loa*. Created with BioRender.com.

c) Onchocerciasis

The primary organ affected by *Onchocerca volvulus* infection is the skin, but it can also cause severe visual impairment, making it the world's second leading cause of blindness after trachoma (Gyasi, Okonkwo, et Tripathy 2024). Infectious microfilarial larvae (stage L3) are introduced into the human body during a blood meal taken by a female black fly (Figure 5). Over a period of 6 to 12 months, these larvae develop into adult worms (macrofilariae) that reside in subcutaneous fibrous nodules, typically located around the iliac crest, head, and torso. Female worms are fertilized by males within the nodules and produce large quantities of microfilariae. These nodules are irrigated by blood and lymphatic vessels (Attout et al. 2009). Microfilariae are unsheathed and measure approximately 300 µm in length. Some of these microfilariae are found in lymphatic vessels, which they may use to reach the dermis, where they become accessible to Simulium vectors (Attout et al. 2009; Allen et al. 2008). Microfilariae can also be detected in the subconjunctival tissues of the eye.

The pathogenesis of onchocerciasis is thought to result from the chronic inflammatory immune response triggered by the death of microfilariae in the skin or eyes. This reaction is likely exacerbated by their Wolbachia endosymbionts, which contribute to the inflammatory process (Ngwewondo, Scandale, et Specht 2021). Symptoms include debilitating pruritus linked to microfilariae in the dermis (Desoubeaux, Chesnay, et Chandenier 2020) and may progress to permanent blindness. Onchocerciasis has also been associated with neurological disorders, including epilepsy (Chesnais et al. 2020), nodding syndrome (Geelhand De Merxem et al. 2019), and growth retardation. Adult worms have a lifespan of up to 15 years, during which the infection may remain asymptomatic or progress to various clinical manifestations such as papules and inflammatory lesions, skin changes (leopard skin, loose skin, skin atrophy and fibrosis), punctate keratitis, sclerosing keratitis, subcutaneous nodules, systemic signs and psychological and social complications (Hartley 2021; CDC 2024).

The incubation period ranges from 1–2 weeks on average but can extend up to 4 years (Antinori et al. 2017). In sub-Saharan Africa, onchocerciasis remains a leading cause of visual disturbances and blindness (Thylefors 2004).




Figure 5. Life cycle of *Onchocerca volvulus*. Created with BioRender.com.

d) Mansonellosis

An infected *Culicoides* or *Simulium* female introduces infective filarial larvae (L3) into the human host during a blood meal. The larvae develop into adult filariae, which typically reside in the pleural cavity, peritoneal cavity, or subcutaneous dermal layer (Figure 6). Female worms produce microfilariae, which are found in: Peripheral blood (*M. perstans* and *M. ozzardi*), Skin (*M. streptocerca* and *M. ozzardi*). When an arthropod ingests microfilariae during a blood meal, the larvae undergo two molts within the vector to become infectious L3 larvae (Tamarozzi et al. 2022). These three parasite species (*M. perstans*, *M. ozzardi*, and *M. streptocerca*) have similar life cycles and epidemiology. They generally cause mild, non-specific symptoms or remain asymptomatic (Ta-Tang et al. 2018; Medeiros et al. 2017). Adult *M. perstans* worms are primarily found in serosal cavities, visceral adipose tissue, or skin tissue, while microfilariae develop in the blood and lymphatic system (Ta-Tang et al. 2018; Mediannikov et Ranque 2018a).

The pathogenicity of *Mansonella* species remains under debate. Infections are often asymptomatic, but certain clinical manifestations are associated with specific species:

M. perstans and *M. ozzardi*: linked to pruritus, headaches, edema of the lower limbs, arthralgia, inguinal lymphadenopathy, hydrocele, and ocular pathologies (Medeiros et al. 2017; Carme et Esterre 2012). *M. streptocerca*: primarily associated with skin lesions and inguinal damage. Although *M. perstans* is often asymptomatic, it may cause symptoms such as: Pruritus (itching), Urticaria (hives), Fatigue, Skin rashes, arthralgia, myalgia, fever, and neurological disorders (Tamarozzi et al. 2022).

In some cases, *M. perstans* infections present with non-specific manifestations similar to loiasis or other filarial infections, such as: localized or generalized pruritus, transient edema in various parts of the body, Headache, lymphadenopathy, and rash (Bottieau et al. 2022; Puente et al. 2020). In non-endemic areas, *M. perstans* infections have been associated with abdominal pain, asthma-like symptoms, pericarditis, pleuritis, and granulomatous nodules surrounding dead adult worms (Puente et al. 2020; Ferreira, Crainey, et Gobbi 2023; Gobbi et al. 2017).




Figure 6. Life cycle of *Mansonella perstans*. Created with BioRender.com.

IV. Diagnostic methods

Classically, the diagnosis of filariasis is based on the detection (direct or indirect) of adult worms or microfilariae produced by adult females. Several diagnostic methods are available to detect these parasites :

- Blood microscopy: Used for species found in humans (*L. loa*, *W. bancrofti*, *Brugia malayi*, and *M. perstans*).
- Skin biopsy: Used for *O. volvulus* and *M. streptocerca*.
- Both methods: Utilized for *M. ozzardi* species (Kamgno et al. 2017).

In some cases, microfilariae can even be found in urine (Gaillard et al. 2020). Various other techniques, including antigen/antibody detection and polymerase chain reaction (PCR), are used for individual diagnoses and to assess infection endemicity in communities (Drame et al. 2016).

i. Lymphatic Filariasis

The diagnosis of lymphatic filariasis involves the detection of microfilariae in the blood or adult worms through direct and indirect examination.

1. Microscopy

Fresh blood examination: A drop of blood is collected after finger disinfection using a vaccinostyle (10–20 μ L). The sample is placed on a slide and examined under low magnification (x50 or x100). Mobile microfilariae can be visualized among red blood cells (Desoubeaux, Chesnay, et Chandenier 2020; Duong et Richard-Lenoble 2008).

- Leucoconcentration technique: This involves enrichment by hemolysis using a 2% saponin solution, followed by centrifugation. The pellet is smeared, stained, and examined under a microscope (Herbeuval, Herbeuval, et Duheille 1962; Walther et Muller 2003). This technique increases sensitivity and detects parasite loads as low as 1 mf/ μ L (Melrose 2002).

Thin and thick smears: Blood volumes of 5–20 μ L are used to prepare smears, stained with May-Grünwald Giemsa. This technique helps in identifying species based on morphological characteristics:

- Microfilariae length: 250–300 μ m.
- Thickness: 7–8 μ m with a sheath and visible internal body (Frédéric 2018; Duong et Richard-Lenoble 2008).

432 ○ Posterior end: Tapered.

433 ○ Cephalic end: Large or small.

434 The limit for microfilariae detection using this method is 15 mf/mL, making it less sensitive in
435 low-density infections (Melrose 2002; Frédéric 2018).

436 • Knott's technique: This involves adding 1 mL of blood to 9 mL of 1% formalin,
437 followed by centrifugation. The sediment is then examined under a microscope. This
438 method allows precise measurement and preserves microfilariae morphology (Duong et
439 Richard-Lenoble 2008).

440 2. Imaging Techniques

441 • Lymph node biopsy: In rare cases, *W. bancrofti* macrofilariae can be identified during
442 histopathological examination.

443 • Lymphatic ultrasound: Detects adult worms through the “filaria dance sign”
444 movements. This is a highly sensitive method for detecting worms in asymptomatic or
445 amicrofilaremic patients (Desoubieux, Chesnay, et Chandenier 2020). However, it
446 becomes ineffective in advanced stages of lymphedema, as adult worms are no longer
447 present (Mendoza et al. 2009).

448 3. Immunological Tests

449 • Antigen and antibody detection:

450 ○ Techniques include ELISA, indirect immunofluorescence (IFI), and
451 immuno-electrophoresis (IELP), using crude antigens from animal filarial
452 species (*Dirofilaria immitis*, *Dipetalonema viteae*, etc.) (Desoubieux, Chesnay,
453 et Chandenier 2020).

454 ○ These tests can detect pan-filarial antibodies, but they lack specificity and may
455 cross-react with other nematodes (Melrose 2002).

456 • ICT (Immunochromatographic Card Test) and FTS (Filariasis Test Strip):
457 These tests detect circulating filarial antigens in blood samples collected from the
458 fingertip at any time of day. They are highly sensitive and effective for diagnosing

lymphatic filariasis but cannot differentiate between past and active infections (Knopp et al. 2012; Cromwell et al. 2020).

4. Molecular Techniques

- Polymerase chain reaction (PCR): PCR is highly sensitive and specific, capable of detecting *W. bancrofti* and *B. malayi* in blood, urine, and tissue samples. It is primarily used in research to monitor infection prevalence and evaluate control (Walther et Muller 2003; Supali et al. 2006).

ii. Loiasis

The diagnosis of loiasis relies on the detection of microfilariae in blood samples. Key methods include:

Microscopic Examination: Detection is performed on fresh blood or after staining with thin or thick smear techniques. For cases of low microfilaremia or negative initial examinations, advanced methods such as the Knott technique and nucleopore membrane filtration are employed (CAMPILLO et al. 2023).

Sample Timing: Due to the diurnal periodicity of *Loa loa* microfilariae, blood samples should be collected around midday.

Adult Worm Detection: Adult helminths can sometimes be identified migrating under the skin or in the conjunctiva. Their extraction may require tools like a vaccinostyle or clawless forceps, or through scarification (Díaz-Menéndez et al. 2011; Varenne et al. 2016).

Morphological Identification: Microfilariae have a sheath, poorly stained by Giemsa, and exhibit visible ovoid nuclei overlapping to the tail but sparing a 5 µm space at the head (Bourée 2005).

Antibody Detection: Methods include ELISA, indirect immunofluorescence (IFI), and co-electrosyneresis (COES), with a focus on Pan-filarial antigens.

Advanced techniques like Luciferase Immunoprecipitation Systems (LIPS) detect IgG4 antibodies specific to *L. loa* (Burbelo et al. 2008).

Molecular Techniques: PCR is used for specific *L. loa* DNA detection and as an epidemiological tool in control programs (Knopp et al. 2012).

iii. Onchocerciasis

For onchocerciasis, several diagnostic techniques exist:

1. Microfilariae Visualization:

490 ○ Skin Snip Test: Small skin biopsies (from iliac crests, shoulder blades, or other
491 regions) are placed in saline to observe microfilaria emerging from the dermis.

492 Staining: Using May-Grünwald Giemsa, microfilariae with elongated nuclei and curved tails
493 are identified (Desoubeaux, Chesnay, et Chandenier 2020).

494 2. Histological Examination:

495 ○ Biopsies or nodules are examined post-staining with hematoxylin and eosin to
496 identify adult worms (Carme et Esterre 2012).

497 3. Immunological Tests:

498 ○ ELISA tests detect IgG4 antibodies against recombinant *O. volvulus* antigen Ov-
499 16.

500 ○ Rapid diagnostic tests (RDTs) or LIPS tests are also available ((Vlaminck,
501 Fischer, et Weil 2015).

502 4. Molecular Techniques:

503 ○ PCR amplifies repetitive O-150 DNA sequences from skin snips or nodules,
504 significantly increasing diagnostic sensitivity (Alhassan et al. 2015).

505 5. Specialized Tests:

506 ○ Patch Test: Using diethylcarbamazine to provoke localized erythema, indicating
507 onchocerciasis presence.

508 ○ Ultrasound: Detects adult worms in nodules through their characteristic
509 movement (Unnasch et al. 2018).

510 **iv. Mansonellosis**

511 The diagnosis of mansonellosis is species-specific:

512 1. Microscopy:

513 ○ *M. perstans*: Detected in blood samples.

514 ○ *M. streptocerca*: Identified through skin biopsies.

515 ○ *M. ozzardi*: Found in both blood and dermal samples (Frédéric 2018).

516 2. Molecular Techniques:

517 ○ PCR allows the differential diagnosis of *M. streptocerca* from *O. volvulus*, *W.*
518 *bancrofti*, and *L. loa* (Walther et Muller 2003).

519 ○ LAMP (Loop-mediated Isothermal Amplification) offers a more rapid
520 alternative for species detection (Poole et al. 2017; Deng et al. 2019).

521 3. Other Sample Sources:

522 ○ Microfilariae of *M. perstans* and *M. ozzardi* have been detected in urine,
523 cerebrospinal fluid, and other body fluids (Gaillard et al. 2020).

524 **v. Prevention**

525 ➤ **Lymphatic Filariasis**

526 Lymphatic filariasis (LF) is a neglected tropical disease whose prevention focuses on
527 raising community awareness about transmission and the importance of preventive measures.
528 Key strategies include:

529 1. Community Awareness and Hygiene: Promoting personal and environmental hygiene
530 to reduce exposure to mosquito vectors.

531 2. Mass Drug Administration (MDA):

532 ○ The Global Programme to Eliminate Lymphatic Filariasis (GPELF), launched
533 in 2000 by WHO, aims to eliminate LF as a public health problem.

534 ○ MDA involves annual administration of albendazole in combination with either
535 diethylcarbamazine (DEC) or ivermectin for at least 5 years in regions with $\geq 1\%$
536 prevalence of filarial antigens (Ottesen 2006; World Health Organization 2011).

537 ○ The goal is to kill circulating microfilariae, thereby reducing transmission
538 (Simonsen et al. 2014).

539 ○ There is no macrofilaricidal drug for MDA, which would eliminate adult worms.

540 3. Vector Control:

541 ○ Intensive use of long-lasting insecticide-treated nets (LLINs) and
542 xenomonitoring (PCR detection of LF parasites in mosquito vectors) has proven
543 effective (Bockarie, Taylor, et Gyapong 2009; Njoku et al. 2013).

544 ○ Addressing stagnant water sources and improving drainage systems also helps
545 control mosquito breeding sites.

546 4. Psychosocial Support:

547 ○ Community programs to combat stigma and provide psychosocial assistance for
548 patients, improving their well-being and reintegration into society.

549

➤ **Onchocerciasis**, or river blindness, is the second leading infectious cause of preventable blindness. Preventive strategies include:

1. Vector Control Programs:

- The Onchocerciasis Control Program (OCP) in West Africa and the African Programme for Onchocerciasis Control (APOC) used aerial spraying of insecticides and mass drug distribution to reduce blackfly populations (Boatin 2008).

2. Mass Drug Administration:

- Ivermectin is distributed annually to affected communities to control skin and eye symptoms and reduce microfilariae levels (Amazigo 2008).

3. Challenges in *L. loa*-Endemic Areas:

- In areas co-endemic with *Loa loa*, ivermectin can cause severe adverse events. Measures are needed to detect high microfilaremia levels before administering the drug (Vinkeles Melchers et al. 2018).

4. Personal Protection:

- Use of insect repellents and protective clothing when traveling through endemic areas.
- There is currently no vaccine for onchocerciasis.

➤ **Loiasis**

Loiasis, caused by *Loa loa*, lacks a global eradication program. Preventive measures focus on:

1. Vector Control:

- Reducing exposure to Chrysops vectors using insect repellents (DEET), permethrin-treated clothing, and mosquito nets, although the latter are less effective due to the diurnal activity of vectors.

2. Chemoprophylaxis:

- Weekly doses of diethylcarbamazine (DEC) can reduce risk but carry the potential for side effects, particularly in high microfilaremia cases (Frédéric 2018).
- Initial treatment with albendazole, a less potent microfilaricide, is recommended to minimize adverse reactions before stronger drugs like ivermectin or DEC are administered.

3. Monitoring:

- Identifying and monitoring high-risk individuals with elevated microfilaremia to minimize complications.

➤ **Mansonellosis** remains poorly studied and lacks targeted control programs. Preventive efforts include:

Insecticide Use:

Indoor residual spraying (IRS) has shown promise in reducing *Culicoides* populations, although its effectiveness for mansonellosis control depends on vector behavior (González et al. 2020).

Limitations of Bed Nets:

Bed nets, effective for malaria and LF, are less impactful for mansonellosis because *Culicoides* are small and typically bite outdoors (Brugman et al. 2017; Harrup et al. 2016).

Research Needs:

More studies are needed to understand vector ecology and optimize control strategies globally.

VI. Treatment

❖ Lymphatic Filariasis

Since the launch of the Global Programme to Eliminate Lymphatic Filariasis (GPELF) by the WHO in 2000, the main strategy has been to stop the spread of the disease through Mass Drug Administration (MDA). This involves large-scale preventive chemotherapy, which is primarily aimed at eliminating microfilariae in the bloodstream rather than adult worms.

Treatment Regimens:

1. Ivermectin (150-200 µg/kg) + Albendazole (400 mg): Used in regions co-endemic with onchocerciasis.
2. Diethylcarbamazine (6 mg/kg) + Albendazole (400 mg): Recommended in areas free of onchocerciasis.
3. Albendazole alone (400 mg): Administered twice a year in regions where loiasis is endemic.

These treatments do not kill the adult parasites but remove microfilariae from the blood, thereby preventing further transmission. For the treatment to be effective, it must be administered annually for at least five years with complete coverage of the at-risk population (Rebollo et Bockarie 2017).

- Doxycycline (200 mg/day for 4–6 weeks) targets *Wolbachia* bacteria, which live in symbiosis with adult worms, leading to the gradual death of the parasites (Ottesen 2000).

❖ Mansonellosis is difficult to treat due to the resistance of *M. perstans* to standard antifilarial drugs. Some treatments have shown varying levels of success:

1. Combination therapy with diethylcarbamazine and mebendazole is effective in reducing microfilaremia.
2. Thiabendazole alone has been shown to significantly reduce microfilariae in the blood.
3. Ivermectin and praziquantel are not effective for *M. perstans* infections.
4. Mebendazole alone is an alternative, more effective than diethylcarbamazine alone in clearing *M. perstans* infections (Medeiros et al. 2017; Enrico Rino Bregani et al. 2006).

Additionally, doxycycline has been found effective for *M. perstans* infections, especially in West Africa, where *M. perstans* carries Wolbachia bacteria. However, its efficacy is uncertain in regions where Wolbachia is absent in *M. perstans* populations (Yaya I. Coulibaly et al. 2009a).

- Ivermectin has shown effectiveness in reducing microfilariae of *M. ozzardi* in short and long terms (Shelley et Coscarón 2001).
- Diethylcarbamazine and ivermectin are also effective against *M. streptocerca*, with ivermectin suppressing microfilariae for up to one year (Pérez-Molina et al. 2010).

❖ Treating onchocerciasis is essential to prevent skin damage and long-term blindness (Antinori et al. 2017).

1. Ivermectin kills microfilariae and temporarily halts their production by adult worms.
 - It is administered annually to populations at risk, but it does not eliminate adult worms.
 - Doxycycline combined with ivermectin has shown promise in significantly reducing both microfilarial loads and adult worm burden, by targeting Wolbachia bacteria that are essential for the survival of adult worms (A. Y. Debrah et al. 2015).

2. Moxidectin, a newer single-dose antiparasitic, has also shown efficacy against onchocerciasis in a study across Ghana, Liberia, and the Democratic Republic of Congo (Opoku et al. 2018).

❖ The treatment of loiasis must be done under medical supervision due to the risk of serious adverse effects:

Diethylcarbamazine is typically used, but it can cause dangerous side effects in patients with high microfilarial densities (encephalopathy).

Albendazole is prescribed first when the microfilarial load exceeds 8000/mL; it is followed by ivermectin or diethylcarbamazine for lower microfilarial densities (Michel Boussinesq 2012).

Ivermectin and diethylcarbamazine are used in the treatment of high microfilarial density, but they may trigger encephalopathy in patients with very high microfilarial counts (Whittaker et al. 2018).

Challenges in Treatment

Albendazole-induced encephalopathy has been reported, particularly in the context of high *L. loa* microfilarial densities, emphasizing the need for caution when administering these drugs (Volpicelli et al. 2020; Métais, Michalak, et Rousseau 2021).

Table 2. Main characteristics of recommended medications for *Loa loa* infection.

	Action mechanism	Recommended use	Dosage	Benefits
Diethylcarbamazine	Microfilaricide and Macrofilaricide	Number of Mf < 2000 mf/ml	1 mg/kg/day, increasing over 3 days to 9 mg/kg/day for 3 weeks	Rapid macrofilaricidal effect
Ivermectine	Microfilaricide	Number of Mf between 2000/ml and 8000/ml	150-200 µg/kg, single or multiple dose	Rapid microfilaricidal effect, high efficiency in combination with ALB
Albendazole	Macrofilaricide (Embryotoxic effect), Slow effect	Mf rate greater than 8000/ml	200-400 mg twice daily for 21-28 days	Fewer adverse events, high efficacy in combination with IVM

Conclusion

This review highlights the crucial importance of filariasis as a public health problem in tropical areas, particularly in West Africa, where it imposes a heavy health and socio-economic burden on affected populations. The wide spectrum of filariasis, including lymphatic infections such as lymphatic filariasis, skin infections such as onchocerciasis, and serous infections such as mansonellosis, illustrates the diversity and complexity of this pathology. Current diagnostic and control strategies, although they have made significant progress, show limits in the sustainable elimination of transmission, partly due to the complexity of the biological cycles of filariae and the diversity of vectors involved. Massive treatment interventions with drugs like

ivermectin have been key to reducing morbidity rates, but challenges persist, particularly in co-endemic areas where some treatments can cause serious adverse effects.

For more effective and long-term control of filariasis in West Africa, several perspectives and recommendations are necessary. It is essential to: Strengthen diagnostic capacities: Developing more specific and less expensive diagnostic methods, including rapid field tests, will help better target interventions and reduce transmission in endemic areas. Develop integrated prevention strategies: This includes controlling vectors through means such as insecticide-treated bed nets and introducing community awareness programs to encourage the adoption of personal protection measures. Support research into new treatments: Exploring new therapeutic agents less likely to cause adverse effects in areas co-endemic with *Loa loa*, for example, could strengthen adherence to mass treatment programs and improve outcomes sanitary facilities.

Improve epidemiological surveillance: Continuous and detailed monitoring of the prevalence of infections will make it possible to quickly detect resurgences and adjust interventions, accordingly, thus ensuring the effectiveness of long-term control programs. Strengthen international partnerships: Collaboration between governments, international organizations, and local communities is essential to support efforts to eliminate filariasis in West Africa and mobilize the resources necessary to achieve this ambitious goal. Adopting these integrated approaches could transform filariasis control efforts, reducing transmission and easing the socio-economic burden in affected regions.

Références

- Adam, Nouhou Diori, Yacoubou Soumana, Ali Saley, et Idrissa Saley. 2020. « Une Loase Oculaire: À Propos D'un Cas ». *Pan African Medical Journal* 36. <https://doi.org/10.11604/pamj.2020.36.302.25124>.
- Adewole, S.O., et S.K. Ayeni. 2009. « Clinical Manifestation of Onchocerciasis in Ise - Orun Local Government, Ekiti State, Nigeria ». *Pakistan Journal of Nutrition* 8 (2): 122-24. <https://doi.org/10.3923/pjn.2009.122.124>.
- Adhikari, Ram Kumar, Jeevan Bahadur Sherchand, Shiva Raj Mishra, Kamal Ranabhat, Amrit Pokharel, Pramila Devkota, Durga Mishra, et al. 2015. « Health-Seeking Behaviors and Self-Care Practices of People with Filarial Lymphoedema in Nepal: A Qualitative Study ». *Journal of Tropical Medicine* 2015:1-6. <https://doi.org/10.1155/2015/260359>.
- Adolph, Paul E., Irving G. Kagan, et Russell M. McQuay. 1962. « Diagnosis and Treatment of Acanthocheilonema Perstans Filariasis », janvier. <https://doi.org/10.4269/ajtmh.1962.11.76>.
- Agbolade, Olufemi Moses, Dora Olufunmilola Akinboye, Taiwo Monroof Olateju, Oluwatoyin Adepeju Ayanbiyi, Olukayode Olakunle Kuloyo, et Oluwalolami Oluwaseun Fenuga. 2006. « Biting of Anthropophilic Culicoides Fulvithorax (Diptera: Ceratopogonidae), a Vector of Mansonella Perstans in Nigeria ». *The Korean Journal of Parasitology* 44 (1): 67-72. <https://doi.org/10.3347/kjp.2006.44.1.67>.
- Akue, Jean Paul, Dieudonné Nkoghe, Cindy Padilla, Ghislain Moussavou, Hubert Moukana, Roger Antoine Mbou, Benjamin Ollomo, et Eric Maurice Leroy. 2011. « Epidemiology of Concomitant Infection Due to Loa Loa and Mansonella Perstans in Gabon ». Édité par Rachel Lisa Pullan. *PLoS Neglected Tropical Diseases* 5 (10): e1329. <https://doi.org/10.1371/journal.pntd.0001329>.
- Alhassan, Andy, Zhiru Li, Catherine B. Poole, et Clotilde K.S. Carlow. 2015. « Expanding the MDx Toolbox for Filarial Diagnosis and Surveillance ». *Trends in Parasitology* 31 (8): 391-400. <https://doi.org/10.1016/j.pt.2015.04.006>.
- Allen, Judith E., Ohene Adjei, Odile Bain, Achim Hoerauf, Wolfgang H. Hoffmann, Benjamin L. Makepeace, Hartwig Schulz-Key, et al. 2008. « Of Mice, Cattle, and Humans: The Immunology and Treatment of River Blindness ». Édité par Sara Lustigman. *PLoS Neglected Tropical Diseases* 2 (4): e217. <https://doi.org/10.1371/journal.pntd.0000217>.
- Alsarraf, Mustafa, Viktoria Levytska, Ewa J. Mierzejewska, Vasyl Poliukhovych, Anna Rodo, Mohammed Alsarraf, Dziyana Kavalevich, Dorota Dwużnik-Szarek, Jerzy M. Behnke, et Anna Bajer. 2021. « Emerging Risk of Dirofilaria Spp. Infection in Northeastern

Europe: High Prevalence of *Dirofilaria Repens* in Sled Dog Kennels from the Baltic Countries ». *Scientific Reports* 11 (1): 1068. <https://doi.org/10.1038/s41598-020-80208-1>.

Amazigo, U. 2008. « The African Programme for Onchocerciasis Control (APOC) ». *Annals of Tropical Medicine & Parasitology* 102 (supl): 19-22. <https://doi.org/10.1179/136485908X337436>.

Anderson, Roy C. 2000. « R.C. Anderson: Nematode Parasites of Vertebrates. Their Development and Transmission. 2nd Edition. CABI Publishing, Wallingford, Oxon (UK) 2000. 650 Pp., 43 Figs. Price £ 99.50 . », 1.

Anosike, Jude C, Ikechukwu N S Dozie, Celestine O E Onwuliri, Bertram E B Nwoke, et Viola A Onwuliri. 2005. « PREVALENCE OF MANSONELLA PERSTANS INFECTIONS AMONG THE NOMADIC FULANIS OF NORTHERN NIGERIA », p.35-39. <https://pubmed.ncbi.nlm.nih.gov/16028864/>.

Antinori, Spinello, Carlo Parravicini, Laura Galimberti, Antonella Tosoni, Paolo Giunta, Massimo Galli, Mario Corbellino, et Anna Lisa Ridolfo. 2017. « Is Imported Onchocerciasis a Truly Rare Entity? Case Report and Review of the Literature ». *Travel Medicine and Infectious Disease* 16 (mars):11-17. <https://doi.org/10.1016/j.tmaid.2017.02.006>.

Asio, S. M., P. E. Simonsen, et A. W. Onapa. 2009. « *Mansonella Perstans* : Safety and Efficacy of Ivermectin Alone, Albendazole Alone and the Two Drugs in Combination ». *Annals of Tropical Medicine & Parasitology* 103 (1): 31-37. <https://doi.org/10.1179/136485909X384929>.

Asio, Santa Maria, Paul E. Simonsen, et Ambrose W. Onapa. 2009a. « Analysis of the 24-h Microfilarial Periodicity of *Mansonella Perstans* ». *Parasitology Research* 104 (4): 945-48. <https://doi.org/10.1007/s00436-008-1312-x>.

———. 2009b. « *Mansonella Perstans* Filariasis in Uganda: Patterns of Microfilaraemia and Clinical Manifestations in Two Endemic Communities ». *Transactions of the Royal Society of Tropical Medicine and Hygiene* 103 (3): 266-73. <https://doi.org/10.1016/j.trstmh.2008.08.007>.

Attout, Tarik, Achim Hoerauf, Gaëlle Dénécé, Alexander Yaw Debrah, Yeboah Marfo-Debrekyei, Michel Boussinesq, Samuel Wanji, et al. 2009. « Lymphatic Vascularisation and Involvement of Lyve-1+ Macrophages in the Human *Onchocerca* Nodule ». Édité par Derya Unutmaz. *PLoS ONE* 4 (12): e8234. <https://doi.org/10.1371/journal.pone.0008234>.

- Aubry, Arthur, Charles Hervochon, Marie Chauveau, Marie Denis-Musquer, Fakhri Jeddi, Rose-Anne Lavergne, Patrice Le Pape, et Florent Morio. 2022. « Une expulsion bien singulière d'un ver adulte de *Loa loa* ». *Revue Francophone des Laboratoires* 2022 (538): 72-75. [https://doi.org/10.1016/S1773-035X\(21\)00381-6](https://doi.org/10.1016/S1773-035X(21)00381-6).
- Aubry, Professeur Pierre, et Docteur Bernard-Alex Gaüzère. 2023. « Filarioses lymphatiques Actualités 2023 », P8.
- Babu, Subash, et Thomas B. Nutman. 2012. « Immunopathogenesis of Lymphatic Filarial Disease ». *Seminars in Immunopathology* 34 (6): 847-61. <https://doi.org/10.1007/s00281-012-0346-4>.
- Badia-Rius, Xavier, Salissou Adamou, Mark J. Taylor, et Louise A. Kelly-Hope. 2023. « Morbidity Hotspot Surveillance: A Novel Approach to Detect Lymphatic Filariasis Transmission in Non-Endemic Areas of the Tillabéry Region of Niger ». *Parasite Epidemiology and Control* 21 (mai):e00300. <https://doi.org/10.1016/j.parepi.2023.e00300>.
- Bassene, Hubert, Masse Sambou, Florence Fenollar, Siân Clarke, Sawdiatou Djiba, Gaël Mourembou, Alioune Badara L Y, Didier Raoult, et Oleg Mediannikov. 2015a. « High Prevalence of *Mansonella Perstans* Filariasis in Rural Senegal ». *The American Journal of Tropical Medicine and Hygiene* 93 (3): 601-6. <https://doi.org/10.4269/ajtmh.15-0051>.
- . 2015b. « High Prevalence of *Mansonella Perstans* Filariasis in Rural Senegal ». *The American Journal of Tropical Medicine and Hygiene* 93 (3): 601-6. <https://doi.org/10.4269/ajtmh.15-0051>.
- Bhalla, Devender, Michel Dumas, et Pierre-Marie Preux. 2013. « Neurological Manifestations of Filarial Infections ». In *Handbook of Clinical Neurology*, 114:235-42. Elsevier. <https://doi.org/10.1016/B978-0-444-53490-3.00018-2>.
- Boatin, B. 2008. « The Onchocerciasis Control Programme in West Africa (OCP) ». *Annals of Tropical Medicine & Parasitology* 102 (sup1): 13-17. <https://doi.org/10.1179/136485908X337427>.
- Bockarie, Moses J, et Rinki M Deb. 2010. « Elimination of Lymphatic Filariasis: Do We Have the Drugs to Complete the Job? ». *Current Opinion in Infectious Diseases* 23 (6): 617-20. <https://doi.org/10.1097/QCO.0b013e32833fdee5>.
- Bockarie, Moses J, Mark J Taylor, et John O Gyapong. 2009. « Current Practices in the Management of Lymphatic Filariasis ». *Expert Review of Anti-Infective Therapy* 7 (5): 595-605. <https://doi.org/10.1586/eri.09.36>.

795 Bottieau, Emmanuel, Ralph Huits, Steven Van Den Broucke, Ula Maniewski, Steven Declercq,
 796 Isabel Brosius, Caroline Theunissen, et al. 2022. « Human Filariasis in Travelers and
 797 Migrants: A Retrospective 25-Year Analysis at the Institute of Tropical Medicine,
 798 Antwerp, Belgium ». *Clinical Infectious Diseases* 74 (11): 1972-78.
 799 <https://doi.org/10.1093/cid/ciab751>.

800 Bouchery, T., E. Lefoulon, G. Karadjian, A. Nieguitsila, et C. Martin. 2013. « The Symbiotic
 801 Role of Wolbachia in Onchocercidae and Its Impact on Filariasis ». *Clinical
 802 Microbiology and Infection* 19 (2): 131-40. <https://doi.org/10.1111/1469-0691.12069>.

803 Bourée, P. 2005. « Parasitoses urinaires ». *Annales d'Urologie* 39 (6): 232-46.
 804 <https://doi.org/10.1016/j.anuro.2005.09.005>.

805 Boussinesq, M. 2003. « L'onchocercose humaine en Afrique ». In *Trente ans de lutte contre
 806 l'onchocercose en Afrique de l'Ouest. Traitements larvicides et protection de
 807 l'environnement : Thirty years of onchocerciasis control in West Africa. Blackfly
 808 larviciding and environmental protection*, édité par Laurent Yaméogo, Christian
 809 Levêque, et Jean-Marc Hougard, 389-400. Synthèses. Marseille: IRD Éditions.
 810 <https://doi.org/10.4000/books.irdeditions.28767>.

811 ———. 2006. « Loiasis ». *Annals of Tropical Medicine & Parasitology* 100 (8): 715-31.
 812 <https://doi.org/10.1179/136485906X112194>.

813 Boussinesq, Michel. 2012. « Loiasis: New Epidemiologic Insights and Proposed Treatment
 814 Strategy ». *Journal of Travel Medicine* 19 (3): 140-43. [https://doi.org/10.1111/j.1708-
 815 8305.2012.00605.x](https://doi.org/10.1111/j.1708-8305.2012.00605.x).

816 Brattig, Norbert W., Robert A. Cheke, et Rolf Garms. 2021. « Onchocerciasis (River Blindness)
 817 – More than a Century of Research and Control ». *Acta Tropica* 218 (juin):105677.
 818 <https://doi.org/10.1016/j.actatropica.2020.105677>.

819 Bregani, E. R., F. Tantardini, et A. Rovellini. 2007. « [Mansonella perstans filariasis] ». *Parassitologia* 49 (1-2): 23-26.

820
 821 Bregani, Enrico Rino, Angelo Rovellini, Narasem Mbaïdoun, et Maria Giovanna Magnini.
 822 2006. « Comparison of Different Anthelmintic Drug Regimens against Mansonella
 823 Perstans Filariasis ». *Transactions of the Royal Society of Tropical Medicine and
 824 Hygiene* 100 (5): 458-63. <https://doi.org/10.1016/j.trstmh.2005.07.009>.

825 Brengues, J., R. Subra, J. Mouchet, et G. S. Nelson. 1968. « [Transmission of Wuchereria
 826 bancrofti Cobbold in West Africa. Preliminary study of a focus in the savanna of north
 827 Guinea] ». *Bulletin of the World Health Organization* 38 (4): 595-608.

828 Brugman, Victor A., Marion E. England, Joanne Stoner, Laura Tugwell, Lara E. Harrup,
829 Anthony J. Wilson, Jolyon M. Medlock, et al. 2017. « How Often Do Mosquitoes Bite
830 Humans in Southern England? A Standardised Summer Trial at Four Sites Reveals
831 Spatial, Temporal and Site-Related Variation in Biting Rates ». *Parasites & Vectors* 10
832 (1): 420. <https://doi.org/10.1186/s13071-017-2360-9>.

833 Burbelo, Peter D., Roshan Ramanathan, Amy D. Klion, Michael J. Iadarola, et Thomas B.
834 Nutman. 2008. « Rapid, Novel, Specific, High-Throughput Assay for Diagnosis of *Loa*
835 *Loa* Infection ». *Journal of Clinical Microbiology* 46 (7): 2298-2304.
836 <https://doi.org/10.1128/JCM.00490-08>.

837 Büttner, Dietrich W., Samuel Wanji, Chiara Bazzocchi, Odile Bain, et Peter Fischer. 2003.
838 « Obligatory Symbiotic Wolbachia Endobacteria Are Absent from *Loa Loa* ». *Filaria*
839 *Journal* 2 (1): 10. <https://doi.org/10.1186/1475-2883-2-10>.

840 CAMPILLO, Jérémy T., Frédéric LOUYA, Paul BIKITA, François MISSAMOU, Michel
841 BOUSSINESQ, Sebastien D. S. PION, Sébastien BERTOUT, et Cédric B CHESNAIS.
842 2023. « Évaluation en milieu rural au Congo de la fiabilité de la méthode de la goutte
843 épaisse calibrée pour évaluer le niveau de la microfilarémie à *Loa loa* ». *MTSI*, mars,
844 Vol. 3 No 1 (2023): MTSIRevue. <https://doi.org/10.48327/MTSI.V3I1.2023.297>.

845 Carne, B., et P. Esterre. 2012. « Filarioses ». *EMC - Maladies infectieuses* 9 (2): 1-19.
846 [https://doi.org/10.1016/S1166-8598\(12\)41786-0](https://doi.org/10.1016/S1166-8598(12)41786-0).

847 CDC. 2024. « Symptoms of Onchocerciasis ». Filarial Worms. 5 juin 2024.
848 <https://www.cdc.gov/filarial-worms/signs-symptoms/onchocerciasis.html>.

849 Chakraborty, Sanjukta, Manokaran Gurusamy, David C. Zawieja, et Mariappan Muthuchamy.
850 2013. « Lymphatic Filariasis: Perspectives on Lymphatic Remodeling and Contractile
851 Dysfunction in Filarial Disease Pathogenesis ». *Microcirculation* 20 (5): 349-64.
852 <https://doi.org/10.1111/micc.12031>.

853 Chandy, Anish, Alok Singh Thakur, Mukesh Pratap Singh, et Ashish Manigauha. 2011. « A
854 Review of Neglected Tropical Diseases: Filariasis ». *Asian Pacific Journal of Tropical*
855 *Medicine* 4 (7): 581-86. [https://doi.org/10.1016/S1995-7645\(11\)60150-8](https://doi.org/10.1016/S1995-7645(11)60150-8).

856 Chelsea, Marie, et A. Petri William. 2022. « Mansonellose-Maladies infectieuses ». [https://www.msdmanuals.com/fr/professional/maladies-](https://www.msdmanuals.com/fr/professional/maladies-infectieuses/n%C3%A9matodes-vers-ronds/mansonellose)
857 [infectieuses/n%C3%A9matodes-vers-ronds/mansonellose](https://www.msdmanuals.com/fr/professional/maladies-infectieuses/n%C3%A9matodes-vers-ronds/mansonellose).

859 Chenna, Ramu, Hideaki Sugawara, Tadashi Koike, Rodrigo Lopez, Toby J. Gibson, Desmond
860 G. Higgins, et Julie D. Thompson. 2003. « Multiple Sequence Alignment with the

- Clustal Series of Programs ». *Nucleic Acids Research* 31 (13): 3497-3500.
<https://doi.org/10.1093/nar/gkg500>.
- Chesnais, Cédric B, Charlotte Bizet, Jérémy T Campillo, Wepnyu Y Njamnshi, Jean Bopda, Philippe Nwane, Sébastien D Pion, Alfred K Njamnshi, et Michel Boussinesq. 2020. « A Second Population-Based Cohort Study in Cameroon Confirms the Temporal Relationship Between Onchocerciasis and Epilepsy ». *Open Forum Infectious Diseases* 7 (6): ofaa206. <https://doi.org/10.1093/ofid/ofaa206>.
- Coffeng, Luc E., Wilma A. Stolk, Achim Hoerauf, Dik Habbema, Roel Bakker, Adrian D. Hopkins, et Sake J. De Vlas. 2014. « Elimination of African Onchocerciasis: Modeling the Impact of Increasing the Frequency of Ivermectin Mass Treatment ». Édité par James M. McCaw. *PLoS ONE* 9 (12): e115886. <https://doi.org/10.1371/journal.pone.0115886>.
- Colebunders, Robert, Jane Y Carter, Peter Claver Olore, Kai Puok, Samit Bhattacharyya, Sonia Menon, Gasim Abd-Elfarag, et al. 2018. « High Prevalence of Onchocerciasis-Associated Epilepsy in Villages in Maridi County, Republic of South Sudan: A Community-Based Survey ». *Seizure* 63 (décembre):93-101. <https://doi.org/10.1016/j.seizure.2018.11.004>.
- Coreil, Jeannine, et Gladys Mayard. 2009. « Indigenization of Illness Support Groups for Lymphatic Filariasis in Haiti ». In *Anthropology and Public Health*, édité par Robert A Hahn et Marcia Inborn, 2^e éd., 245-65. Oxford University Press New York. <https://doi.org/10.1093/acprof:oso/9780195374643.003.0010>.
- Coulibaly, Yaya I., Benoit Dembele, Abdallah A. Diallo, Ettie M. Lipner, Salif S. Doumbia, Siaka Y. Coulibaly, Siaka Konate, et al. 2009a. « A Randomized Trial of Doxycycline for *Mansonella Perstans* Infection ». *New England Journal of Medicine* 361 (15): 1448-58. <https://doi.org/10.1056/NEJMoa0900863>.
- . 2009b. « A Randomized Trial of Doxycycline for *Mansonella Perstans* Infection ». *The New England Journal of Medicine* 361 (15): 1448-58. <https://doi.org/10.1056/NEJMoa0900863>.
- Coulibaly, Yaya Ibrahim, Benoit Dembele, Abdallah Amadou Diallo, Sibylle Kristensen, Siaka Konate, Housseini Dolo, Ilo Dicko, et al. 2013. « Wuchereria Bancrofti Transmission Pattern in Southern Mali Prior to and Following the Institution of Mass Drug Administration ». *Parasites & Vectors* 6 (1): 247. <https://doi.org/10.1186/1756-3305-6-247>.

- Crawford, Katie E, Shannon M Hedtke, Stephen R Doyle, Annette C Kuesel, Samuel Armoo, Mike Y Osei-Atweneboana, et Warwick N Grant. 2024. « Genome-Based Tools for Onchocerciasis Elimination: Utility of the Mitochondrial Genome for Delineating Onchocerca Volvulus Transmission Zones ». *International Journal for Parasitology* 54 (3-4): 171-83. <https://doi.org/10.1016/j.ijpara.2023.11.002>.
- Cromwell, Elizabeth A, Chris A Schmidt, Kevin T Kwong, David M Pigott, Denise Mupfasoni, Gautam Biswas, Shreya Shirude, et al. 2020. « The Global Distribution of Lymphatic Filariasis, 2000–18: A Geospatial Analysis ». *The Lancet Global Health* 8 (9): e1186-94. [https://doi.org/10.1016/S2214-109X\(20\)30286-2](https://doi.org/10.1016/S2214-109X(20)30286-2).
- Davis, Emma L., Lisa J. Reimer, Lorenzo Pellis, et T. Deirdre Hollingsworth. 2019. « Evaluating the Evidence for Lymphatic Filariasis Elimination ». *Trends in Parasitology* 35 (11): 860-69. <https://doi.org/10.1016/j.pt.2019.08.003>.
- De Sole, G., J. Giese, F.M. Keita, et J. Remme. 1991. « Detailed Epidemiological Mapping of Three Onchocerciasis Foci in West Africa ». *Acta Tropica* 48 (3): 203-13. [https://doi.org/10.1016/0001-706X\(91\)90048-O](https://doi.org/10.1016/0001-706X(91)90048-O).
- Debrah, Alexander Yaw, Sabine Specht, Ute Klarmann-Schulz, Linda Batsa, Sabine Mand, Yeboah Marfo-Debrekyei, Rolf Fimmers, et al. 2015. « Doxycycline Leads to Sterility and Enhanced Killing of Female *Onchocerca Volvulus* Worms in an Area With Persistent Microfilaridemia After Repeated Ivermectin Treatment: A Randomized, Placebo-Controlled, Double-Blind Trial ». *Clinical Infectious Diseases* 61 (4): 517-26. <https://doi.org/10.1093/cid/civ363>.
- Debrah, Linda Batsa, Norman Nausch, Vera Serwaa Opoku, Wellington Owusu, Yusif Mubarik, Daniel Antwi Berko, Samuel Wanji, et al. 2017. « Epidemiology of *Mansonella perstans* in the middle belt of Ghana ». *Parasites & Vectors* 10 (janvier):15. <https://doi.org/10.1186/s13071-016-1960-0>.
- Demanou, M., S. D. Pion, et M. Boussinesq. 2001. « [Entomologic study of loaiasis transmission in the Lekie area (Cameroon)] ». *Bulletin De La Societe De Pathologie Exotique (1990)* 94 (4): 347-52.
- Dembélé, Pascal, Mady Cissoko, Adama Zan Diarra, Lassana Doumbia, Aïssata Koné, Mahamadou H. Magassa, Maissane Mehadji, Mahamadou A. Thera, et Stéphane Ranque. 2024. « Evaluation of the Performance of Rapid Diagnostic Tests for Malaria Diagnosis and Mapping of Different Plasmodium Species in Mali ». *International Journal of Environmental Research and Public Health* 21 (2): 228. <https://doi.org/10.3390/ijerph21020228>.

928 Deng, Miao-Han, Lan-Yi Zhong, Okanurak Kamolnetr, Yanin Limpanont, et Zhi-Yue Lv.
 929 2019. « Detection of Helminths by Loop-Mediated Isothermal Amplification Assay: A
 930 Review of Updated Technology and Future Outlook ». *Infectious Diseases of Poverty* 8
 931 (1): 20. <https://doi.org/10.1186/s40249-019-0530-z>.

932 Desjardins, Christopher A, Gustavo C Cerqueira, Jonathan M Goldberg, Julie C Dunning
 933 Hotopp, Brian J Haas, Jeremy Zucker, José M C Ribeiro, et al. 2013. « Genomics of
 934 Loa Loa, a Wolbachia-Free Filarial Parasite of Humans ». *Nature Genetics* 45 (5):
 935 495-500. <https://doi.org/10.1038/ng.2585>.

936 Desoubeaux, Guillaume, Adélaïde Chesnay, et Jacques Chandenier. 2020. « Approches
 937 diagnostiques des filarioses en France métropolitaine ». *Revue Francophone des*
 938 *Laboratoires* 2020 (519): 72-80. [https://doi.org/10.1016/S1773-035X\(20\)30066-6](https://doi.org/10.1016/S1773-035X(20)30066-6).

939 Devi, RRenuka, et V Raju. 2018. « Information Technology in Morbidity Management of
 940 Human Lymphatic Filariasis—A Promising Tool in Global Programme for Elimination
 941 of Lymphatic Filariasis ». *Journal of Vector Borne Diseases* 55 (1): 20.
 942 <https://doi.org/10.4103/0972-9062.234622>.

943 Diagne, Jean Pierre, Aly Mbara Ka, Joseph Matar Ndiaye, ABoubacry Sadikh Sow, Aissatou
 944 Maguatte Wane, et Paule Aida Ndoeye Roth. 2018. « MIGRATION SOUS CUTANEE
 945 PALPEBRALE D'UNE LOASE: A PROPOS D'UN CAS », n° 1, 4.
 946 https://www.revues-ufhb-ci.org/fichiers/FICHIR_ARTICLE_2511.pdf.

947 Diallo, S., M. Sarr, S. Diagne, et L. Konate. 1977. « Filariasis in the Nianing région" Senegal.
 948 Prevalence, clinical aspects, entomological findings. » *Medecine d'Afrique Noire* 24
 949 (4): 233-42. <https://www.cabdirect.org/cabdirect/abstract/19772902103>.

950 Díaz-Menéndez, Marta, Francesca Norman, Begoña Monge-Maillo, José Antonio Pérez-
 951 Molina, et Rogelio López-Vélez. 2011. « Las filariasis en la práctica clínica ». *Enfermedades Infecciosas y Microbiología Clínica* 29 (décembre):27-37.
 952 [https://doi.org/10.1016/S0213-005X\(11\)70041-6](https://doi.org/10.1016/S0213-005X(11)70041-6).

954 Diwakar, Deepak Kumar, Nikita Wadhwani, Nikhil Prasad, et Ashwani Gupta. 2018. « Primary
 955 Retroperitoneal Filariasis: A Common Disease of Tropics with Uncommon Presentation
 956 and Review of Literature ». *BMJ Case Reports*, septembre, bcr-2018-226217.
 957 <https://doi.org/10.1136/bcr-2018-226217>.

958 Dolo, Housseini, Yaya Ibrahim Coulibaly, Louise Kelly-Hope, Siaka Konate, Benoit Dembele,
 959 Siaka Yamoussa Coulibaly, Dramane Sanogo, et al. 2018. « Factors Associated with
 960 Wuchereria Bancrofti Microfilaremia in an Endemic Area of Mali ». *The American*

Journal of Tropical Medicine and Hygiene 98 (6): 1782-87.
<https://doi.org/10.4269/ajtmh.17-0902>.

Downes, B, et K Jacobsen. 2010. « A systematic review of the epidemiology of mansonelliasis ». *African Journal of Infectious Diseases* 4 (1).
<https://doi.org/10.4314/ajid.v4i1.55085>.

Downes, Barbara L., et Kathryn H. Jacobsen. 2010. « A Systematic Review of the Epidemiology of Mansonelliasis ». *African Journal of Infectious Diseases* 4 (1): 7-14.
<https://doi.org/10.4314/ajid.v4i1.55085>.

Drame, Papa M., Céline Montavon, Sébastien D. Pion, Joseph Kubofcik, Michael P. Fay, et Thomas B. Nutman. 2016. « Molecular Epidemiology of Blood-Borne Human Parasites in a Loa Loa-, Mansonella Perstans-, and Plasmodium Falciparum-Endemic Region of Cameroon ». *The American Journal of Tropical Medicine and Hygiene* 94 (6): 1301-8.
<https://doi.org/10.4269/ajtmh.15-0746>.

Duong, Thanh Hai, et Dominique Richard-Lenoble. 2008. « Diagnostic des parasitoses à parasites sanguicoles ». *Revue Francophone des Laboratoires* 2008 (399): 29-39.
[https://doi.org/10.1016/S1773-035X\(08\)70180-1](https://doi.org/10.1016/S1773-035X(08)70180-1).

Ehrens, Alexandra, Achim Hoerauf, et Marc P. Hübner. 2022. « Eosinophils in filarial infections: Inducers of protection or pathology? » *Frontiers in Immunology* 13 (octobre):983812. <https://doi.org/10.3389/fimmu.2022.983812>.

El Setouhy, Maged, Reda M. R. Ramzy, Ehab S. Ahmed, Amr M. Kandil, Omar Hussain, Hoda A. Farid, Hanan Helmy, et Gary J. Weil. 2004. « A Randomized Clinical Trial Comparing Single- and Multi-Dose Combination Therapy with Diethylcarbamazine and Albendazole for Treatment of Bancroftian Filariasis ». *The American Journal of Tropical Medicine and Hygiene* 70 (2): 191-96.

Emukah, Emmanuel, Lindsay J. Rakers, Barminas Kahansim, Emmanuel S. Miri, Bertram E. B. Nwoke, Emily Griswold, Yisa Saka, et al. 2018. « In Southern Nigeria Loa loa Blood Microfilaria Density is Very Low Even in Areas with High Prevalence of Loiasis: Results of a Survey Using the New LoaScope Technology ». *The American Journal of Tropical Medicine and Hygiene* 99 (1): 116-23. <https://doi.org/10.4269/ajtmh.18-0163>.

Erickson, Sara M., Edward K. Thomsen, John B. Keven, Naomi Vincent, Gussy Koimbu, Peter M. Siba, Bruce M. Christensen, et Lisa J. Reimer. 2013. « Mosquito-Parasite Interactions Can Shape Filariasis Transmission Dynamics and Impact Elimination Programs ». Édité par Jesus G. Valenzuela. *PLoS Neglected Tropical Diseases* 7 (9): e2433. <https://doi.org/10.1371/journal.pntd.0002433>.

- Famakinde, Damilare. 2018. « Mosquitoes and the Lymphatic Filarial Parasites: Research Trends and Budding Roadmaps to Future Disease Eradication ». *Tropical Medicine and Infectious Disease* 3 (1): 4. <https://doi.org/10.3390/tropicalmed3010004>.
- Faye, O., B. Faye, J. L. Ndiaye, R. C. K. Tine, D. Ndiaye, et A. Haydara Mbacké. 2009. « [Epidemiological study on lymphatic filariasis (*Wuchereria bancrofti*) in Sénégal] ». *Medecine Tropicale: Revue Du Corps De Sante Colonial* 69 (1): 101-2.
- Ferreira, Marcelo U., James Lee Crainey, et Federico G. Gobbi. 2023. « The Search for Better Treatment Strategies for Mansonellosis: An Expert Perspective ». *Expert Opinion on Pharmacotherapy* 24 (15): 1685-92. <https://doi.org/10.1080/14656566.2023.2240235>.
- Ferreira, Marcelo U., James Lee Crainey, et Sérgio L.B. Luz. 2021. « *Mansonella Ozzardi* ». *Trends in Parasitology* 37 (1): 90-91. <https://doi.org/10.1016/j.pt.2020.03.005>.
- Frédéric, NAHMIAS. 2018. « Haute Autorité de santé ».
- Gaillard, Colette Marie, Sebastien David Pion, Hadjira Hamou, Constant Sirima, Charlotte Bizet, Thomas Lemarcis, Jules Rodrigues, et al. 2020. « Detection of DNA of Filariae Closely Related to *Mansonella Perstans* in Faecal Samples from Wild Non-Human Primates from Cameroon and Gabon ». *Parasites & Vectors* 13 (1): 313. <https://doi.org/10.1186/s13071-020-04184-1>.
- Gardon, Jacques, Nathalie Gardon-Wendel, Demanga-Ngangue, Joseph Kamgno, Jean-Philippe Chippaux, et Michel Boussinesq. 1997. « Serious Reactions after Mass Treatment of Onchocerciasis with Ivermectin in an Area Endemic for Loa Loa Infection ». *The Lancet* 350 (9070): 18-22. [https://doi.org/10.1016/S0140-6736\(96\)11094-1](https://doi.org/10.1016/S0140-6736(96)11094-1).
- Geelhand De Merxem, D., J. N. Siewe Fodjo, S. Menon, A. Hotterbeekx, et R. Colebunders. 2019. « Nodding Syndrome Research, Lessons Learned from the NSETHIO Project ». *Global Mental Health* 6:e26. <https://doi.org/10.1017/gmh.2019.24>.
- Gentilini, M., J.M. Pinon, et M. Danis. 1973. « Diagnostic biologique des filarioses humaines ». *Médecine et Maladies Infectieuses* 3 (8-9): 351-53. [https://doi.org/10.1016/S0399-077X\(73\)80130-1](https://doi.org/10.1016/S0399-077X(73)80130-1).
- Gobbi, Federico, Anna Beltrame, Dora Buonfrate, Silvia Staffolani, Monica Degani, Maria Gobbo, Andrea Angheben, Stefania Marocco, et Zeno Bisoffi. 2017. « Imported Infections with *Mansonella perstans* Nematodes, Italy ». *Emerging Infectious Diseases* 23 (9): 1539-42. <https://doi.org/10.3201/eid2309.170263>.
- González, Mikel A., Erin Dilger, María M. Ronderos, Gustavo R. Spinelli, Orin Courtenay, et James G. C. Hamilton. 2020. « Significant Reduction in Abundance of Peridomestic

1029 Mosquitoes (Culicidae) and Culicoides Midges (Ceratopogonidae) after Chemical
 1030 Intervention in Western São Paulo, Brazil ». *Parasites & Vectors* 13 (1): 549.
 1031 <https://doi.org/10.1186/s13071-020-04427-1>.

1032 Gyasi, Michael E., Ogugua N. Okonkwo, et Koushik Tripathy. 2024. « Onchocerciasis ». In
 1033 *StatPearls. Treasure Island (FL): StatPearls Publishing.*
 1034 <http://www.ncbi.nlm.nih.gov/books/NBK559027/>.

1035 Haddix, Anne C., et Andrew Kestler. 2000. « Lymphatic Filariasis: Economic Aspects of the
 1036 Disease and Programmes for Its Elimination ». *Transactions of the Royal Society of*
 1037 *Tropical Medicine and Hygiene* 94 (6): 592-93. [https://doi.org/10.1016/S0035-](https://doi.org/10.1016/S0035-9203(00)90199-8)
 1038 [9203\(00\)90199-8](https://doi.org/10.1016/S0035-9203(00)90199-8).

1039 Harrup, Fabio Macchioni, Luca Chelucci, et Beatrice Torracca. 2016. « Advances in Control
 1040 Techniques for Culicoides and Future Prospects ». *Veterinaria Italiana* 52 (3): 247-64.
 1041 <https://doi.org/10.12834/VetIt.741.3602.3>.

1042 Hartley, Marie. 2021. « Onchocerciasis (river blindness) », 2021.
 1043 <https://dermnetnz.org/topics/onchocerciasis>.

1044 Herbeuval, H., R. Herbeuval, et J. Duheille. 1962. « [The passage, in the circulating blood, of
 1045 hematopoietic masses, erythroblastic islands, megakaryoblastic islands, mixed
 1046 islands] ». *Nouvelle Revue Francaise D'hematologie* 2:616-19.

1047 Herricks, Jennifer R., Peter J. Hotez, Valentine Wanga, Luc E. Coffeng, Juanita A. Haagsma,
 1048 María-Gloria Basáñez, Geoffrey Buckle, et al. 2017. « The Global Burden of Disease
 1049 Study 2013: What Does It Mean for the NTDs? » Édité par Xiao-Nong Zhou. *PLOS*
 1050 *Neglected Tropical Diseases* 11 (8): e0005424.
 1051 <https://doi.org/10.1371/journal.pntd.0005424>.

1052 Hotez, Peter J., et Aruna Kamath. 2009. « Neglected Tropical Diseases in Sub-Saharan Africa:
 1053 Review of Their Prevalence, Distribution, and Disease Burden ». Édité par Michael
 1054 Cappello. *PLoS Neglected Tropical Diseases* 3 (8): e412.
 1055 <https://doi.org/10.1371/journal.pntd.0000412>.

1056 Hotez, Peter J., David H. Molyneux, Alan Fenwick, Jacob Kumaresan, Sonia Ehrlich Sachs,
 1057 Jeffrey D. Sachs, et Lorenzo Savioli. 2007. « Control of Neglected Tropical Diseases ». *New*
 1058 *England Journal of Medicine* 357 (10): 1018-27.
 1059 <https://doi.org/10.1056/NEJMra064142>.

1060 Ibe, Ogochukwu, Obinna Onwujekwe, Benjamin Uzochukwu, Miriam Ajuba, et Paul
 1061 Okonkwo. 2015. « Exploring Consumer Perceptions and Economic Burden of
 1062 Onchocerciasis on Households in Enugu State, South-East Nigeria ». Édité par Samuel

1063 V. Scarpino. *PLOS Neglected Tropical Diseases* 9 (11): e0004231.
 1064 <https://doi.org/10.1371/journal.pntd.0004231>.

1065 Jacobsen, Kathryn H, Bailey C Andress, Elina A Bhagwat, Ciera A Bryant, Vipusha R
 1066 Chandrapu, Caroline G Desmonts, Tania M Matthews, Akeem Ogunkoya, Tristan J
 1067 Wheeler, et A'kayla S Williams. 2022. « A Call for Loiasis to Be Added to the WHO
 1068 List of Neglected Tropical Diseases ». *The Lancet Infectious Diseases* 22 (10):
 1069 e299-302. [https://doi.org/10.1016/S1473-3099\(22\)00064-0](https://doi.org/10.1016/S1473-3099(22)00064-0).

1070 Jain, Rajeev, Jern Y. Chen, Andrew R. Butcher, Robert Casson, et Dinesh Selva. 2008.
 1071 « Subconjunctival Loa Loa Worm ». *International Journal of Infectious Diseases* 12
 1072 (6): e133-35. <https://doi.org/10.1016/j.ijid.2008.03.001>.

1073 James, Spencer L, Degu Abate, Kalkidan Hassen Abate, Solomon M Abay, Cristiana Abbafati,
 1074 Nooshin Abbasi, Hedayat Abbastabar, et al. 2018. « Global, Regional, and National
 1075 Incidence, Prevalence, and Years Lived with Disability for 354 Diseases and Injuries
 1076 for 195 Countries and Territories, 1990–2017: A Systematic Analysis for the Global
 1077 Burden of Disease Study 2017 ». *The Lancet* 392 (10159): 1789-1858.
 1078 [https://doi.org/10.1016/S0140-6736\(18\)32279-7](https://doi.org/10.1016/S0140-6736(18)32279-7).

1079 Kamgno, Joseph, Sébastien D. Pion, Cédric B. Chesnais, Matthew H. Bakalar, Michael V.
 1080 D'Ambrosio, Charles D. Mackenzie, Hugues C. Nana-Djeunga, et al. 2017. « A Test-
 1081 and-Not-Treat Strategy for Onchocerciasis in *Loa Loa* –Endemic Areas ». *New England*
 1082 *Journal of Medicine* 377 (21): 2044-52. <https://doi.org/10.1056/NEJMoa1705026>.

1083 Katawa, Gnatoulma, Laura E. Layland, Alex Y. Debrah, Charlotte Von Horn, Linda Batsa,
 1084 Alexander Kwarteng, Sandra Arriens, et al. 2015. « Hyperreactive Onchocerciasis Is
 1085 Characterized by a Combination of Th17-Th2 Immune Responses and Reduced
 1086 Regulatory T Cells ». Édité par Benjamin L. Makepeace. *PLoS Neglected Tropical*
 1087 *Diseases* 9 (1): e3414. <https://doi.org/10.1371/journal.pntd.0003414>.

1088 Katiyar, D., et L. K. Singh. 2011. « Filariasis: Current Status, Treatment and Recent Advances
 1089 in Drug Development ». *Current Medicinal Chemistry* 18 (14): 2174-85.
 1090 <https://doi.org/10.2174/092986711795656234>.

1091 Keiser, Paul B., Yaya I. Coulibaly, Falaye Keita, Diakaridia Traore, Abdallah Diallo, Dapa A.
 1092 Diallo, Roshanak T. Semnani, et al. 2003. « Clinical Characteristics of Post-Treatment
 1093 Reactions to Ivermectin/Albendazole for *Wuchereria Bancrofti* in a Region Co-
 1094 Endemic for *Mansonella Perstans* ». *The American Journal of Tropical Medicine and*
 1095 *Hygiene* 69 (3): 331-35.

1096 Keiser, Paul B., Yaya Coulibaly, Joseph Kubofcik, Abdallah A. Diallo, Amy D. Klion, Sekou
 1097 F. Traoré, et Thomas B. Nutman. 2008. « Molecular Identification of Wolbachia from
 1098 the Filarial Nematode *Mansonella Perstans* ». *Molecular and Biochemical Parasitology*
 1099 160 (2): 123-28. <https://doi.org/10.1016/j.molbiopara.2008.04.012>.

1100 Keita, Mamadou Fodé. 1979. « Aspects épidémiologiques des filarioses à onchocerca volvulus,
 1101 wuchereria bancrofti et tetrapetatonema perstans ». Thesis, Ecole Nationale de
 1102 Médecine et de Pharmacie. <https://www.bibliosante.ml/handle/123456789/9605>.

1103 Kelly-Hope, Louise, Rossely Paulo, Brent Thomas, Miguel Brito, Thomas R. Unnasch, et
 1104 David Molyneux. 2017. « Loa Loa Vectors Chrysops Spp.: Perspectives on Research,
 1105 Distribution, Bionomics, and Implications for Elimination of Lymphatic Filariasis and
 1106 Onchocerciasis ». *Parasites & Vectors* 10 (1): 172. [https://doi.org/10.1186/s13071-017-](https://doi.org/10.1186/s13071-017-2103-y)
 1107 2103-y.

1108 Khan, A. M., P. Dutta, S. Das, A. K. Pathak, P. Sarmah, M. E. Hussain, et J. Mahanta. 2015.
 1109 « Microfilarial Periodicity of Wuchereria Bancrofti in Assam, Northeast India ». *Journal of Vector Borne Diseases* 52 (3): 208-12.

1110 Kima, A., K.T. Guiguemde, M. Serme, Z.C. Meda, R. Bougma, J.P. Djiatsa, C. Bougouma, et
 1111 F. Drabo. 2021. « Évaluation de la Transmission de la Filariose Lymphatique au
 1112 Burkina Faso: À Propos de 4 Districts Sanitaires ». *Meédecine Tropicale et Santéé*
 1113 *Internationale* 1 (2): mtsibulletin.n1.2021.83.
 1114 <https://doi.org/10.48327/mtsibulletin.n1.2021.83>.

1115 Knopp, Stefanie, Peter Steinmann, Christoph Hatz, Jennifer Keiser, et Jürg Utzinger. 2012.
 1116 « Nematode Infections »: *Infectious Disease Clinics of North America* 26 (2): 359-81.
 1117 <https://doi.org/10.1016/j.idc.2012.02.005>.

1118 Komlan, Kossi, Patrick S. Vossberg, Richard G. Gantim, Tchalim Solim, Francois Korbmacher,
 1119 Méba Banla, Koffi Padjoudoum, Potchoziou Karabou, Carsten Köhler, et Peter T.
 1120 Soboslay. 2018. « Onchocerca Volvulus Infection and Serological Prevalence, Ocular
 1121 Onchocerciasis and Parasite Transmission in Northern and Central Togo after Decades
 1122 of Simulium Damnosum s.l. Vector Control and Mass Drug Administration of
 1123 Ivermectin ». Édité par Matthew C. Freeman. *PLOS Neglected Tropical Diseases* 12
 1124 (3): e0006312. <https://doi.org/10.1371/journal.pntd.0006312>.

1125 Kwarteng, Alexander, Samuel Terkper Ahuno, et Freda Osei Akoto. 2016. « Killing Filarial
 1126 Nematode Parasites: Role of Treatment Options and Host Immune Response ». *Infectious Diseases of Poverty* 5 (1): 86. <https://doi.org/10.1186/s40249-016-0183-0>.

1129 Laidoudi, Younes, Jean-Lou Marie, Djamel Tahir, Stéphanie Watier-Grillot, Oleg
1130 Mediannikov, et Bernard Davoust. 2020. « Detection of Canine Vector-Borne Filariasis
1131 and Their Wolbachia Endosymbionts in French Guiana ». *Microorganisms* 8 (5): 770.
1132 <https://doi.org/10.3390/microorganisms8050770>.

1133 Lima, Nathália F., Cecilia A. Veggiani Aybar, María J. Dantur Juri, et Marcelo U. Ferreira.
1134 2016. « *Mansonella Ozzardi* : A Neglected New World Filarial Nematode ». *Pathogens*
1135 *and Global Health* 110 (3): 97-107. <https://doi.org/10.1080/20477724.2016.1190544>.

1136 Lustigman, Sara, Alexandra Grote, et Elodie Ghedin. 2017. « The Role of “omics” in the Quest
1137 to Eliminate Human Filariasis ». Édité par Judd L. Walson. *PLOS Neglected Tropical*
1138 *Diseases* 11 (4): e0005464. <https://doi.org/10.1371/journal.pntd.0005464>.

1139 M. Boussinesq. 2007. « Actualités thérapeutiques dans les filarioses – Recent advances in the
1140 treatment of filariases ». *mise au point*, n° 2, 55.

1141 Mandal, Debasish, Dipankar Roy, D. K. Bera, et Buddhadeb Manna. 2012. « Occurrence of
1142 Gravid Loa Loa in Subconjunctival Space of Man: A Case Report from West Bengal,
1143 India ». *Journal of Parasitic Diseases*, juillet. [https://doi.org/10.1007/s12639-012-](https://doi.org/10.1007/s12639-012-0130-3)
1144 [0130-3](https://doi.org/10.1007/s12639-012-0130-3).

1145 Manguin, S., M.J. Bangs, J. Pothikasikorn, et T. Chareonviriyaphap. 2010. « Review on Global
1146 Co-Transmission of Human Plasmodium Species and Wuchereria Bancrofti by
1147 Anopheles Mosquitoes ». *Infection, Genetics and Evolution* 10 (2): 159-77.
1148 <https://doi.org/10.1016/j.meegid.2009.11.014>.

1149 McGarry, Helen F, Ken Pfarr, Gill Egerton, Achim Hoerauf, Jean-Paul Akue, Peter Enyong,
1150 Samuel Wanji, et al. 2003. « [No title found] ». *Filaria Journal* 2 (1): 9.
1151 <https://doi.org/10.1186/1475-2883-2-9>.

1152 Meda, N., M.A. Daboue, W.P. Djigumde, C. Meda, et S. Konate/Ouedraogo. 2009. « 571
1153 Manifestations oculaires de la loase ». *Journal Français d’Ophtalmologie* 32
1154 (avril):1S173. [https://doi.org/10.1016/S0181-5512\(09\)73695-2](https://doi.org/10.1016/S0181-5512(09)73695-2).

1155 Medeiros, Jansen Fernandes, James Lee Crainey, Felipe Arley Costa Pessoa, et Sérgio Luis
1156 Bessa Luz. 2017. « Mansonelliasis ». In *Arthropod Borne Diseases*, édité par Carlos
1157 Brisola Marcondes, 405-26. Cham: Springer International Publishing.
1158 https://doi.org/10.1007/978-3-319-13884-8_26.

1159 Mediannikov, O., et S. Ranque. 2018a. « Mansonellosis, the Most Neglected Human
1160 Filariasis ». *New Microbes and New Infections* 26 (novembre):S19-22.
1161 <https://doi.org/10.1016/j.nmni.2018.08.016>.

1162 ———. 2018b. « Mansonellosis, the Most Neglected Human Filariasis ». *New Microbes and*
1163 *New Infections* 26 (novembre):S19-22. <https://doi.org/10.1016/j.nmni.2018.08.016>.

1164 Melrose, Wayne D. 2002. « Lymphatic Filariasis: New Insights into an Old Disease ». *International Journal for Parasitology* 32 (8): 947-60. [https://doi.org/10.1016/S0020-](https://doi.org/10.1016/S0020-7519(02)00062-0)
1165 [7519\(02\)00062-0](https://doi.org/10.1016/S0020-7519(02)00062-0).

1167 Mendoza, Natalia, Alexa Li, Amber Gill, et Stephen Tying. 2009. « Filariasis: Diagnosis and
1168 Treatment ». *Dermatologic Therapy* 22 (6): 475-90. [https://doi.org/10.1111/j.1529-](https://doi.org/10.1111/j.1529-8019.2009.01271.x)
1169 [8019.2009.01271.x](https://doi.org/10.1111/j.1529-8019.2009.01271.x).

1170 Métais, Alice, Sophie Michalak, et Audrey Rousseau. 2021. « Albendazole-Related Loa Loa
1171 Encephalopathy ». *IDCases* 23:e01033. <https://doi.org/10.1016/j.idcr.2020.e01033>.

1172 Metzger, Wolfram Gottfried, et Benjamin Mordmüller. 2014. « Loa Loa—Does It Deserve to
1173 Be Neglected? » *The Lancet Infectious Diseases* 14 (4): 353-57.
1174 [https://doi.org/10.1016/S1473-3099\(13\)70263-9](https://doi.org/10.1016/S1473-3099(13)70263-9).

1175 Molyneux, David H., et Mark J. Taylor. 2001. « Current Status and Future Prospects of the
1176 Global Lymphatic Filariasis Programme »: *Current Opinion in Infectious Diseases* 14
1177 (2): 155-59. <https://doi.org/10.1097/00001432-200104000-00008>.

1178 Mourembou, Gaël, Florence Fenollar, Jean Bernard Lekana-Douki, Angelique Ndjoi
1179 Mbiguino, Sydney Maghendji Nzondo, Pierre Blaise Matsiegui, Rella Zoleko Manego,
1180 et al. 2015a. « Mansonella, Including a Potential New Species, as Common Parasites in
1181 Children in Gabon ». Édité par Dunstan Mukoko. *PLOS Neglected Tropical Diseases* 9
1182 (10): e0004155. <https://doi.org/10.1371/journal.pntd.0004155>.

1183 ———. 2015b. « Mansonella, including a Potential New Species, as Common Parasites in
1184 Children in Gabon ». *PLoS Neglected Tropical Diseases* 9 (10): e0004155.
1185 <https://doi.org/10.1371/journal.pntd.0004155>.

1186 Ngwewondo, Adela, Ivan Scandale, et Sabine Specht. 2021. « Onchocerciasis Drug
1187 Development: From Preclinical Models to Humans ». *Parasitology Research* 120 (12):
1188 3939-64. <https://doi.org/10.1007/s00436-021-07307-4>.

1189 Nizeyimana, Jean Baptiste. 2021. « ÉTUDE DES CONDITIONS DE DEPISTAGE
1190 CLINIQUE ET LES EFFETS PSYCHO SOCIOECONOMIQUE DES PERSONNES
1191 PORTEUSES DE L'ELEPHANTIASIS DES JAMBES. CAS DE
1192 L'ARRONDISSEMENT DE GUIDIGUIS CAS DE LA REGION DE L'EXTREME-
1193 NORD CAMEROUN DE 2017-2020 Par » 26 (03).

1194 Njim, Tsi, Joel Mbigha Ngum, et Leopold Ndemnge Aminde. 2015. « Cutaneous
1195 Onchocerciasis in Dumbu, a Pastoral Area in the North-West Region of Cameroon:

Diagnostic Challenge and Socio-Economic Implications ». *Pan African Medical Journal* 22. <https://doi.org/10.11604/pamj.2015.22.298.7707>.

Njoku, Chidiebere, Lawrence Nwankwo, Masayo Ozaki, Andrew Obasi, Chinyere N. Ukaga, Emmanuel Emukah, Kenrick Nwodu, et al. 2013. « Community-Wide Distribution of Long-Lasting Insecticidal Nets Can Halt Transmission of Lymphatic Filariasis in Southeastern Nigeria ». *The American Journal of Tropical Medicine and Hygiene* 89 (3): 578-87. <https://doi.org/10.4269/ajtmh.12-0775>.

Noroës, Joaquim, David Addiss, Agnaldo Cedenho, Jose Figueredo-Silvan, Guilherme Lima, et Gerusa Dreyer. 2003. « Pathogenesis of Filarial Hydrocele: Risk Associated with Intrascrotal Nodules caused by Death of Adult *Wuchereria Bancrofti* ». *Transactions of the Royal Society of Tropical Medicine and Hygiene* 97 (5): 561-66. [https://doi.org/10.1016/S0035-9203\(03\)80029-9](https://doi.org/10.1016/S0035-9203(03)80029-9).

Nutman, Thomas B. 2013. « Insights into the Pathogenesis of Disease in Human Lymphatic Filariasis ». *Lymphatic Research and Biology* 11 (3): 144-48. <https://doi.org/10.1089/lrb.2013.0021>.

O'Hanlon, Simon J., Hannah C. Slater, Robert A. Cheke, Boakye A. Boatın, Luc E. Coffeng, Sébastien D. S. Pion, Michel Boussinesq, Honorat G. M. Zouré, Wilma A. Stolk, et María-Gloria Basáñez. 2016. « Model-Based Geostatistical Mapping of the Prevalence of *Onchocerca Volvulus* in West Africa ». Édité par Ricardo J. Soares Magalhaes. *PLOS Neglected Tropical Diseases* 10 (1): e0004328. <https://doi.org/10.1371/journal.pntd.0004328>.

Ojurongbe, Olusola, Akeem Abiodun Akindele, Monsuru Adebayo Adeleke, Matthew Oyebo Oyebo, Samuel Adeyinka Adedokun, Josephine Folashade Ojo, Callistus Adewale Akinleye, Oloyede Samuel Bolaji, Olusegun Adelowo Adefioye, et Oluwaseyi Adegboyega Adeyeba. 2015. « Co-endemicity of Loiasis and Onchocerciasis in Rain Forest Communities in Southwestern Nigeria ». *PLOS Neglected Tropical Diseases* 9 (3): e0003633. <https://doi.org/10.1371/journal.pntd.0003633>.

« OMS MALI Rapport annuel 2018.pdf ». s. d. Consulté le 24 juillet 2024. <https://www.afro.who.int/sites/default/files/2019-01/OMS%20MALI%20Rapport%20annuel%202018.pdf>.

Opoku, Millicent, Corrado Minetti, Worlasi D. Kartey-Attipoe, Sampson Otoo, Joseph Otchere, Bruno Gomes, Dziedzom K. De Souza, et Lisa J. Reimer. 2018. « An Assessment of Mosquito Collection Techniques for Xenomonitoring of Anopheline-Transmitted

1229 Lymphatic Filariasis in Ghana ». *Parasitology* 145 (13): 1783-91.
 1230 <https://doi.org/10.1017/S0031182018000938>.

1231 Ottesen, Eric A. 2000. « Editorial: The Global Programme to Eliminate Lymphatic Filariasis ». *Tropical Medicine & International Health* 5 (9): 591-94. <https://doi.org/10.1046/j.1365-3156.2000.00620.x>.

1232 ———. 2006. « Lymphatic Filariasis: Treatment, Control and Elimination ». In *Advances in Parasitology*, 61:395-441. Elsevier. [https://doi.org/10.1016/S0065-308X\(05\)61010-X](https://doi.org/10.1016/S0065-308X(05)61010-X).

1233 Ouedraogo, Amina Nomtongo, Emile Bégnouriti Somda, Fagnima Traoré, Muriel Sidnoma
 1234 Ouédraogo, Gilbert Patrice Tapsoba, Angèle Ouangre/Ouédraogo, Salamata Bara, et al.
 1235 2016. « Impact du traitement de masse de la filariose lymphatique par l'albendazole-
 1236 ivermectine en zone de savane : cas de la région de l'Est du Burkina ». *HEALTH SCIENCES AND DISEASE* 17 (4). <http://hsd-fmsb.org/index.php/hsd/article/view/731>.

1237 Pampiglione, S., G. Canestri Trotti, et F. Rivasi. 1995. « Human Dirofilariasis Due to
 1238 Dirofilaria (Noctiella) Repens: A Review of World Literature ». *Parassitologia* 37
 1239 (2-3): 149-93.

1240 Pérez-Molina, José Antonio, Marta Díaz-Menéndez, Ana Pérez-Ayala, Federico Ferrere,
 1241 Begoña Monje, Francesca Norman, et Rogelio López-Vélez. 2010. « Tratamiento de las
 1242 enfermedades causadas por parásitos ». *Enfermedades Infecciosas y Microbiología Clínica* 28 (1): 44-59. <https://doi.org/10.1016/j.eimc.2009.11.003>.

1243 Phillips, Richard O., Michael Frimpong, Fred S. Sarfo, Birte Kretschmer, Marcus Beissner,
 1244 Alexander Debrah, Yaw Ampem-Amoako, et al. 2014. « Infection with *Mansonella*
 1245 *perstans* Nematodes in Buruli Ulcer Patients, Ghana ». *Emerging Infectious Diseases*
 1246 20 (6): 1000-1003. <https://doi.org/10.3201/eid2006.131501>.

1247 Pion, S. D. S., J. A. N. Filipe, J. Kamgno, J. Gardon, M.-G. Basáñez, et M. Boussinesq. 2006.
 1248 « Microfilarial Distribution of Loa Loa in the Human Host: Population Dynamics and
 1249 Epidemiological Implications ». *Parasitology* 133 (01): 101.
 1250 <https://doi.org/10.1017/S0031182006000035>.

1251 Pion, Sébastien Ds, Hugues Nana-Djeunga, Yannick Niamsi-Emalio, Cédric B Chesnais, Hugo
 1252 Deléglise, Charles Mackenzie, Wilma Stolk, et al. 2020. « Implications for Annual
 1253 Retesting after a Test-and-Not-Treat Strategy for Onchocerciasis Elimination in Areas
 1254 Co-Endemic with Loa Loa Infection: An Observational Cohort Study ». *The Lancet Infectious Diseases* 20 (1): 102-9. [https://doi.org/10.1016/S1473-3099\(19\)30554-7](https://doi.org/10.1016/S1473-3099(19)30554-7).

1255 Poole, Catherine B., Zhiru Li, Andy Alhassan, Dylan Guelig, Steven Diesburg, Nathan A.
 1256 Tanner, Yinhua Zhang, et al. 2017. « Colorimetric Tests for Diagnosis of Filarial

1263 Infection and Vector Surveillance Using Non-Instrumented Nucleic Acid Loop-
 1264 Mediated Isothermal Amplification (NINA-LAMP) ». Édité par Matty Knight. *PLOS*
 1265 *ONE* 12 (2): e0169011. <https://doi.org/10.1371/journal.pone.0169011>.

1266 Porter, Suzanne, et Adrian Hopkins. 2015. « Élimination de l'onchocercose et de la filariose
 1267 lymphatique » 12.

1268 Puente, Sabino, Mar Lago, Mercedes Subirats, Ismael Sanz-Esteban, Marta Arsuaga, Belén
 1269 Vicente, Montserrat Alonso-Sardon, Moncef Belhassen-Garcia, et Antonio Muro. 2020.
 1270 « Imported Mansonella perstans infection in Spain ». *Infectious Diseases of Poverty* 9
 1271 (1): 105. <https://doi.org/10.1186/s40249-020-00729-9>.

1272 Ramaiah, K. D., et Eric A. Ottesen. 2014. « Progress and Impact of 13 Years of the Global
 1273 Programme to Eliminate Lymphatic Filariasis on Reducing the Burden of Filarial
 1274 Disease ». Édité par Moses Bockarie. *PLoS Neglected Tropical Diseases* 8 (11): e3319.
 1275 <https://doi.org/10.1371/journal.pntd.0003319>.

1276 Ramharter, Michael, Joanna Butler, Ghyslain Mombo-Ngoma, Tamara Nordmann, Saskia
 1277 Dede Davi, et Rella Zoleko Manego. 2023. « The African Eye Worm: Current
 1278 Understanding of the Epidemiology, Clinical Disease, and Treatment of Loiasis ». *The*
 1279 *Lancet Infectious Diseases*, octobre, S1473309923004383.
 1280 [https://doi.org/10.1016/S1473-3099\(23\)00438-3](https://doi.org/10.1016/S1473-3099(23)00438-3).

1281 Rebollo, Maria P., et Moses J. Bockarie. 2017. « Can Lymphatic Filariasis Be Eliminated by
 1282 2020? » *Trends in Parasitology* 33 (2): 83-92. <https://doi.org/10.1016/j.pt.2016.09.009>.

1283 Richardson, Eugene T., Robert Luo, Doran L. Fink, Thomas B. Nutman, John K. Geisse, et
 1284 Michele Barry. 2012. « Transient Facial Swellings in a Patient With a Remote African
 1285 Travel History ». *Journal of Travel Medicine* 19 (3): 183-85.
 1286 <https://doi.org/10.1111/j.1708-8305.2012.00612.x>.

1287 Risch, Frederic, Manuel Ritter, Achim Hoerauf, et Marc P. Hübner. 2021. « Human Filariasis—
 1288 Contributions of the Litomosoides Sigmodontis and Acanthocheilonema Viteae Animal
 1289 Model ». *Parasitology Research* 120 (12): 4125-43. [https://doi.org/10.1007/s00436-](https://doi.org/10.1007/s00436-020-07026-2)
 1290 [020-07026-2](https://doi.org/10.1007/s00436-020-07026-2).

1291 Robert, Vincent, E.H. Ndiaye, Nil Rahola, Gilbert Le Goff, Philippe Boussès, D. Diallo,
 1292 Voahirana Le Goff, Lauriane Mariamé, et M. Diallo. 2022. « Clés dichotomiques
 1293 illustrées d'identification des femelles et des larves de moustiques (Diptera : Culicidae)
 1294 du Burkina Faso, Cap-Vert, Gambie, Mali, Mauritanie, Niger, Sénégal et Tchad », 181
 1295 p. <https://doi.org/10.23708/FDI:010084866>.

- Rodi, Miriam, Caspar Gross, Thaisa Lucas Sandri, Lilith Berner, Marina Marcet-Houben, Ersoy Kocak, Michaela Pogoda, et al. 2023a. « Whole genome analysis of two sympatric human *Mansonella*: *Mansonella perstans* and *Mansonella* sp “DEUX” ». *Frontiers in Cellular and Infection Microbiology* 13 (avril):1159814. <https://doi.org/10.3389/fcimb.2023.1159814>.
- . 2023b. « Whole Genome Analysis of Two Sympatric Human *Mansonella*: *Mansonella Perstans* and *Mansonella* Sp “DEUX” ». *Frontiers in Cellular and Infection Microbiology* 13:1159814. <https://doi.org/10.3389/fcimb.2023.1159814>.
- Sandri, Thaisa Lucas, Andrea Kreidenweiss, Simon Cavallo, David Weber, Sascha Juhas, Miriam Rodi, Tamirat Gebru Woldearegai, et al. 2021. « Molecular Epidemiology of *Mansonella* Species in Gabon ». *The Journal of Infectious Diseases* 223 (2): 287-96. <https://doi.org/10.1093/infdis/jiaa670>.
- Sangare, Moussa Brema, Yaya Ibrahim Coulibaly, Siaka Yamoussa Coulibaly, Michel Emmanuel Coulibaly, Bourama Traore, Ilo Dicko, Ibrahim Moussa Sissoko, et al. 2018. « A cross-sectional study of the filarial and *Leishmania* co-endemicity in two ecologically distinct settings in Mali ». *Parasites & Vectors* 11 (1): 18. <https://doi.org/10.1186/s13071-017-2531-8>.
- Sawadogo, Pm. 2019. « >Onchocercose au Burkina Faso : une endémie à transmission vectorielle en voie de réémergence ». *Bulletin de la société de pathologie exotique* 112 (5): 260-74. <https://doi.org/10.3166/bspe-2020-0106>.
- Shelley, Aj, et S Coscarón. 2001. « Simuliid blackflies (Diptera: Simuliidae) and ceratopogonid midges (Diptera: Ceratopogonidae) as vectors of *Mansonella ozzardi* (Nematoda: Onchocercidae) in northern Argentina ». *Memórias do Instituto Oswaldo Cruz* 96 (4): 451-58. <https://doi.org/10.1590/S0074-02762001000400003>.
- Shenoy, R. K. 2008. « Clinical and Pathological Aspects of Filarial Lymphedema and Its Management ». *The Korean Journal of Parasitology* 46 (3): 119. <https://doi.org/10.3347/kjp.2008.46.3.119>.
- Simón, Fernando, Mar Siles-Lucas, Rodrigo Morchón, Javier González-Miguel, Isabel Mellado, Elena Carretón, et Jose Alberto Montoya-Alonso. 2012. « Human and Animal Dirofilariasis: The Emergence of a Zoonotic Mosaic ». *Clinical Microbiology Reviews* 25 (3): 507-44. <https://doi.org/10.1128/CMR.00012-12>.
- Simonsen, Paul E., Yahya A. Derua, Stephen M. Magesa, Erling M. Pedersen, Anna-Sofie Stensgaard, Mwelecele N. Malecela, et William N. Kisinza. 2014. « Lymphatic filariasis control in Tanga Region, Tanzania: status after eight rounds of mass drug

administration ». *Parasites & Vectors* 7 (1): 507. <https://doi.org/10.1186/s13071-014-0507-5>.

Simonsen, Paul E., Ambrose W. Onapa, et Santa Maria Asio. 2011a. « Mansonella Perstans Filariasis in Africa ». *Acta Tropica*, The Diagnostics and Control of Neglected Tropical Helminth Diseases, 120 (septembre):S109-20. <https://doi.org/10.1016/j.actatropica.2010.01.014>.

———. 2011b. « Mansonella Perstans Filariasis in Africa ». *Acta Tropica* 120 Suppl 1 (septembre):S109-120. <https://doi.org/10.1016/j.actatropica.2010.01.014>.

Small, Scott T., Daniel J. Tisch, et Peter A. Zimmerman. 2014. « Molecular Epidemiology, Phylogeny and Evolution of the Filarial Nematode Wuchereria Bancrofti ». *Infection, Genetics and Evolution* 28 (décembre):33-43. <https://doi.org/10.1016/j.meegid.2014.08.018>.

Supali, Taniawati, Is S. Ismid, Heri Wibowo, Yenny Djuardi, Esther Majawati, Praba Ginanjar, et Peter Fischer. 2006. « Estimation of the Prevalence of Lymphatic Filariasis by a Pool Screen PCR Assay Using Blood Spots Collected on Filter Paper ». *Transactions of the Royal Society of Tropical Medicine and Hygiene* 100 (8): 753-59. <https://doi.org/10.1016/j.trstmh.2005.10.005>.

Tahita, Marc Christian, Thuy-Huong Ta-Tang, Bérénger Kaboré, Raquel Capote-Morales, Irene Molina De La Fuente, Raquel Cruces, Hamidou Ilboudo, et al. 2024. « Molecular Detection of Wuchereria Bancrofti, Loa Loa and Mansonella Perstans from Dried Blood Spots Taken from Pregnant Women in Rural Burkina Faso ». *African Journal of Parasitology, Mycology and Entomology* 1 (2): 1. <https://doi.org/10.35995/ajpme2010002>.

Tamarozzi, Francesca, Paola Rodari, Joaquín Salas-Coronas, Emmanuel Bottieau, Fernando Salvador, Manuel Jesús Soriano-Pérez, María Isabel Cabeza-Barrera, et al. 2022. « A Large Case Series of Travel-Related *Mansonella Perstans* (Vector-Borne Filarial Nematode): A TropNet Study in Europe ». *Journal of Travel Medicine*, avril, taac048. <https://doi.org/10.1093/jtm/taac048>.

Ta-Tang, Thuy-Huong, James Crainey, Rory J. Post, Sergio L.B. Luz, et José Rubio. 2018. « Mansonellosis: Current Perspectives ». *Research and Reports in Tropical Medicine* Volume 9 (janvier):9-24. <https://doi.org/10.2147/RRTM.S125750>.

Ta-Tang, Thuy-Huong, Sergio L B Luz, James L Crainey, et José M Rubio. 2021a. « An Overview of the Management of Mansonellosis ». *Research and Reports in Tropical Medicine* 12 (mai):93-105. <https://doi.org/10.2147/RRTM.S274684>.

- 1364 Ta-Tang, Thuy-Huong, Sergio Lb Luz, James L Crainey, et José M Rubio. 2021b. « An
1365 Overview of the Management of Mansonellosis ». *Research and Reports in Tropical*
1366 *Medicine* Volume 12 (mai):93-105. <https://doi.org/10.2147/RRTM.S274684>.
- 1367 Tatuene, J. Kamtchum, R. Gounoue Fotsing, T. Nkoa, JB Tchateng Mbougua, HC Nana
1368 Djeunga, J. Bopda, et Joseph Kamgno. 2014a. « Epidemiology of Loa Loa and
1369 Mansonella Perstans Filariasis in the Akonolinga Health District, Centre Region,
1370 Cameroon. » *HEALTH SCIENCES AND DISEASE* 15 (1). [https://www.hsd-](https://www.hsd-fmsb.org/index.php/hsd/article/view/297)
1371 [fmsb.org/index.php/hsd/article/view/297](https://www.hsd-fmsb.org/index.php/hsd/article/view/297).
- 1372 ———. 2014b. « Epidemiology of Loa Loa and Mansonella Perstans Filariasis in the
1373 Akonolinga Health District, Centre Region, Cameroon. » *HEALTH SCIENCES AND*
1374 *DISEASE* 15 (1). <https://doi.org/10.5281/hsd.v15i1.297>.
- 1375 Tavares da Silva, Lucyane Bastos, James Lee Crainey, Túllio Romão Ribeiro da Silva, Uziel
1376 Ferreira Suwa, Ana Carolina Paulo Vicente, Jansen Fernandes de Medeiros, Felipe
1377 Arley Costa Pessoa, et Sérgio Luiz Bessa Luz. 2017. « Molecular Verification of New
1378 World Mansonella perstans Parasitemias ». *Emerging Infectious Diseases* 23 (3):
1379 545-47. <https://doi.org/10.3201/eid2303.161159>.
- 1380 Taylor, Mark J, Achim Hoerauf, et Moses Bockarie. 2010. « Lymphatic Filariasis and
1381 Onchocerciasis ». *The Lancet* 376 (9747): 1175-85. [https://doi.org/10.1016/S0140-](https://doi.org/10.1016/S0140-6736(10)60586-7)
1382 [6736\(10\)60586-7](https://doi.org/10.1016/S0140-6736(10)60586-7).
- 1383 Thylefors, B. 2004. « Eliminating Onchocerciasis as a Public Health Problem ». *Tropical*
1384 *Medicine & International Health* 9 (4). [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-3156.2004.01226.x)
1385 [3156.2004.01226.x](https://doi.org/10.1111/j.1365-3156.2004.01226.x).
- 1386 Unnasch, Thomas R, Allison Golden, Vitaliano Cama, et Paul T Cantey. 2018. « Diagnostics
1387 for Onchocerciasis in the Era of Elimination ». *International Health* 10 (suppl_1):
1388 i20-26. <https://doi.org/10.1093/inthealth/ihx047>.
- 1389 Varenne, F., J. Fillaux, M. Porterie, J. Soler, M. Cassagne, et V. Soler. 2016. « Loase sous-
1390 conjonctivale : à propos d'un cas ». *Journal Français d'Ophtalmologie* 39 (8): e193-94.
1391 <https://doi.org/10.1016/j.jfo.2015.08.018>.
- 1392 Vinkeles Melchers, Natalie V. S., Sarah Mollenkopf, Robert Colebunders, Michael Edlinger,
1393 Luc E. Coffeng, Julia Irani, Trésor Zola, et al. 2018. « Burden of Onchocerciasis-
1394 Associated Epilepsy: First Estimates and Research Priorities ». *Infectious Diseases of*
1395 *Poverty* 7 (1). <https://doi.org/10.1186/s40249-018-0481-9>.
- 1396 Vinkeles Melchers, Natalie V. S., Wilma A. Stolk, Welmoed Van Loon, Belén Pedrique, Roel
1397 Bakker, Michele E. Murdoch, Sake J. De Vlas, et Luc E. Coffeng. 2021. « The Burden

1398 of Skin Disease and Eye Disease Due to Onchocerciasis in Countries Formerly under
1399 the African Programme for Onchocerciasis Control Mandate for 1990, 2020, and
1400 2030 ». Édité par Alberto Novaes Ramos. *PLOS Neglected Tropical Diseases* 15 (7):
1401 e0009604. <https://doi.org/10.1371/journal.pntd.0009604>.

1402 Vlaminck, Johnny, Peter U. Fischer, et Gary J. Weil. 2015. « Diagnostic Tools for
1403 Onchocerciasis Elimination Programs ». *Trends in Parasitology* 31 (11): 571-82.
1404 <https://doi.org/10.1016/j.pt.2015.06.007>.

1405 Volpicelli, Lorenzo, Maurizio De Angelis, Alessandra Morano, Elisa Biliotti, Cristiana Franchi,
1406 Simona Gabrielli, Simonetta Mattiucci, Carlo Di Bonaventura, et Gloria Taliani. 2020.
1407 « Encephalopathy in a Patient with Loiasis Treated with Albendazole: A Case Report ». *Parasitology International* 75 (avril):102006.
1408 <https://doi.org/10.1016/j.parint.2019.102006>.

1410 Walther, Michael, et Ralph Muller. 2003. « Diagnosis of Human Filariases (except
1411 Onchocerciasis) ». In *Advances in Parasitology*, 53:149-93. Elsevier.
1412 [https://doi.org/10.1016/S0065-308X\(03\)53004-4](https://doi.org/10.1016/S0065-308X(03)53004-4).

1413 Wanji, S., N. Tendongfor, M. Esum, S. Ndindeng, et P. Enyong. 2003. « Epidemiology of
1414 Concomitant Infections Due to Loa Loa, Mansonella Perstans, and Onchocerca
1415 Volvulus in Rain Forest Villages of Cameroon ». *Medical Microbiology and Immunology* 192 (1): 15-21. <https://doi.org/10.1007/s00430-002-0154-x>.

1417 Whittaker, Charles, Martin Walker, Sébastien D.S. Pion, Cédric B. Chesnais, Michel
1418 Boussinesq, et María-Gloria Basáñez. 2018. « The Population Biology and
1419 Transmission Dynamics of Loa Loa ». *Trends in Parasitology* 34 (4): 335-50.
1420 <https://doi.org/10.1016/j.pt.2017.12.003>.

1421 World Health Organization (WHO). 2020. « Onchocerciasis factsheet. Geneva: World Health
1422 Organization », 16. <https://www.who.int/news-room/factsheets/detail/onchocerciasis>.

1423 World Health Organization. 2011. « Agir pour réduire l'impact mondial des maladies tropicales
1424 négligées : premier rapport de l'OMS sur les maladies tropicales négligées ». *Working
1425 to overcome the global impact of neglected tropical diseases: first WHO report on
1426 neglected tropical diseases*. <https://iris.who.int/handle/10665/44694>.

1427 ———. 2022. « Global programme to eliminate lymphatic filariasis: progress report, 2021 »,
1428 2022. <http://www.who.int/wer>.

1429 World Health Organization. *Bench Aids for the Diagnosis of Filarial Infection*. Geneves:
1430 1997. <https://iris.who.int/handle/10665/37156> Consulté Le 2024-07-11. s. d.

- World Health Organization (WHO). 2021. « Global programme to eliminate lymphatic filariasis: progress report, 2020 ». *Weekly Epidemiological Record = Relevé épidémiologique hebdomadaire* 96 (41): 497-508. CORREA.
- World Health Organization (WHO). 2022. « Plan stratégique national de lutte contre les maladies tropicales négligées 2022 - 2025 Sénégal », 144. <https://espen.afro.who.int/system/files/content/resources/Senegal%20Plan%20National%20stratégie%20MTN%202022-2025.pdf>.
- World Health Organization (WHO). 2023. « Lymphatic Filariasis ». 1 juin 2023. <https://www.who.int/fr/news-room/fact-sheets/detail/lymphatic-filariasis>.
- Wynd, Shona. 2007. « Understanding the community impact of lymphatic filariasis: a review of the sociocultural literature ». *Bulletin of the World Health Organization* 85 (6): 493-98. <https://doi.org/10.2471/BLT.06.031047>.
- Yokoly, Firmain N., Julien B. Z. Zahouli, Aboulaye Méite, Millicent Opoku, Bernard L. Kouassi, Dziedzom K. De Souza, Moses Bockarie, et Benjamin G. Koudou. 2020. « Low Transmission of Wuchereria Bancrofti in Cross-Border Districts of Côte d'Ivoire: A Great Step towards Lymphatic Filariasis Elimination in West Africa ». Édité par Justin V. Remais. *PLOS ONE* 15 (4): e0231541. <https://doi.org/10.1371/journal.pone.0231541>.
- Zouré, Honorat Gustave Marie, Samuel Wanji, Mounkaïla Noma, Uche Veronica Amazigo, Peter J. Diggle, Afework Hailemariam Tekle, et Jan H. F. Remme. 2011. « The Geographic Distribution of Loa Loa in Africa: Results of Large-Scale Implementation of the Rapid Assessment Procedure for Loiasis (RAPLOA) ». Édité par Giovanna Raso. *PLoS Neglected Tropical Diseases* 5 (6): e1210. <https://doi.org/10.1371/journal.pntd.0001210>.

Funding

The study was funded by the Mediterranean Infection Foundation.

Informed Consent Statement

Not applicable.

Data Availability Statement

All data are provided in the manuscript.

Conflicts of Interest

The authors declare that they have no competing interests.

1463 **Abbreviations**

1464	NTD	Neglected Tropical Diseases
1465	FTS	Filariasis Test Strip
1466	DNA	Deoxyribonucleic Acid
1467	ELISA	Enzyme-Linked Immuno Assay
1468	MDA	Mass Drug Administration
1469	PCR	Polymerase Chain Reaction
1470	LAMP	Loop-mediated amplification
1471	GE	thick drop
1472	IFI	Immunofluorescence
1473	COES	Co-Electrosyneresis
1474	IELP	Immuno-electrophoresis
1475	ICT	Immunochromatographic
1476	LIPS	Luciferase Immunoprecipitation System
1477	WHO	World Health Organization
1478	BCE	Bloodless skin biopsy
1479	MGG	May Grunwald Giemsa
1480	GPELF	Global program for the elimination of lymphatic filariasis
1481	OCP	Onchocerciasis in West Africa
1482	APOC	Africa onchocerciasis control program
1483	DEET	NN-diethyl-3-methylbenzamide
1484		

PARTIE 2

Diagnosis of filariasis in patients and identification of potential vectors

Mansonella perstans in southern Senegal

ARTICLE 2

Elephantiasis and *Mansonella perstans* in Senegal (West Africa)

Manthia DIALLO, Hubert BASSENE, Cheikh SOKHNA, Georges DIATTA, Mousse SAMBOU, Jean-Noel Magloire MANGA, Ousseynou BA, Florence FENOLLAR, Oleg MEDIANNIKOV

Article soumis au journal: The American Journal of Tropical Medicine and Hygiene



Elephantiasis and *Mansonella perstans* in West Africa

Journal:	<i>American Journal of Tropical Medicine & Hygiene</i>
Manuscript ID	Draft
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	n/a
Complete List of Authors:	DIALLO, Manthia; Aix-Marseille Universite, biologie Bassene, Hubert; URMITE, UMR IRD 198 Sokhna, Cheikh; URMITE, UMR IRD 198 Diatta, Georges; IRD, VITROME, ex-URMITE, HEALTH SAMBOU, Masse; IRD Senegal, Dakar Manga, Noel; Hôpital de la Paix, Service de maladies infectieuses et tropicales BA, Ousseynou; Ministry of Healthcare and Social Action, Cap Skirring, Senegal , Cap Skirring Fenollar, Florence; Faculté de Médecine, Unité des Rickettsies Mediannikov, Oleg; Aix Marseille Université, URMITE UMR 6236, IRD 198; IRD, URMITE UMR198
Key Words:	Elephantiasis, <i>Mansonella perstans</i> , rapid diagnostic tests, Mosquitoes, biting midges

SCHOLARONE™
Manuscripts

Résumé

Les filarioses sont des maladies parasitaires fréquentes qui touchent des millions de personnes dans les régions tropicales et subtropicales engendrées par différentes espèces de vers ronds connus sous le nom de filaires. L'éléphantiasis est une affection clinique, grave due à l'obstruction du système lymphatique par les vers adultes. *Mansonella perstans* est plus négligé car il est considéré comme bénigne causant principalement des infections asymptomatiques. L'infection par l'une de ces maladies se produit lorsque les parasites filaires responsables sont transmis à l'Homme par des arthropodes hématophages des genres Moustiques et Culicoïdes. Le but de cette étude était de rechercher l'étiologie des cas d'éléphantiasis, d'explorer la faisabilité de l'utilisation des cassettes tests de diagnostic rapide (TDR) négatifs pour le paludisme et d'identifier si les arthropodes hématophages sont vecteurs potentiels de la filariose au Sénégal.


La collecte des échantillons a été réalisée au Sud du Sénégal qui est endémique par la filariose. Nous avons analysé 23 échantillons de sang des patients atteints d'éléphantiasis résidant à Kolda et Oussouye en 2021, 279 échantillons de sang de sujets fébriles à Mlomp entre 2018 - 2020 et 3714 cassettes TDR *Plasmodium falciparum* (*P.f*) négatives collectées dans les postes et centres de santé de 2020 à 2021. En plus, 254 moustiques collectés à Kolda, Oussouye et Sédhiou en 2021 à l'aide de pièges lumineux CDC, OVI et faune matinale résiduelle et 1233 moustiques collectés en 2007 et 2012, et 991 spécimens de culicoides ont été analysés.

La présence de *M. perstans* a été détectée chez 8,7 % (2/23) des patients malades et chez 4,3 % (12/279) des sujets fébriles de Mlomp. Parmi les 3714 cassettes TDR négatives *P.f* testées, 0,24 % (9/3714) se sont révélées positives.

Nos résultats montrent la présence de *M. perstans* dans les TDR, chez les sujets fébriles et chez les individus atteints d'éléphantiasis. Aucun arthropode n'a été trouvé infecter par *M. perstans*. Il est nécessaire de poursuivre la recherche du vecteur responsable de la filariose à *M. perstans* dans les sites endémiques du sud du Sénégal.

Keywords: Elephantiasis, *Mansonella perstans*, TDR, Arthropodes, filarioses, qPCR, Sénégal

Résumé graphique du projet



Abstract

Background: Filariases are parasitic diseases that infect millions of people living mostly in less developed areas, mainly in West Africa. Infection occurs when the filarial larvae are transmitted to humans by mosquitoes or other dipterans. Elephantiasis is a clinical condition, often severe, usually associated with mosquito-transmitted filarial worms of *Wuchereria* or *Brugia* genera. *Mansonella perstans* is considered as clinically mild, mainly asymptomatic filariasis and not usually associated with elephantiasis. The aim of this study was to investigate the etiology and epidemiology of elephantiasis cases and to screen the filaria hosting in population in Senegal.

Methods: This study was carried out in July-November and December 2021 in endemic sites in southern Senegal. We analyzed 23 blood samples from elephantiasis patients residing in Kolda and Oussouye in 2021, 279 blood samples from febrile subjects in Mlomp examined at the dispensary between 2018 - 2020 and 3714 nitro-cellulose strips from malaria rapid diagnostic tests collected in health posts and centres from 2020 to 2021. 254 mosquitoes and 991 *Culicoides* biting midges were collected in residences of elephantiasis patients in Kolda, Oussouye and Sédhiou in 2021 using light traps CDC, OVI and residual morning fauna. In addition, 1233 mosquitoes collected in other regions of Senegal in 2007 and 2012 were analyzed.

Results: The presence of *Mansonella perstans* was detected in 8.7% (2/23) of ill patients and 4.3% (12/279) of febrile subjects from Mlomp. Among the 3714 negative *P. falciparum* RDT cassettes tested, 0.24% (9/3714) were found to be positive. None of the mosquitoes and *Culicoides* tested were infected with *M. perstans*, but other probably animal filarial species were detected.

Conclusion: Our results show the presence of only *M. perstans* in both elephantiasis and febrile patients in Senegal, no other filarial nematodes were detected. No arthropods were infected with human pathogen filaria. It is necessary to continue studies of etiology of elephantiasis in Senegal and the search for the vector responsible for *M. perstans* filariasis in endemic sites in southern Senegal.

Keywords: Elephantiasis, *Mansonella perstans*, RDT, mosquitoes, biting midges, filariasis, qPCR, Senegal

INTRODUCTION

The World Health Organization (WHO) estimates that about 1.7 billion people suffer from neglected tropical disease and that 49 African countries bear 40% of the global burden. Elephantiasis or lymphatic filariasis is one of the oldest and most debilitating neglected tropical disease affecting more than 51.4 million people worldwide in 72 endemic countries [1]. According to the WHO, in 2023, more than 882 million people in 44 countries are at risk of lymphatic filariasis [2]. Since the beginning of the Global Program for the Elimination of Lymphatic Filariasis (GPELF) in 2000, the goal was to stop the parasite transmission by mass drug administration, management of morbidity and prevention of disabilities [3]. Between 2007 and 2019, 34 African countries had lymphatic filariasis, with 26 countries of them reporting lymphoedema and hydrocele with 148,114 and 137,696 cases, respectively. Lymphatic filariasis results from infestation by *Wuchereria bancrofti* and *Brugia* sp. [4]. These are transmitted to humans by the bite of mosquitoes belonging to the genera *Culex*, *Anopheles*, *Mansonia* and *Aedes* [1]. Hydrocele, lymphedema and elephantiasis are the visible chronic consequences that do not appear until later in life [1,5]. Lymphatic filariasis causes an apparent alteration of the lymphatic system and can lead to an abnormal increase in the volume of certain parts of the body, resulting in pain, severe disability, and social stigma.

Elephantiasis has been historically reported in Senegal. Published in 1821 detailed description of the frigate “Méduse” shipwreck mentioned the existence of elephantiasis and probably onchocerciasis in Senegal [6]. Interesting, for the best of our knowledge, *W. bancrofti* was neither characterized nor even detected by molecular methods in Senegal. Based on very few studies, it is considered to be present in certain areas, but diagnostic relies only on microscopic studies [7–10].

Recent studies on children in Uganda have revealed co-infections between *W. bancrofti* and *Mansonella perstans*, the agent responsible for mansonellosis, in 33 individuals (0.3%) in six schools. [11].

Despite the wide distribution of mansonellosis, very few studies have investigated its epidemiology and clinical picture [12]. Simonsen estimated that 114 million people were infected. In Africa alone, 581 million people in 33 countries are at risk of *M. perstans* [13]. Prevalence is often very high in endemic areas and children are not spared [12]. Mansonellosis is globally caused by three species of filarial nematodes, generally producing chronic human microfilaremia and with variable geographical distribution: (1) *M. perstans*, generally found in

sub-Saharan Africa, but also in South America; (2) *Mansonella streptocerca* endemic to sub-Saharan Africa; (3) *Mansonella ozzardii*, found exclusively in America and the Caribbean islands [13–15]. Recently, genetic studies showed that at least one other genetic variant *Mansonella* sp. DEUX infect humans in Africa (Gabon) [16–18]. Whole genome analysis clearly showed that *Mansonella* sp. “DEUX” is a separate species, [16] but it is yet not clear if it is a new, yet undescribed species or one of old ape-associated *Mansonella* species that were described only morphologically [19]. These are *Mansonella vanhoofi* from bonobos and gorillas, *Mansonella rodhaini* from bonobos and chimpanzees, *Mansonella lopeensis*, *Mansonella leopoldi* and *Mansonella gorilla*, all three from gorillas.

The recent study in rural Senegal (Kedougou) showed an overall prevalence of 14.5% of *M. perstans* in febrile and afebrile subjects [20]. Although infection with *M. perstans* is considered a low pathogenic, often asymptomatic, various clinical manifestations have been associated with its presence, such as asthenia, skin swelling, itching, ocular or palpebral pruritus, rashes, angioedema, headache, arthralgia, abdominal pain, chest pain, pericarditis, pleurisy, neurological manifestations, visual impairment, inflammatory granulomatous nodules surrounding dead adult worms, and fever in some cases [12,20,21]. Eosinophilia can also be observed.

M. perstans is considered to be transmitted by *Culicoides* biting midges [22]: *Culicoides milnei* and *Culicoides austani* in Cameroon [23,24] and *Culicoides fulvithorax* and *Culicoides grahamii* in Nigeria [25,26]. However, in Senegal, no *Culicoides* vector species has yet been identified [20].

The aim of this study carried out in rural Senegal was to determine the etiology of cases of elephantiasis, to demonstrate the usefulness of nitro-cellulose strips of malaria rapid diagnostic tests (RDTs) to the screening of filariasis, and to identify whether mosquitoes could be potential vectors of *M. perstans* in the regions studied.

Methods

Ethical statement

This study was approved by the National Committee on Health Research Ethics of Senegal (CNERS) and the Research Ethics Committee (CER) of UCAD under the numbers SEN21/81 and O75/2021/CER/UCAD. Local authorizations were also obtained by the chief medical officers of the districts of Kolda, Sedhiou, Kedougou and Ziguinchor (Oussouye).

Individual written informed consent was obtained from adult participants and, for minors, from their parents or legal guardians, and all were informed of the procedure. Data was recorded and stored to preserve patient anonymity.

Study areas

The study was conducted in 4 regions (Ziguinchor, Kolda, Sedhiou and Kedougou) in Senegal (**Fig. 1**). In the Ziguinchor region, the study was conducted in the department of Oussouye (12°29'06"N - 16°32'49"E), in the villages of Kagnout, Diakene diola, Djibelene, Niambalang and Diakene ouolof. Oussouye is one of the 3 departments of the Ziguinchor region, located in Lower Casamance on the left bank of the Casamance River. The average temperature in Ziguinchor is 26.0°C in the coldest month (January) and 29.8°C in the hottest month (June). Rainfall reaches 1,400 millimeters per year, making it quite abundant. During the least rainy months (February, March, April, December), it is 0 mm, while during the wettest month (August), it reaches 481 mm [27].

In the Kolda region (12°53'00"N - 14°57'00"E), the study was conducted in the villages of Bagadadji, Demba korbali, Sare pathe, Missira kamarang, Guiro yero alpha, Sare sandiong, Missara Ousmane and Kampissa. Kolda is in southeastern Senegal on the southern border of Gambia. Its climate is Sudano Guinean with an average temperature of 28°C, with a minimum of 18.3°C and a maximum of 37.8°C. Annual rainfall is 1185 mm [28]

In the Sedhiou region (12°42'29"N – 15°33'25"E), the study was conducted in the villages of Tambanaba and Badiary. Sedhiou is bounded to the north by Gambia, to the south by the republics of Guinea and Guinea Bissau, to the east by the region of Kolda, to the west by the region of Ziguinchor. Temperatures range from 25 - 30°C with an average precipitation of 1000 mm per year [29].

In the Kedougou region (12°33'N – 12°11'E), the study was conducted in the localities of Mako, Ndiormi, Fadiga, Dalaba and Tomboronkoto. Kedougou is limited to the West and North by the Tambacounda region, to the South-East by the Republic of Guinea and Mali. The rainfall is very important which makes it one of the rainiest regions of Senegal with at least 1300 mm per year. The climate is Sudano-Guinean, and temperatures are relatively high ranging from a maximum of 42°C to a minimum of 21°C [30].




Figure 1. Localization of villages and health posts in study areas (QGIS Desktop 2.2.0).

Samples collection

Capillary blood sampling in patients with elephantiasis

Capillary blood samples were collected in July 2021 in patients with symptoms of elephantiasis and residing in 8 localities of Kolda (Bagadadji, Demba korbali, Sare pathe, Missira kamarang, Guiro yero alpha, Sare sandiong, Missara Ousmane and Kampissa) and 4 localities of Oussouye (Kagnout, Diakene diola, Djibelene, Niambalang and Diakene ouolof). (Table 1) A questionnaire designed to collect the patient's personal and clinical information was completed by the team's doctors prior to sample collection. Wherever possible, two capillary blood samples were taken from patients by finger prick at 10 a.m. (day) and 10 p.m. (night) in EDTA tubes. The samples were then stored in a cooler at 4°C and transferred to Dakar (Senegal's capital) under the same conditions.

Table 1: Clinical data on certain cases observed in participants in the study areas

									qPCR Pan Filaria		qPCR M. perstans	
Sites	Villages	Codes	Sex	Ages	Symptoms	Duration of illness (YEARS)	Hospitalization	Specific treatment	Diurnal	Nocturnal	Diurnal	Nocturnal
Kolda	Bagadadji	KL 1	F	68	Leg edema with recurrent inflammations, pruritus, vomiting, headache	~ 40	YES	Extencillin 1.2 MUT, mycostere 1 application, Local care	0	0	0	0
	Missira kamarang	KL 2	M	71	Hydrocele, vomiting, headache, recurrent inflammation	~ 18	NO	NO	0	0	0	0
	Missira ousmane	KL 3	M	58	Leg/foot edema, pruritus, headache	25	NO	Betadine dermique, Mycooster pomade, Fleming (Amoxy Acide Clavulanique)	0	0	0	0
	Kampissa	KL 4	M	67	Leg edema with recurrent inflammation, pain in the liver area, muscle pain	~ 60	YES	Extencillin, Amoxicillin, Acide clavulanique, Permaganate, SAT, VAT, Mycooster	0	0	0	0
	Demba korbali	KL 5	M	76	Leg/foot edema, periorbital edema, bilateral inguinal hernia	55	NO	Flucanazole, Mycooster, Vaselinesalicyle	31,08	28,13	32,58	32,08
	Saré Sandiong	KL 6	F	~ 95	Leg edema, recurrent erysipelas	~ 5	YES	NS*	0	0	0	0
	Bagadadji	KL 7	M	51	Leg/foot edema with recurrent infection, pruritus, omitting, headache	25	NO	NO (Surgery recommended)	0	0	0	0
	Guïro yero alpha	KL 8	F	70	Edema of groin area-Pruritus-Muscle Pain-Headache-Arthralgia	50	NO	Anti-fungal cream	0	0	0	0
	Saré Pathé	KL 9	F	65	Leg edema, muscle pain, headache, arthralgia, recurrent erysipelas	> 20	NO	NO	0	0	0	0
	Saré Pathé	KL 10	F	65	Quincke's edema, periorbital edema, arm/hand edema with recurrent inflammations, muscle pain, pruritus, stomach pain	More 40	YES	Hospital treatment in Gambia	0	0	0	0
	Saré Pathé	KL 11	F	90	Quincke's edema, hypogastric pain, myalgia, pruritis, arthralgie, lower limb swelling with recurrent inflammation	> 60	NO	NO	0	0	0	0
Oussouye	Kagnout Ebrouaye	KL 12	F	69	Leg edema, pruritus, stomach pain, arthralgia, itching, frequent erysipelas	> 40	NO	NO	0	0	0	0
	Kagnout Ebrouaye	KL 13	M	79	Leg/foot edema, periorbital edema, myalgia, pruritus, arthralgia, erysipelas	8	YES	NS*	0	26,73	0	31,14
	Diakène Diola	KL 14	F	55	Leg edema, muscle pain, pruritus, headache, arthralgia, erysipelas sometimes	35	NR	NS*	0	0	0	0
	Diakène Diola	KL 15	M	70	Leg edema, periorbital edema, myalgia, stomachache, arthralgia, headache, erysipelas	8	NO	1 time Ivermectin / Albendazol	0	0	0	0
	Diakène Diola	KL 16	F	61	Muscle pain-Arthralgia-Phlebitis odor	35	NO	2 times Ivermectin / Albendazol	0	0	0	0
	Djibetène / Mlomp	KL 17	F	67	Leg swelling, muscle pain, arthralgia	birth	NO	NO	0	0	0	0
	Diakène	KL 18	M	76	Leg edema, muscle pain, pruritus, arthralgia, cough for 7 years	1	NO	NO	0	0	0	0

	Diakène Diola	KL 19	M	64	Leg edema, muscle pain, pruritus, arthralgia, loss of sensation in the feet	3	NO	2 times mass treatment with Ivecmectin	0	0	0	0
	Niambalang	KL 20	M	66	Leg/foot edema, myalgia and arthralgia	28	NO	NO	0	0	0	0
	Niambalang	KL 21	M	57	Hydrocele, periorbital edema, muscle pain, pruritis	29	NR	NS*	0	0	0	0
	Niambalang	KL 22	M	76	Hydrocele, periorbital edema, headache	28	NR	NS*	0	0	0	0
	Diakène ouolof	KL 23	M	61	Leg edema, myalgia, pruritus, headache, erysipelas	41	NO	NO	0	0	0	0

*not specified

Capillary blood sampling in febrile patients attending the Mlomp health post

Capillary blood samples were taken daily from 2018 to 2020 in febrile patients consulting in the health post of Mlomp. A fever was defined by an axillary temperature of $\geq 37.5^{\circ}\text{C}$. The samples were kept at 4°C and then the nucleic acids were extracted in the laboratory of the Mlomp dispensary.

Negative RDT samples for *P. falciparum* in targeted stations

With the agreement of the National Malaria Control Programme (PNLP), all RDT cassettes from malaria-negative patients attending 18 health facilities were analysed. RDTs were therefore sorted at each site, and those that were negative were placed in plastic bags at room temperature. Once at the laboratory, each sample was assigned an internal code to preserve patient anonymity.

Arthropods collection and identification

In addition to Residual Morning Fauna (RMF), two types of light traps were used: Center for Diseases Control (CDC) modified (Bio Quip Products, Compton, CA, USA) and Onderspoort Veterinary Institute (OVI) developed by (Onderstepoort Veterinary Institute, Pretoria, South Africa). The RMF allowed arthropods to collect at rest in the rooms. The light traps were used in July, November and December 2021 in 12 villages in the districts of Kolda, Sedhiou and Oussouye. In the households where patients with elephantiasis lived, we placed one inside the patient's room that slept under a mosquito net and another outside. The traps were set at dusk at 6 p.m and removed in the morning at 8 a.m for two consecutive days. All collected arthropods were sorted, morphologically identified using dichotomic keys [31] and placed in individual 1.5 ml Eppendorf tubes containing silica gel [32,33] and carded cotton. Once they arrived at the laboratory, the mosquitoes were dissected, the head, thorax and abdomen were placed in an Eppendorf tube for filarial research, the legs and wings in the original tube. *Culicoides* species were not dissected, they were identified morphologically using identification keys based primarily on the characters described in the identification keys [34,35].

DNA extraction

According to the manufacturer's instructions, samples were extracted using different protocols:

1. EZ1®DNA tissue kit (Qiagen, Hilden, Germany) for blood samples collected from elephantiasis patients, and mosquitoes and culicoides captured in patients' homes in 2021,

2. E.Z.N.A® tissue DNA kit (Omega Bio-Tek, Norcross, GA, USA), blood samples from febrile Mlomp subjects.

3. 2% cetyltrimethylammonium bromide (CTAB) method to extract DNA from blood deposits on nitrocellulose membrane from negative TDRs [36] and mosquitoes collected in 2007 and 2012 in Oussouye and Kedougou.

Molecular screening of filariae by real-time quantitative PCR

Real-time quantitative PCR (qPCR) assays were used to screen for the presence of filariae. The 28S gene (pan-filarial 28S qPCR system) was first targeted to screen for the presence of all filarial species [37].

Then, when the sample was positive, the internal transcribed spacer 1 (ITS) of *M. perstans* was specifically targeted [20] as well as cytochrome c oxidase I mitochondrial subunit I (COX1) (Table 2).

Table 2. Primers and probes used in this study.

PCR types	Target sequences	Name of Primers and Probes	Sequences (5'-3')	Ref
qPCR	ITS	M. p-F	5'-AGGATCATTAACGAGCTTCC-3'	[20]
		M. p-R	5'-CGAATATCACCGTTAATTCAGT-3'	
		M. p-P	6-FAM-TTCACTTTTATTTAGCAACATGCA-TAMRA	
	28S	qFil-28S-F	5'-TTGTTTGAGATTGCAGCCCA-3'	[37]
		qFil-28S-R	5'-GTTTCCATCTCAGCGGTTTC-3'	
		qFil-28S-P	6-FAM-CAAGTACCGTGAGGGAAAGT-TAMRA	
	COX1(nuclear copy)	Nfil - F	5'-TACAGCGAGTCAACCAGAGG-3'	(this study)
		Nfil - R	5'-TGAAGAATCGGCCAAAACAGG-3'	
		Nfil - P	6-FAM-AATATGCGGTCTGTTGCTGC-TAMRA	
Sequencing	ITS1	ITS1-F	5'-GGTGAACCTGCGGAAGGATC-3'	[17]
		ITS1-R	5'-CTCAATGCGTCTGCAATTCGC-3'	
	5S	5S-S2	5'-GTTAAGCAACGTTGGGCCTGG-3'	
		5S-S16	5'-TTGACAGATCGGACGAGATG-3'	[38]
	COI	COLint-F	5'-TGATTGGTGGTTTTGGTAA-3'	
		COLint-R	5'-ATAAGTACGAGTATCAATATC-3'	[37]
	12S	Fwd-12S-Nem-1	5'-AAGTTTGATTTTGGTTTTGGTTG-3'	
		1Rwd-12S-Nem-681	5'-CCATTGACGGATGTTTGTA-3'	
		R-2198	5'-TASACTTCAGGGTGSCCRAARAAYCA-3'	

Amplification was performed using a CFX96 thermocycler (BioRad Laboratories, Foster City, CA, USA) in a final reaction volume of 20 µl, containing 10 µl of Master Mix Roche (Eurogentec, Seraing, Belgium), 0.5 µL of each primer at a concentration of 20 µM, 0.5 µL of the probe at a concentration of 5 µM, 0.5 µL of UDG, 3 µL of water DNase/RNase free and 5 µL of the DNA sample to be tested. The amplification program consisted of an initial cleaning step of 2 minutes at 50°C followed by a denaturation of 5 minutes at 95°C, then 40 denaturation cycles at 95°C for 5 seconds and a hybridization and elongation at 60°C for 30 seconds. For each series, negative (the mix) and positive controls were used. All samples positive for pan-filarial 28S qPCR system and for *M. perstans* and Nfil qPCR assays were sequenced to confirm.

Sequencing

Standard PCRs targeting COX1, ITS1, 5S and 12S genes using the primers listed in (Table 2) were performed using a thermal cycler (Applied Biosystems, 2720, Foster City, USA) and AmpliTaq Gold® 360 Master Mix (Applied Biosystems, Waltham, MA, USA). The amplification protocols consisted of a 15-minute incubation step at 95°C, 40 cycles (30 seconds at 95°C, 30 seconds at the hybridization temperature depending on the target gene, one minute at 72°C elongation and a final step of extension from 5 minutes to 72°C. Migration of the amplicons was carried out on a 2% agarose gel for 25 to 30 minutes at 180V. The bands were visualized under UV light. PCR products were purified using NucleoFast 96 PCR plates (Macherey Nagel, Hoerd, France) according to the manufacturer's instructions. Then, they were sequenced using the Big Dye Terminator Cycle sequencing kit (Perkin Elmer Applied Biosystems, Foster City, CA, USA) with an automated ABI sequencer (Applied Biosystems). The resulting sequences were assembled and analyzed using Chromas Pro v.1.77 software (Technelysium Pty. Ltd, Tewantin, Australia) in order to compare them to the reference sequences of filarial species available on the National Center for Biotechnology Information (NCBI) online database (<http://blast.ncbi.nlm.nih.gov>) through a Basic Local Alignment Search Tool (BLAST) search. We have deduced evolutionary relationships between wireline types identified by phylogenetic analysis using the neighbor junction method by Molecular Evolutionary Genetics Analysis MEGA version 7 (<https://www.megasoftware.net/>).

Statistical analysis

In the absence of up-to-date data on the prevalence of *M. perstans*, the sample size for screening of RDT needed to estimate the prevalence of the disease in each site at 5% of its true value, and with a 95% confidence interval was determined by adopting a proportion (p) of

14.5% for the Kedougou area and 50% (maximum prevalence) for the other areas, to calculate sample sizes. Sample size (n) was calculated using the following formula with sample size to be determined (n), $z = 1.96$; interval width (2δ), last known prevalence 14.5%

$$n = (z1 - \alpha/\delta)^2 p (1 - p)$$

Data were entered into a Microsoft Excel (version 13) spreadsheet.

Results

Selected patients with elephantiasis

In 2021, according to information obtained from the head nurse, 13 villages were visited, 8 in Kolda and 5 in the department of Oussouye. Patients were selected and sampled according to their local presence, with 11 patients living in the Kolda region and 12 patients in the department of Oussouye. Capillary blood samples were collected twice (at day and at night) from 23 patients with symptoms of elephantiasis (**fig. 2**) living in the regions of Kolda (11) and Oussouye (12). In Kolda region, 9 patients had leg or arm lymphedema (right or left) swelling and 2 cases of hydrocele. In Oussouye, 8 patients had leg or arm lymphedema (right or left) swelling, 3 cases hydroceles and 1 case who had lymphedema of both legs and hydrocele. Patient ages ranged from 51 to 95 years. The male/female sex ratio was 1.3% (13/10). These patients complained of symptoms such as groin pain, muscle pain, pruritus, headaches, arthralgia, recurrent erysipelas and pain in the liver area and others. All 23 patients were sampled during the day between 9 and 10 a.m., while only 14 were sampled at night between 10 and 11 p.m.



KL 9



KL 11



KL 12



KL 8



KL 4



KL 5

Fig. 2 Patients with clinical manifestations of elephantiasis in the study areas (Kolda and Oussouye). Among the study participants, **(KL8)** Swollen left arm, **(KL 9,4)** lymphedema affecting one leg (right and left), and **(KL 11,12,5)** lymphedema of both legs.

At the Mlomp dispensary, 279 capillary blood samples were collected from acute febrile patients between 2018 - 2020. The sampling has been performed within the framework of diagnostic activities. A total of 3,714 malaria-negative RDTs were retrieved from health facilities in the study sites and analyzed according to the sample size calculations, including 762 (20.5%) in Kedougou, 762 (20.5%) in Kolda, 863 (23.3%) in Oussouye and 1,327 (35.7%) in Sedhiou. The detailed numbers of RDTs tested per health post and center are shown in **Table 3**.

Table 3. RDTs cassettes collected and prevalence by study area.

Study areas	Sanitary facilities	Number of RDTs collected	Number of RDTs tested	Pos qfil 28S (%)	Pos <i>M. perstans</i> (%)	Prevalence of <i>M. perstans</i> by area
Kedougou	Mako*	675	124	2 (1.6%)	1 (0.08%)	1.0%
	Ndiormi§	1676	307	5 (1.6%)	5 (1.6%)	
	Fadiga*	443	81	0	0	
	Dalaba*	580	106	0	0	
	Tomboronkoto*	786	144	0	0	
Kolda	Bagadadji*	922	252	1 (0.39%)	1 (0.39%)	0.1%
	Sare Bilaly*	299	81	0	0	
	Sare Kemo*	1,571	429	0	0	
Oussouye	Oussouye§	1,743	352	1 (0.28%)	1 (0.28%)	0.1%
	Carouate*	258	253	0	0	
	Louda Ouoloff*	55	55	0	0	
	Diakene Djola*	154	54	0	0	
	Mlomp*	145	149	0	0	
Sedhiou	Diana Malary*	1,492	300	1 (0.33%)	1 (0.33%)	0.2%
	Oudoucar*	1,614	238	1 (0.4%)	1 (0.4%)	
	Dembo Coly*	904	300	1 (0.33%)	1 (0.3%)	
	Djende*	354	300	0	0	
	Dianna Ba*	207	189	1 (0.5%)	0	
Total		13,878	3,714	13 (0.35%)	11 (0.30%)	

* Health post

§ Health center

Arthropods

Overall, 254 mosquitoes were collected and analyzed for filariae in 2021 in 11 villages (6 in Kolda region [Missira ousmane, Missira mamadou, Demba korbali, Sare sandiong,

Bagadadji and Missira kamarang], 3 in Oussouye region [Diakene diola, Niambalang and Kagnout] and 2 in Sedhiou region [Tambanaba, Badiary]. Using dichotomous keys [31], the following proportions were determined: 35% *Anopheles gambiae* s.l. (89/254), 5.1% *Anopheles funestus* (13/254), 1.1% *Anopheles rufipes* (3/254), 0.3% *Anopheles squamosus* (1/254), 1.1% *Culex* sp (3/254), 55.9% *Mansonia uniformis* (142/254) and 1.5% *Mansonia africana* (4/254). The same traps also captured 991 biting midges of *Culicoides* genus. Using the identification keys from Cornet et al, all midges were identified *C. fulvithorax*, *C. imicola*, *C. milnei*, *C. moreli*, *C. enderleini* and *C. oxystoma*.

In addition, 1,233 mosquitoes collected using CDC traps were selected for molecular analysis of filariae: Kedougou 36% (444/1233) were *Aedes* collected in 2007 and 64% (789/1233) represented *An. gambiae*, *An. ziemanni*, *Culex* sp, *An. rufipes*, *Mansonia* sp, *Aedes* sp, *An. squamosus*, *An. pharonsis* in Oussouye in 2012.

Molecular detection of filariae

Patients

A total of 37 blood samples were collected (23 daytime and 14 night-time) from the 23 elephantiasis patients and analyzed by qPCR. The pan-filarial 28S qPCR system yielded two positive results in different patient samples, a positive elephantiasis patient (day and night) in Kolda with both legs swollen and another patient who has hydrocele is positive only at night in Oussouye, representing an infection prevalence of 9% (2/23). These two samples were also positive for qPCR targeting the *M. perstans*-specific ITS gene (**Table 2**).

Of the 279 samples taken from febrile Mlomp patients, 16 were positive by pan-filarial 28S qPCR, a prevalence of 6%. Among these samples, 12 were positive to qPCR targeting ITS sequences specific to *M. perstans* and standard PCR targeting mitochondrial cox1 gene giving a prevalence of *M. perstans* of 4.3% (12/279). (**Table 2**). Other four samples (all with elevated ct) did not produce positive results neither with *M. perstans* qPCR nor with pan-filarial standard PCR targeting cox1 and were considered as negative.

Of 3,714 RDTs analyzed, 13 were positive for the 28S pan-filarial qPCR system, for an overall prevalence of 0.35%. Among them, 11 were also positive for qPCR targeting *M. perstans*, for a prevalence of 0.3% (11/3,714). Prevalence by regions were as follows: 1% (6/762) in Kedougou, 0.1% (1/762) in Kolda, 0.1% (1/863) in Oussouye and 0.2% (3/1327) in Sedhiou (**Table 3**).

Probable mitochondrial cytochrome oxidase copy of *M. perstans* in nuclear genome

Amplification targeting the cox 1 gene using primers COIntF (5'-TGATTGGTGGTTTTGGTAA-3') and COIntR (5'-ATAAGTACGAGTATCAATATC-3') was performed on *M. perstans* positive samples. Most samples produced double bands visualized during gel migration.

All bands obtained were re-sequenced individually. The heavy bands appear to be the *M. perstans* species and for the light bands, BLAST showed the similarity of 82-83% with different filarial species, including *Loa loa*, *Brugia timori*, *Brugia malayi* and *W. bancrofti* (**Fig. 3**). Based on the obtained sequence we have developed the specific qPCR; all samples positive for *M. perstans* were also positive for this second “neofilarial” cox1 filarial variant. Moreover, analyzing the 617 bps cox1 fragment alignments we have detected at least 4 single nucleotide deletions causing multiple stop codons.

These results suggested that the detected “neofilarial” DNA represent NUMT [39] gene, a non-functional copy of the mitochondrial genome present in the nuclear genome.




Fig. 3 Phylogenetic tree based on the functional mitochondrial and NUMT copies of *cox1* sequences of *Mansonella perstans* identified in blood samples from febrile subjects and other filarial species. The sequences were aligned using ClustalW. The tree was deduced using the maximum likelihood method based on 100 bootstraps. The tree is drawn to scale, with branch lengths in the same units as the evolutionary distances used to infer the phylogenetic tree. Evolutionary distances were calculated using the Tamura-Nei method and are expressed in units of the number of base substitutions per site. Evolutionary analyses were performed using MEGA7.

Arthropods

Among 254 mosquitoes collected in 2021 in the bedrooms of elephantiasis patients we detected 5 (5/254, 2%) individuals positive with 28S pan-filarial qPCR system. All 5 specimens were identified as *Mansonia uniformis*, that resulted in the infection rate of *M. uniformis* at 3.5% (5/142).

5 specimens of *Aedes* sp. among 444 (1.1%) in Kedougou collected in 2007 and 8 mosquitoes including 5 *An. ziemanni*, 2 *Culex* sp. and 1 *An. gambiae* among 789 (1%) collected in Oussouye in 2012 were also positive in pan-filaria qPCR.

None of 991 *Culicoides* spp. collected in 2021 in patient's bedrooms was detected positive for *M. perstans* by specific qPCR.

Sequencing confirmation of filariae

Patients

The presence of *M. perstans* in the 2 elephantiasis patients were confirmed by sequencing the following genes ITS1 (443pb; PQ041263-PQ041267), COX1 (643pb; PQ097864-PQ097865) and 5S (379pb; PQ043272-PQ043273) that were identical among each other and shared 100% identity with different *M. perstans* samples in Genbank.

In febrile patients from Mlomp health dispensary, sequencing also confirmed the presence of *M. perstans* with 100% identity targeting the ITS1 (443pb; PQ041264) and 5S (379pb; PQ043274) genes. With regard to the RDTs, sequencing of the ITS1 gene confirmed the presence of *M. perstans* (PQ041650) in 9 samples with 100% identity (**Fig. 4**).





Fig. 4. Phylogenetic tree based on ITS1 sequence showing the clustering of the *Mansonella* identified in malaria-negative RDTs with other filarial species. Sequence alignment was performed using ClustalW. The tree was deduced using the maximum likelihood method based on 100 bootstraps. The tree is drawn to scale, with branch lengths in the same units as the evolutionary distances used to infer the phylogenetic tree. Evolutionary distances were calculated using the Tamura-Nei method and are expressed in units of the number of base substitutions per site. Evolutionary analyses were carried out using MEGA7.

Arthropods

We have amplified and sequenced a portion of filarial mitochondrial 12S from all mosquito samples positive by pan-filarial qPCR. Sanger sequencing of these fragments showed that different mosquito species harbored at least 5 different filarial species (Figure 5). *Setaria digitata* (PQ047250) and *Setaria labiাপapillosa* (PQ047258) were identified in *Mansonia africana*, and *An. coustani*; *Onchocerca* sp. (PQ047260), *Onchocercidea* sp. (PQ047256) and

375 Filarioidea sp. (PQ047257) in *Culex cf. sitiens* and *Culex* sp. mosquitoes collected in 2021-
 376 2007 and 2012 (**Fig. 5**). All sequences obtained have been submitted to the GenBank database.



377 **Fig. 5** Phylogenetic tree based on the 12S gene sequence of 31 specimens showing filarial
 378 species found in arthropods collected between 2021 - 2007 and in 2012. Sequence alignment
 379 was performed using ClustalW. The evolutionary history was inferred using the maximum
 380 likelihood method based on the Tamura-Nei model. Nodes are associated with bootstrap values
 381 based on 100 replicates. Bootstrap support values <70 is not shown. Initial trees for heuristic
 382 search were obtained automatically by applying the Neighbor-Join and BioNJ algorithms to a
 383 matrix of pairwise distances estimated using the Composite Maximum Likelihood (CML)
 384 approach, then selecting the topology with the highest log-likelihood value. The tree is drawn
 385 to scale, with branch lengths measured as the number of substitutions per site. Evolutionary
 386 analyses were performed with MEGA7.

387

Discussion

Our study enabled us to detect *M. perstans* in patients suffering from elephantiasis and hydrocele in the Kolda and Oussouye zones in Senegal. Neither *W. bancrofti*, nor any other filarial parasite responsible for lymphatic filariasis were detected in patients with elephantiasis. Patients were sampled either once or twice (day and/or night, taking into account the circadian rhythms of certain filaria). In the majority of patients (21/23), no filarial parasites were identified in the blood.

It is known that in endemic areas the prevalence of lymphatic filaria in blood of the total population and even in elephantiasis patients may vary significantly. Often it is about 0.2-2% [40] but sometimes may be as high as 18.5% [41]. PCR is a very sensitive and specific test to detect filaria in blood samples; usually all cases positive in microscopy and by antigen tests are also positive in specific qPCR [42].

A possible explanation may be that the examined patients may be already treated by anti-filarial drugs. Indeed, the implementation of the 2016-2020 and 2022-2025 Strategic Plans to combat neglected tropical diseases (NTDs) in Senegal (NTD, 2016-2020) (NTD, 2022-2025), targeted lymphatic filariasis among other diseases [43,44].

However, in our cohort of patient, only two were treated by antifilarial drugs, others denied any treatment in the pass. Two patients positive for *M. perstans* also refused to receive any anti-filarial treatment. *M. perstans* infection is, however, known to be tenacious due to its insensitivity to filaricidal drugs [21].

Our results of the screening of febrile patients for filarial parasites were concordant. Neither in DNA extracted directly from patients, nor in DNA extracted from nitrocellulose strips from malaria RDT *W. bancrofti* DNA was not identified; the only filarial species identified was *M. perstans*. In febrile subjects from Mlomp *M. perstans* was detected in 4,3% of samples. A hydrocele patient with *M. perstans* microfilariae lived in several kilometers from the Mlomp dispensary.

Malaria-negative RDT cassettes were used to determine the prevalence of *M. perstans* in febrile patients with malaria symptoms in the different study areas. Overall prevalence was relatively low. In Kedougou, endemic region for *M. perstans* [20], we found 1% of RDT samples positive. The total prevalence of confirmed cases in RDT was 0.24%. However, these results may be impacted by several factors as small volume of the blood samples conserved in

nitrocellulose strip and unstable conditions and long terms of storage of RDT cassettes that may impact the DNA conservation.

Taking together, none of three sampling series (elephantiasis patients, febrile patients with DNA extracted in Mlomp dispensary laboratory and RDT collected from other dispensaries) detected filaria other than *M. perstans*.

W. bancrofti lymphatic filariasis in Senegal has been described only morphologically, without any genetic confirmation. Our data provide substrate for further research on the presence of Bancroftian filariasis in Senegal and probable role of *M. perstans* in elephantiasis. In Uganda, a similar study carried out between 2000 and 2003 on elephantiasis patients reported a low proportion (0.3%) of *W. bancrofti* and *M. perstans* co-infection in an area where the two parasites coexisted [11]. Both elephantiasis patients with confirmed *M. perstans* in our study were older men between 76 and 79 years of age living in rural areas. These findings confirm previous studies [45–47].

Characterizing *M. perstans* positive samples we occasionally amplified and sequenced possible NUMT of *cox1* gene. All samples positive for *M. perstans* were positive for this unusual “neofilarial” *cox1* gene by specific qPCR, and the gene itself was not functional containing many deletions and stop-codons. The NUMTs in *M. ozzardi* were already reported [48] and in *M. perstans* were recently identified during sequencing of a whole genome of *M. perstans* from Cameroon [49]. Interesting that in case of *Mansonella*, the NUMTs are poorly conserved and present a weak identity with functional mitochondrial genes in 86-87% [49].

We tried to identify the vectors of filarial parasites in Senegal. None of those tested were positive for *W. bancrofti* or *M. perstans*. Previously, the vector had not been detected in *Culicoides* biting midges in rural Senegal (Kedougou region). In our study we also tested biting midges collected in endemic regions for *M. perstans*; we tested also mosquitoes trying to identify another vector. We have identified three undescribed filarial genotypes that have never been identified in humans. One is obviously grouped within *Onchocerca* genus two others are identified as *Filarioidea* sp and *Onchocercidea* sp. Our genotypes are close to those previously identified in domestic cattle of the "Ndama" breed in the Guinean savannah regions of Kedougou, Senegal [50]. Other filaria identified in mosquitoes, *S. digitata* and *S. labiopallosa* are mosquito-borne filaria found in the abdominal cavity of cattle [51]. *S. digitata* and *S. labiopallosa* have been detected in *An. coustani* and *Mansonia africana*, an exophilic and exophagous mosquito [52]. This species prefers to bite cattle, sheep and poultry. Females are

very aggressive and can bite during the day [53]. *M. africana*, a mosquito found in swampy areas that bites outdoors at night, although some are endophilic and behave exophilically after a blood meal. Like other zoonotic vectors, *Culex sitiens* [54] is an important coastal vector of many mosquito-borne diseases [55], including Japanese encephalitis [56], found in mangroves, salt marshes and crab holes. It is also a species that bites humans at all hours, but especially at dusk and at night.

Conclusion

Overall, for the first time in Senegal, *M. perstans* was found in elephantiasis patients in the absence of “classical” lymphatic filaria. Screening of blood samples (based on DNA extracted from malaria RDT and from febrile patients in dispensary) documented wide distribution of *M. perstans* and virtual absence of other human pathogenic filaria. Our data suggest that *M. perstans* mansonellosis, although asymptomatic, remains endemic in the southern regions of Senegal. A cost-effective approach to continue studies of filariases may be to reuse *P. falciparum* RDTs as a diagnostic tool for monitoring prevalence, parasite density and other blood-borne diseases. The role of *M. perstans* a causative agent of elephantiasis should be further examined.

Références

- Adam, Nouhou Diori, Yacoubou Soumana, Ali Saley, et Idrissa Saley. 2020. « Une Loase Oculaire: À Propos D'un Cas ». *Pan African Medical Journal* 36. <https://doi.org/10.11604/pamj.2020.36.302.25124>.
- Adewole, S.O., et S.K. Ayeni. 2009. « Clinical Manifestation of Onchocerciasis in Ise - Orun Local Government, Ekiti State, Nigeria ». *Pakistan Journal of Nutrition* 8 (2): 122-24. <https://doi.org/10.3923/pjn.2009.122.124>.
- Adhikari, Ram Kumar, Jeevan Bahadur Sherchand, Shiva Raj Mishra, Kamal Ranabhat, Amrit Pokharel, Pramila Devkota, Durga Mishra, et al. 2015. « Health-Seeking Behaviors and Self-Care Practices of People with Filarial Lymphoedema in Nepal: A Qualitative Study ». *Journal of Tropical Medicine* 2015:1-6. <https://doi.org/10.1155/2015/260359>.
- Adolph, Paul E., Irving G. Kagan, et Russell M. McQuay. 1962. « Diagnosis and Treatment of Acanthocheilonema Perstans Filariasis », janvier. <https://doi.org/10.4269/ajtmh.1962.11.76>.
- Agbolade, Olufemi Moses, Dora Olufunmilola Akinboye, Taiwo Monroof Olateju, Oluwatoyin Adepeju Ayanbiyi, Olukayode Olakunle Kuloyo, et Oluwalolami Oluwaseun Fenuga. 2006. « Biting of Anthropophilic Culicoides Fulvithorax (Diptera: Ceratopogonidae), a Vector of Mansonella Perstans in Nigeria ». *The Korean Journal of Parasitology* 44 (1): 67-72. <https://doi.org/10.3347/kjp.2006.44.1.67>.
- Akue, Jean Paul, Dieudonné Nkoghe, Cindy Padilla, Ghislain Moussavou, Hubert Moukana, Roger Antoine Mbou, Benjamin Ollomo, et Eric Maurice Leroy. 2011. « Epidemiology of Concomitant Infection Due to Loa Loa and Mansonella Perstans in Gabon ». Édité par Rachel Lisa Pullan. *PLoS Neglected Tropical Diseases* 5 (10): e1329. <https://doi.org/10.1371/journal.pntd.0001329>.
- Alhassan, Andy, Zhiru Li, Catherine B. Poole, et Clotilde K.S. Carlow. 2015. « Expanding the MDx Toolbox for Filarial Diagnosis and Surveillance ». *Trends in Parasitology* 31 (8): 391-400. <https://doi.org/10.1016/j.pt.2015.04.006>.
- Allen, Judith E., Ohene Adjei, Odile Bain, Achim Hoerauf, Wolfgang H. Hoffmann, Benjamin L. Makepeace, Hartwig Schulz-Key, et al. 2008. « Of Mice, Cattle, and Humans: The Immunology and Treatment of River Blindness ». Édité par Sara Lustigman. *PLoS Neglected Tropical Diseases* 2 (4): e217. <https://doi.org/10.1371/journal.pntd.0000217>.
- Alsarraf, Mustafa, Viktoria Levytska, Ewa J. Mierzejewska, Vasyl Poliukhovych, Anna Rodo, Mohammed Alsarraf, Dzyiyana Kavalevich, Dorota Dwużnik-Szarek, Jerzy M. Behnke,

- et Anna Bajer. 2021. « Emerging Risk of *Dirofilaria* Spp. Infection in Northeastern Europe: High Prevalence of *Dirofilaria Repens* in Sled Dog Kennels from the Baltic Countries ». *Scientific Reports* 11 (1): 1068. <https://doi.org/10.1038/s41598-020-80208-1>.
- Amazigo, U. 2008. « The African Programme for Onchocerciasis Control (APOC) ». *Annals of Tropical Medicine & Parasitology* 102 (sup1): 19-22. <https://doi.org/10.1179/136485908X337436>.
- Anderson, Roy C. 2000. « R.C. Anderson: Nematode Parasites of Vertebrates. Their Development and Transmission. 2nd Edition. CABI Publishing, Wallingford, Oxon (UK) 2000. 650 Pp., 43 Figs. Price £ 99.50 . », 1.
- Anosike, Jude C, Ikechukwu N S Dozie, Celestine O E Onwuliri, Bertram E B Nwoke, et Viola A Onwuliri. 2005. « PREVALENCE OF MANSONELLA PERSTANS INFECTIONS AMONG THE NOMADIC FULANIS OF NORTHERN NIGERIA », p.35-39. <https://pubmed.ncbi.nlm.nih.gov/16028864/>.
- Antinori, Spinello, Carlo Parravicini, Laura Galimberti, Antonella Tosoni, Paolo Giunta, Massimo Galli, Mario Corbellino, et Anna Lisa Ridolfo. 2017. « Is Imported Onchocerciasis a Truly Rare Entity? Case Report and Review of the Literature ». *Travel Medicine and Infectious Disease* 16 (mars):11-17. <https://doi.org/10.1016/j.tmaid.2017.02.006>.
- Asio, S. M., P. E. Simonsen, et A. W. Onapa. 2009. « *Mansonella Perstans* : Safety and Efficacy of Ivermectin Alone, Albendazole Alone and the Two Drugs in Combination ». *Annals of Tropical Medicine & Parasitology* 103 (1): 31-37. <https://doi.org/10.1179/136485909X384929>.
- Asio, Santa Maria, Paul E. Simonsen, et Ambrose W. Onapa. 2009a. « Analysis of the 24-h Microfilarial Periodicity of *Mansonella Perstans* ». *Parasitology Research* 104 (4): 945-48. <https://doi.org/10.1007/s00436-008-1312-x>.
- . 2009b. « *Mansonella Perstans* Filariasis in Uganda: Patterns of Microfilaraemia and Clinical Manifestations in Two Endemic Communities ». *Transactions of the Royal Society of Tropical Medicine and Hygiene* 103 (3): 266-73. <https://doi.org/10.1016/j.trstmh.2008.08.007>.
- Attout, Tarik, Achim Hoerauf, Gaëlle Dénécé, Alexander Yaw Debrah, Yeboah Marfo-Debrekyei, Michel Boussinesq, Samuel Wanji, et al. 2009. « Lymphatic Vascularisation and Involvement of Lyve-1+ Macrophages in the Human *Onchocerca* Nodule ». *Édité*

545 par Derya Unutmaz. *PLoS ONE* 4 (12): e8234.
 546 <https://doi.org/10.1371/journal.pone.0008234>.

547 Aubry, Arthur, Charles Hervochon, Marie Chauveau, Marie Denis-Musquer, Fakhri Jeddi,
 548 Rose-Anne Lavergne, Patrice Le Pape, et Florent Morio. 2022. « Une expulsion bien
 549 singulière d'un ver adulte de *Loa loa* ». *Revue Francophone des Laboratoires* 2022
 550 (538): 72-75. [https://doi.org/10.1016/S1773-035X\(21\)00381-6](https://doi.org/10.1016/S1773-035X(21)00381-6).

551 Aubry, Professeur Pierre, et Docteur Bernard-Alex Gaüzère. 2023. « Filarioses lymphatiques
 552 Actualités 2023 », P8.

553 Babu, Subash, et Thomas B. Nutman. 2012. « Immunopathogenesis of Lymphatic Filarial
 554 Disease ». *Seminars in Immunopathology* 34 (6): 847-61.
 555 <https://doi.org/10.1007/s00281-012-0346-4>.

556 Badia-Rius, Xavier, Salissou Adamou, Mark J. Taylor, et Louise A. Kelly-Hope. 2023.
 557 « Morbidity Hotspot Surveillance: A Novel Approach to Detect Lymphatic Filariasis
 558 Transmission in Non-Endemic Areas of the Tillabéry Region of Niger ». *Parasite
 559 Epidemiology and Control* 21 (mai):e00300.
 560 <https://doi.org/10.1016/j.parepi.2023.e00300>.

561 Bassene, Hubert, Masse Sambou, Florence Fenollar, Siân Clarke, Sawdiatou Djiba, Gaël
 562 Mourembou, Alioune Badara L Y, Didier Raoult, et Oleg Mediannikov. 2015a. « High
 563 Prevalence of *Mansonella Perstans* Filariasis in Rural Senegal ». *The American Journal
 564 of Tropical Medicine and Hygiene* 93 (3): 601-6. <https://doi.org/10.4269/ajtmh.15-0051>.
 565 0051.

566 ———. 2015b. « High Prevalence of *Mansonella Perstans* Filariasis in Rural Senegal ». *The
 567 American Journal of Tropical Medicine and Hygiene* 93 (3): 601-6.
 568 <https://doi.org/10.4269/ajtmh.15-0051>.

569 Bhalla, Devender, Michel Dumas, et Pierre-Marie Preux. 2013. « Neurological Manifestations
 570 of Filarial Infections ». In *Handbook of Clinical Neurology*, 114:235-42. Elsevier.
 571 <https://doi.org/10.1016/B978-0-444-53490-3.00018-2>.

572 Boatin, B. 2008. « The Onchocerciasis Control Programme in West Africa (OCP) ». *Annals of
 573 Tropical Medicine & Parasitology* 102 (sup1): 13-17.
 574 <https://doi.org/10.1179/136485908X337427>.

575 Bockarie, Moses J, et Rinki M Deb. 2010. « Elimination of Lymphatic Filariasis: Do We Have
 576 the Drugs to Complete the Job? ». *Current Opinion in Infectious Diseases* 23 (6):
 577 617-20. <https://doi.org/10.1097/QCO.0b013e32833fdee5>.

578 Bockarie, Moses J, Mark J Taylor, et John O Gyapong. 2009. « Current Practices in the
579 Management of Lymphatic Filariasis ». *Expert Review of Anti-Infective Therapy* 7 (5):
580 595-605. <https://doi.org/10.1586/eri.09.36>.

581 Bottieau, Emmanuel, Ralph Huits, Steven Van Den Broucke, Ula Maniewski, Steven Declercq,
582 Isabel Brosius, Caroline Theunissen, et al. 2022. « Human Filariasis in Travelers and
583 Migrants: A Retrospective 25-Year Analysis at the Institute of Tropical Medicine,
584 Antwerp, Belgium ». *Clinical Infectious Diseases* 74 (11): 1972-78.
585 <https://doi.org/10.1093/cid/ciab751>.

586 Bouchery, T., E. Lefoulon, G. Karadjian, A. Nieguitsila, et C. Martin. 2013. « The Symbiotic
587 Role of Wolbachia in Onchocercidae and Its Impact on Filariasis ». *Clinical
588 Microbiology and Infection* 19 (2): 131-40. <https://doi.org/10.1111/1469-0691.12069>.

589 Bourée, P. 2005. « Parasitoses urinaires ». *Annales d'Urologie* 39 (6): 232-46.
590 <https://doi.org/10.1016/j.anuro.2005.09.005>.

591 Boussinesq, M. 2003. « L'onchocercose humaine en Afrique ». In *Trente ans de lutte contre
592 l'onchocercose en Afrique de l'Ouest. Traitements larvicides et protection de
593 l'environnement : Thirty years of onchocerciasis control in West Africa. Blackfly
594 larviciding and environmental protection*, édité par Laurent Yaméogo, Christian
595 Levêque, et Jean-Marc Hougard, 389-400. Synthèses. Marseille: IRD Éditions.
596 <https://doi.org/10.4000/books.irdeditions.28767>.

597 ———. 2006. « Loiasis ». *Annals of Tropical Medicine & Parasitology* 100 (8): 715-31.
598 <https://doi.org/10.1179/136485906X112194>.

599 Boussinesq, Michel. 2012. « Loiasis: New Epidemiologic Insights and Proposed Treatment
600 Strategy ». *Journal of Travel Medicine* 19 (3): 140-43. [https://doi.org/10.1111/j.1708-
601 8305.2012.00605.x](https://doi.org/10.1111/j.1708-8305.2012.00605.x).

602 Brattig, Norbert W., Robert A. Cheke, et Rolf Garms. 2021. « Onchocerciasis (River Blindness)
603 – More than a Century of Research and Control ». *Acta Tropica* 218 (juin):105677.
604 <https://doi.org/10.1016/j.actatropica.2020.105677>.

605 Bregani, E. R., F. Tantardini, et A. Rovellini. 2007. « [Mansonella perstans filariasis] ». *Parassitologia* 49 (1-2): 23-26.

607 Bregani, Enrico Rino, Angelo Rovellini, Narasem Mbaïdoum, et Maria Giovanna Magnini.
608 2006. « Comparison of Different Anthelmintic Drug Regimens against Mansonella
609 Perstans Filariasis ». *Transactions of the Royal Society of Tropical Medicine and
610 Hygiene* 100 (5): 458-63. <https://doi.org/10.1016/j.trstmh.2005.07.009>.

- Brengues, J., R. Subra, J. Mouchet, et G. S. Nelson. 1968. « [Transmission of *Wuchereria bancrofti* Cobbold in West Africa. Preliminary study of a focus in the savanna of north Guinea] ». *Bulletin of the World Health Organization* 38 (4): 595-608.
- Brugman, Victor A., Marion E. England, Joanne Stoner, Laura Tugwell, Lara E. Harrup, Anthony J. Wilson, Jolyon M. Medlock, et al. 2017. « How Often Do Mosquitoes Bite Humans in Southern England? A Standardised Summer Trial at Four Sites Reveals Spatial, Temporal and Site-Related Variation in Biting Rates ». *Parasites & Vectors* 10 (1): 420. <https://doi.org/10.1186/s13071-017-2360-9>.
- Burbelo, Peter D., Roshan Ramanathan, Amy D. Klion, Michael J. Iadarola, et Thomas B. Nutman. 2008. « Rapid, Novel, Specific, High-Throughput Assay for Diagnosis of *Loa Loa* Infection ». *Journal of Clinical Microbiology* 46 (7): 2298-2304. <https://doi.org/10.1128/JCM.00490-08>.
- Büttner, Dietrich W., Samuel Wanji, Chiara Bazzocchi, Odile Bain, et Peter Fischer. 2003. « Obligatory Symbiotic *Wolbachia* Endobacteria Are Absent from *Loa Loa* ». *Filaria Journal* 2 (1): 10. <https://doi.org/10.1186/1475-2883-2-10>.
- CAMPILLO, Jérémy T., Frédéric LOUYA, Paul BIKITA, François MISSAMOU, Michel BOUSSINESQ, Sebastien D. S. PION, Sébastien BERTOUT, et Cédric B CHESNAIS. 2023. « Évaluation en milieu rural au Congo de la fiabilité de la méthode de la goutte épaisse calibrée pour évaluer le niveau de la microfilarémie à *Loa loa* ». *MTSI*, mars, Vol. 3 No 1 (2023): MTSIRevue. <https://doi.org/10.48327/MTSI.V3I1.2023.297>.
- Carne, B., et P. Esterre. 2012. « Filarioses ». *EMC - Maladies infectieuses* 9 (2): 1-19. [https://doi.org/10.1016/S1166-8598\(12\)41786-0](https://doi.org/10.1016/S1166-8598(12)41786-0).
- CDC. 2024. « Symptoms of Onchocerciasis ». *Filarial Worms*. 5 juin 2024. <https://www.cdc.gov/filarial-worms/signs-symptoms/onchocerciasis.html>.
- Chakraborty, Sanjukta, Manokaran Gurusamy, David C. Zawieja, et Mariappan Muthuchamy. 2013. « Lymphatic Filariasis: Perspectives on Lymphatic Remodeling and Contractile Dysfunction in Filarial Disease Pathogenesis ». *Microcirculation* 20 (5): 349-64. <https://doi.org/10.1111/micc.12031>.
- Chandy, Anish, Alok Singh Thakur, Mukesh Pratap Singh, et Ashish Manigauha. 2011. « A Review of Neglected Tropical Diseases: Filariasis ». *Asian Pacific Journal of Tropical Medicine* 4 (7): 581-86. [https://doi.org/10.1016/S1995-7645\(11\)60150-8](https://doi.org/10.1016/S1995-7645(11)60150-8).
- Chelsea, Marie, et A. Petri William. 2022. « Mansonellose-Maladies infectieuses ». <https://www.msmanuals.com/fr/professional/maladies-infectieuses/n%C3%A9matodes-vers-ronds/mansonellose>.

- Chenna, Ramu, Hideaki Sugawara, Tadashi Koike, Rodrigo Lopez, Toby J. Gibson, Desmond G. Higgins, et Julie D. Thompson. 2003. « Multiple Sequence Alignment with the Clustal Series of Programs ». *Nucleic Acids Research* 31 (13): 3497-3500. <https://doi.org/10.1093/nar/gkg500>.
- Chesnais, Cédric B, Charlotte Bizet, Jérémy T Campillo, Wepnyu Y Njamnshi, Jean Bopda, Philippe Nwane, Sébastien D Pion, Alfred K Njamnshi, et Michel Boussinesq. 2020. « A Second Population-Based Cohort Study in Cameroon Confirms the Temporal Relationship Between Onchocerciasis and Epilepsy ». *Open Forum Infectious Diseases* 7 (6): ofaa206. <https://doi.org/10.1093/ofid/ofaa206>.
- Coffeng, Luc E., Wilma A. Stolk, Achim Hoerauf, Dik Habbema, Roel Bakker, Adrian D. Hopkins, et Sake J. De Vlas. 2014. « Elimination of African Onchocerciasis: Modeling the Impact of Increasing the Frequency of Ivermectin Mass Treatment ». Édité par James M. McCaw. *PLoS ONE* 9 (12): e115886. <https://doi.org/10.1371/journal.pone.0115886>.
- Colebunders, Robert, Jane Y Carter, Peter Claver Olore, Kai Puok, Samit Bhattacharyya, Sonia Menon, Gasim Abd-Elfarag, et al. 2018. « High Prevalence of Onchocerciasis-Associated Epilepsy in Villages in Maridi County, Republic of South Sudan: A Community-Based Survey ». *Seizure* 63 (décembre):93-101. <https://doi.org/10.1016/j.seizure.2018.11.004>.
- Coreil, Jeannine, et Gladys Mayard. 2009. « Indigenization of Illness Support Groups for Lymphatic Filariasis in Haiti ». In *Anthropology and Public Health*, édité par Robert A Hahn et Marcia Inborn, 2^e éd., 245-65. Oxford University Press New York. <https://doi.org/10.1093/acprof:oso/9780195374643.003.0010>.
- Coulibaly, Yaya I., Benoit Dembele, Abdallah A. Diallo, Ettie M. Lipner, Salif S. Doumbia, Siaka Y. Coulibaly, Siaka Konate, et al. 2009a. « A Randomized Trial of Doxycycline for *Mansonella Perstans* Infection ». *New England Journal of Medicine* 361 (15): 1448-58. <https://doi.org/10.1056/NEJMoa0900863>.
- . 2009b. « A Randomized Trial of Doxycycline for *Mansonella Perstans* Infection ». *The New England Journal of Medicine* 361 (15): 1448-58. <https://doi.org/10.1056/NEJMoa0900863>.
- Coulibaly, Yaya Ibrahim, Benoit Dembele, Abdallah Amadou Diallo, Sibylle Kristensen, Siaka Konate, Housseini Dolo, Ilo Dicko, et al. 2013. « Wuchereria Bancrofti Transmission Pattern in Southern Mali Prior to and Following the Institution of Mass Drug

- Administration ». *Parasites & Vectors* 6 (1): 247. <https://doi.org/10.1186/1756-3305-6-247>.
- Crawford, Katie E, Shannon M Hedtke, Stephen R Doyle, Annette C Kuesel, Samuel Armoo, Mike Y Osei-Atweneboana, et Warwick N Grant. 2024. « Genome-Based Tools for Onchocerciasis Elimination: Utility of the Mitochondrial Genome for Delineating Onchocerca Volvulus Transmission Zones ». *International Journal for Parasitology* 54 (3-4): 171-83. <https://doi.org/10.1016/j.ijpara.2023.11.002>.
- Cromwell, Elizabeth A, Chris A Schmidt, Kevin T Kwong, David M Pigott, Denise Mupfasoni, Gautam Biswas, Shreya Shirude, et al. 2020. « The Global Distribution of Lymphatic Filariasis, 2000–18: A Geospatial Analysis ». *The Lancet Global Health* 8 (9): e1186-94. [https://doi.org/10.1016/S2214-109X\(20\)30286-2](https://doi.org/10.1016/S2214-109X(20)30286-2).
- Davis, Emma L., Lisa J. Reimer, Lorenzo Pellis, et T. Deirdre Hollingsworth. 2019. « Evaluating the Evidence for Lymphatic Filariasis Elimination ». *Trends in Parasitology* 35 (11): 860-69. <https://doi.org/10.1016/j.pt.2019.08.003>.
- De Sole, G., J. Giese, F.M. Keita, et J. Remme. 1991. « Detailed Epidemiological Mapping of Three Onchocerciasis Foci in West Africa ». *Acta Tropica* 48 (3): 203-13. [https://doi.org/10.1016/0001-706X\(91\)90048-O](https://doi.org/10.1016/0001-706X(91)90048-O).
- Debrah, Alexander Yaw, Sabine Specht, Ute Klarmann-Schulz, Linda Batsa, Sabine Mand, Yeboah Marfo-Debrekyei, Rolf Fimmers, et al. 2015. « Doxycycline Leads to Sterility and Enhanced Killing of Female *Onchocerca Volvulus* Worms in an Area With Persistent Microfilaridermia After Repeated Ivermectin Treatment: A Randomized, Placebo-Controlled, Double-Blind Trial ». *Clinical Infectious Diseases* 61 (4): 517-26. <https://doi.org/10.1093/cid/civ363>.
- Debrah, Linda Batsa, Norman Nausch, Vera Serwaa Opoku, Wellington Owusu, Yusif Mubarik, Daniel Antwi Berko, Samuel Wanji, et al. 2017. « Epidemiology of Mansonella perstans in the middle belt of Ghana ». *Parasites & Vectors* 10 (janvier):15. <https://doi.org/10.1186/s13071-016-1960-0>.
- Demanou, M., S. D. Pion, et M. Boussinesq. 2001. « [Entomologic study of loaiasis transmission in the Lekie area (Cameroon)] ». *Bulletin De La Societe De Pathologie Exotique (1990)* 94 (4): 347-52.
- Dembélé, Pascal, Mady Cissoko, Adama Zan Diarra, Lassana Doumbia, Aïssata Koné, Mahamadou H. Magassa, Maissane Mehadji, Mahamadou A. Thera, et Stéphane Ranque. 2024. « Evaluation of the Performance of Rapid Diagnostic Tests for Malaria Diagnosis and Mapping of Different Plasmodium Species in Mali ». *International*

- Journal of Environmental Research and Public Health* 21 (2): 228.
<https://doi.org/10.3390/ijerph21020228>.
- Deng, Miao-Han, Lan-Yi Zhong, Okanurak Kamolnetr, Yanin Limpanont, et Zhi-Yue Lv. 2019. « Detection of Helminths by Loop-Mediated Isothermal Amplification Assay: A Review of Updated Technology and Future Outlook ». *Infectious Diseases of Poverty* 8 (1): 20. <https://doi.org/10.1186/s40249-019-0530-z>.
- Desjardins, Christopher A, Gustavo C Cerqueira, Jonathan M Goldberg, Julie C Dunning Hotopp, Brian J Haas, Jeremy Zucker, José M C Ribeiro, et al. 2013. « Genomics of Loa Loa, a Wolbachia-Free Filarial Parasite of Humans ». *Nature Genetics* 45 (5): 495-500. <https://doi.org/10.1038/ng.2585>.
- Desoubeaux, Guillaume, Adélaïde Chesnay, et Jacques Chandenier. 2020. « Approches diagnostiques des filarioses en France métropolitaine ». *Revue Francophone des Laboratoires* 2020 (519): 72-80. [https://doi.org/10.1016/S1773-035X\(20\)30066-6](https://doi.org/10.1016/S1773-035X(20)30066-6).
- Devi, R Renuka, et V Raju. 2018. « Information Technology in Morbidity Management of Human Lymphatic Filariasis—A Promising Tool in Global Programme for Elimination of Lymphatic Filariasis ». *Journal of Vector Borne Diseases* 55 (1): 20. <https://doi.org/10.4103/0972-9062.234622>.
- Diagne, Jean Pierre, Aly Mbara Ka, Joseph Matar Ndiaye, ABoubacry Sadikh Sow, Aissatou Maguatte Wane, et Paule Aida Ndoeye Roth. 2018. « MIGRATION SOUS CUTANEE PALPEBRALE D'UNE LOASE: A PROPOS D'UN CAS », n° 1, 4. https://www.revues-ufhb-ci.org/fichiers/FICHIR_ARTICLE_2511.pdf.
- Diallo, S., M. Sarr, S. Diagne, et L. Konate. 1977. « Filariasis in the Nianing région" Senegal. Prevalence, clinical aspects, entomological findings. » *Medecine d'Afrique Noire* 24 (4): 233-42. <https://www.cabdirect.org/cabdirect/abstract/19772902103>.
- Díaz-Menéndez, Marta, Francesca Norman, Begoña Monge-Maillo, José Antonio Pérez-Molina, et Rogelio López-Vélez. 2011. « Las filariasis en la práctica clínica ». *Enfermedades Infecciosas y Microbiología Clínica* 29 (décembre):27-37. [https://doi.org/10.1016/S0213-005X\(11\)70041-6](https://doi.org/10.1016/S0213-005X(11)70041-6).
- Diwakar, Deepak Kumar, Nikita Wadhwani, Nikhil Prasad, et Ashwani Gupta. 2018. « Primary Retroperitoneal Filariasis: A Common Disease of Tropics with Uncommon Presentation and Review of Literature ». *BMJ Case Reports*, septembre, bcr-2018-226217. <https://doi.org/10.1136/bcr-2018-226217>.
- Dolo, Housseini, Yaya Ibrahim Coulibaly, Louise Kelly-Hope, Siaka Konate, Benoit Dembele, Siaka Yamoussa Coulibaly, Dramane Sanogo, et al. 2018. « Factors Associated with

- Wuchereria Bancrofti Microfilaremia in an Endemic Area of Mali ». *The American Journal of Tropical Medicine and Hygiene* 98 (6): 1782-87.
<https://doi.org/10.4269/ajtmh.17-0902>.
- Downes, B, et K Jacobsen. 2010. « A systematic review of the epidemiology of mansonelliasis ». *African Journal of Infectious Diseases* 4 (1).
<https://doi.org/10.4314/ajid.v4i1.55085>.
- Downes, Barbara L., et Kathryn H. Jacobsen. 2010. « A Systematic Review of the Epidemiology of Mansonelliasis ». *African Journal of Infectious Diseases* 4 (1): 7-14.
<https://doi.org/10.4314/ajid.v4i1.55085>.
- Drame, Papa M., Céline Montavon, Sébastien D. Pion, Joseph Kubofcik, Michael P. Fay, et Thomas B. Nutman. 2016. « Molecular Epidemiology of Blood-Borne Human Parasites in a Loa Loa-, Mansonella Perstans-, and Plasmodium Falciparum-Endemic Region of Cameroon ». *The American Journal of Tropical Medicine and Hygiene* 94 (6): 1301-8.
<https://doi.org/10.4269/ajtmh.15-0746>.
- Duong, Thanh Hai, et Dominique Richard-Lenoble. 2008. « Diagnostic des parasitoses à parasites sanguicoles ». *Revue Francophone des Laboratoires* 2008 (399): 29-39.
[https://doi.org/10.1016/S1773-035X\(08\)70180-1](https://doi.org/10.1016/S1773-035X(08)70180-1).
- Ehrens, Alexandra, Achim Hoerauf, et Marc P. Hübner. 2022. « Eosinophils in filarial infections: Inducers of protection or pathology? » *Frontiers in Immunology* 13 (octobre):983812. <https://doi.org/10.3389/fimmu.2022.983812>.
- El Setouhy, Maged, Reda M. R. Ramzy, Ehab S. Ahmed, Amr M. Kandil, Omar Hussain, Hoda A. Farid, Hanan Helmy, et Gary J. Weil. 2004. « A Randomized Clinical Trial Comparing Single- and Multi-Dose Combination Therapy with Diethylcarbamazine and Albendazole for Treatment of Bancroftian Filariasis ». *The American Journal of Tropical Medicine and Hygiene* 70 (2): 191-96.
- Emukah, Emmanuel, Lindsay J. Rakers, Barminas Kahansim, Emmanuel S. Miri, Bertram E. B. Nwoke, Emily Griswold, Yisa Saka, et al. 2018. « In Southern Nigeria Loa loa Blood Microfilaria Density is Very Low Even in Areas with High Prevalence of Loiasis: Results of a Survey Using the New LoaScope Technology ». *The American Journal of Tropical Medicine and Hygiene* 99 (1): 116-23. <https://doi.org/10.4269/ajtmh.18-0163>.
- Erickson, Sara M., Edward K. Thomsen, John B. Keven, Naomi Vincent, Gussy Koimbu, Peter M. Siba, Bruce M. Christensen, et Lisa J. Reimer. 2013. « Mosquito-Parasite Interactions Can Shape Filariasis Transmission Dynamics and Impact Elimination

Programs ». Édité par Jesus G. Valenzuela. *PLoS Neglected Tropical Diseases* 7 (9): e2433. <https://doi.org/10.1371/journal.pntd.0002433>.

Famakinde, Damilare. 2018. « Mosquitoes and the Lymphatic Filarial Parasites: Research Trends and Budding Roadmaps to Future Disease Eradication ». *Tropical Medicine and Infectious Disease* 3 (1): 4. <https://doi.org/10.3390/tropicalmed3010004>.

Faye, O., B. Faye, J. L. Ndiaye, R. C. K. Tine, D. Ndiaye, et A. Haydara Mbacké. 2009. « [Epidemiological study on lymphatic filariasis (*Wuchereria bancrofti*) in Sénégal] ». *Medecine Tropicale: Revue Du Corps De Sante Colonial* 69 (1): 101-2.

Ferreira, Marcelo U., James Lee Crainey, et Federico G. Gobbi. 2023. « The Search for Better Treatment Strategies for Mansonellosis: An Expert Perspective ». *Expert Opinion on Pharmacotherapy* 24 (15): 1685-92. <https://doi.org/10.1080/14656566.2023.2240235>.

Ferreira, Marcelo U., James Lee Crainey, et Sérgio L.B. Luz. 2021. « *Mansonella Ozzardi* ». *Trends in Parasitology* 37 (1): 90-91. <https://doi.org/10.1016/j.pt.2020.03.005>.

Frédéric, NAHMIAS. 2018. « Haute Autorité de santé ».

Gaillard, Colette Marie, Sebastien David Pion, Hadjira Hamou, Constant Sirima, Charlotte Bizet, Thomas Lemarcis, Jules Rodrigues, et al. 2020. « Detection of DNA of *Filariae* Closely Related to *Mansonella Perstans* in Faecal Samples from Wild Non-Human Primates from Cameroon and Gabon ». *Parasites & Vectors* 13 (1): 313. <https://doi.org/10.1186/s13071-020-04184-1>.

Gardon, Jacques, Nathalie Gardon-Wendel, Demanga-Ngangue, Joseph Kamgno, Jean-Philippe Chippaux, et Michel Boussinesq. 1997. « Serious Reactions after Mass Treatment of Onchocerciasis with Ivermectin in an Area Endemic for Loa Loa Infection ». *The Lancet* 350 (9070): 18-22. [https://doi.org/10.1016/S0140-6736\(96\)11094-1](https://doi.org/10.1016/S0140-6736(96)11094-1).

Geelhand De Merxem, D., J. N. Siewe Fodjo, S. Menon, A. Hotterbeekx, et R. Colebunders. 2019. « Nodding Syndrome Research, Lessons Learned from the NSETHIO Project ». *Global Mental Health* 6:e26. <https://doi.org/10.1017/gmh.2019.24>.

Gentilini, M., J.M. Pinon, et M. Danis. 1973. « Diagnostic biologique des filarioses humaines ». *Médecine et Maladies Infectieuses* 3 (8-9): 351-53. [https://doi.org/10.1016/S0399-077X\(73\)80130-1](https://doi.org/10.1016/S0399-077X(73)80130-1).

Gobbi, Federico, Anna Beltrame, Dora Buonfrate, Silvia Staffolani, Monica Degani, Maria Gobbo, Andrea Angheben, Stefania Marocco, et Zeno Bisoffi. 2017. « Imported Infections with *Mansonella perstans* Nematodes, Italy ». *Emerging Infectious Diseases* 23 (9): 1539-42. <https://doi.org/10.3201/eid2309.170263>.

813 González, Mikel A., Erin Dilger, María M. Ronderos, Gustavo R. Spinelli, Orin Courtenay, et
814 James G. C. Hamilton. 2020. « Significant Reduction in Abundance of Peridomestic
815 Mosquitoes (Culicidae) and Culicoides Midges (Ceratopogonidae) after Chemical
816 Intervention in Western São Paulo, Brazil ». *Parasites & Vectors* 13 (1): 549.
817 <https://doi.org/10.1186/s13071-020-04427-1>.

818 Gyasi, Michael E., Ogugua N. Okonkwo, et Koushik Tripathy. 2024. « Onchocerciasis ». In
819 *StatPearls*. Treasure Island (FL): StatPearls Publishing.
820 <http://www.ncbi.nlm.nih.gov/books/NBK559027/>.

821 Haddix, Anne C., et Andrew Kestler. 2000. « Lymphatic Filariasis: Economic Aspects of the
822 Disease and Programmes for Its Elimination ». *Transactions of the Royal Society of*
823 *Tropical Medicine and Hygiene* 94 (6): 592-93. [https://doi.org/10.1016/S0035-](https://doi.org/10.1016/S0035-9203(00)90199-8)
824 [9203\(00\)90199-8](https://doi.org/10.1016/S0035-9203(00)90199-8).

825 Harrup, Fabio Macchioni, Luca Chelucci, et Beatrice Torracca. 2016. « Advances in Control
826 Techniques for Culicoides and Future Prospects ». *Veterinaria Italiana* 52 (3): 247-64.
827 <https://doi.org/10.12834/VetIt.741.3602.3>.

828 Hartley, Marie. 2021. « Onchocerciasis (river blindness) », 2021.
829 <https://dermnetnz.org/topics/onchocerciasis>.

830 Herbeuval, H., R. Herbeuval, et J. Duheille. 1962. « [The passage, in the circulating blood, of
831 hematopoietic masses, erythroblastic islands, megakaryoblastic islands, mixed
832 islands] ». *Nouvelle Revue Francaise D'hematologie* 2:616-19.

833 Herricks, Jennifer R., Peter J. Hotez, Valentine Wanga, Luc E. Coffeng, Juanita A. Haagsma,
834 María-Gloria Basáñez, Geoffrey Buckle, et al. 2017. « The Global Burden of Disease
835 Study 2013: What Does It Mean for the NTDs? » Édité par Xiao-Nong Zhou. *PLOS*
836 *Neglected Tropical Diseases* 11 (8): e0005424.
837 <https://doi.org/10.1371/journal.pntd.0005424>.

838 Hotez, Peter J., et Aruna Kamath. 2009. « Neglected Tropical Diseases in Sub-Saharan Africa:
839 Review of Their Prevalence, Distribution, and Disease Burden ». Édité par Michael
840 Cappello. *PLoS Neglected Tropical Diseases* 3 (8): e412.
841 <https://doi.org/10.1371/journal.pntd.0000412>.

842 Hotez, Peter J., David H. Molyneux, Alan Fenwick, Jacob Kumaresan, Sonia Ehrlich Sachs,
843 Jeffrey D. Sachs, et Lorenzo Savioli. 2007. « Control of Neglected Tropical Diseases ». *New*
844 *England Journal of Medicine* 357 (10): 1018-27.
845 <https://doi.org/10.1056/NEJMra064142>.

- Ibe, Ogochukwu, Obinna Onwujekwe, Benjamin Uzochukwu, Miriam Ajuba, et Paul Okonkwo. 2015. « Exploring Consumer Perceptions and Economic Burden of Onchocerciasis on Households in Enugu State, South-East Nigeria ». Édité par Samuel V. Scarpino. *PLOS Neglected Tropical Diseases* 9 (11): e0004231. <https://doi.org/10.1371/journal.pntd.0004231>.
- Jacobsen, Kathryn H, Bailey C Andress, Elina A Bhagwat, Ciera A Bryant, Vipusha R Chandrapu, Caroline G Desmonts, Tania M Matthews, Akeem Ogunkoya, Tristan J Wheeler, et A'kayla S Williams. 2022. « A Call for Loiasis to Be Added to the WHO List of Neglected Tropical Diseases ». *The Lancet Infectious Diseases* 22 (10): e299-302. [https://doi.org/10.1016/S1473-3099\(22\)00064-0](https://doi.org/10.1016/S1473-3099(22)00064-0).
- Jain, Rajeev, Jern Y. Chen, Andrew R. Butcher, Robert Casson, et Dinesh Selva. 2008. « Subconjunctival Loa Loa Worm ». *International Journal of Infectious Diseases* 12 (6): e133-35. <https://doi.org/10.1016/j.ijid.2008.03.001>.
- James, Spencer L, Degu Abate, Kalkidan Hassen Abate, Solomon M Abay, Cristiana Abbafati, Nooshin Abbasi, Hedayat Abbastabar, et al. 2018. « Global, Regional, and National Incidence, Prevalence, and Years Lived with Disability for 354 Diseases and Injuries for 195 Countries and Territories, 1990–2017: A Systematic Analysis for the Global Burden of Disease Study 2017 ». *The Lancet* 392 (10159): 1789-1858. [https://doi.org/10.1016/S0140-6736\(18\)32279-7](https://doi.org/10.1016/S0140-6736(18)32279-7).
- Kamgno, Joseph, Sébastien D. Pion, Cédric B. Chesnais, Matthew H. Bakalar, Michael V. D'Ambrosio, Charles D. Mackenzie, Hugues C. Nana-Djeunga, et al. 2017. « A Test-and-Not-Treat Strategy for Onchocerciasis in *Loa Loa* –Endemic Areas ». *New England Journal of Medicine* 377 (21): 2044-52. <https://doi.org/10.1056/NEJMoa1705026>.
- Katawa, Gnatoulma, Laura E. Layland, Alex Y. Debrah, Charlotte Von Horn, Linda Batsa, Alexander Kwarteng, Sandra Arriens, et al. 2015. « Hyperreactive Onchocerciasis Is Characterized by a Combination of Th17-Th2 Immune Responses and Reduced Regulatory T Cells ». Édité par Benjamin L. Makepeace. *PLoS Neglected Tropical Diseases* 9 (1): e3414. <https://doi.org/10.1371/journal.pntd.0003414>.
- Katiyar, D., et L. K. Singh. 2011. « Filariasis: Current Status, Treatment and Recent Advances in Drug Development ». *Current Medicinal Chemistry* 18 (14): 2174-85. <https://doi.org/10.2174/092986711795656234>.
- Keiser, Paul B., Yaya I. Coulibaly, Falaye Keita, Diakaridia Traore, Abdallah Diallo, Dapa A. Diallo, Roshanak T. Semnani, et al. 2003. « Clinical Characteristics of Post-Treatment Reactions to Ivermectin/Albendazole for *Wuchereria Bancrofti* in a Region Co-

880 Endemic for *Mansonella Perstans* ». *The American Journal of Tropical Medicine and*
881 *Hygiene* 69 (3): 331-35.

882 Keiser, Paul B., Yaya Coulibaly, Joseph Kubofcik, Abdallah A. Diallo, Amy D. Klion, Sekou
883 F. Traoré, et Thomas B. Nutman. 2008. « Molecular Identification of *Wolbachia* from
884 the Filarial Nematode *Mansonella Perstans* ». *Molecular and Biochemical Parasitology*
885 160 (2): 123-28. <https://doi.org/10.1016/j.molbiopara.2008.04.012>.

886 Keita, Mamadou Fodé. 1979. « Aspects épidémiologiques des filarioses à *onchocerca volvulus*,
887 *wuchereria bancrofti* et *tetrapetatonema perstans* ». Thesis, Ecole Nationale de
888 Médecine et de Pharmacie. <https://www.bibliosante.ml/handle/123456789/9605>.

889 Kelly-Hope, Louise, Rossely Paulo, Brent Thomas, Miguel Brito, Thomas R. Unnasch, et
890 David Molyneux. 2017. « *Loa Loa* Vectors *Chrysops* Spp.: Perspectives on Research,
891 Distribution, Bionomics, and Implications for Elimination of Lymphatic Filariasis and
892 *Onchocerciasis* ». *Parasites & Vectors* 10 (1): 172. [https://doi.org/10.1186/s13071-017-](https://doi.org/10.1186/s13071-017-2103-y)
893 2103-y.

894 Khan, A. M., P. Dutta, S. Das, A. K. Pathak, P. Sarmah, M. E. Hussain, et J. Mahanta. 2015.
895 « Microfilarial Periodicity of *Wuchereria Bancrofti* in Assam, Northeast India ». *Journal of Vector Borne Diseases* 52 (3): 208-12.

897 Kima, A., K.T. Guiguemde, M. Serme, Z.C. Meda, R. Bougma, J.P. Djiatsa, C. Bougouma, et
898 F. Drabo. 2021. « Évaluation de la Transmission de la Filariose Lymphatique au
899 Burkina Faso: À Propos de 4 Districts Sanitaires ». *Meédecine Tropicale et Santéé*
900 *Internationale* 1 (2): mtsibulletin.n1.2021.83.
901 <https://doi.org/10.48327/mtsibulletin.n1.2021.83>.

902 Knopp, Stefanie, Peter Steinmann, Christoph Hatz, Jennifer Keiser, et Jürg Utzinger. 2012.
903 « Nematode Infections »: *Infectious Disease Clinics of North America* 26 (2): 359-81.
904 <https://doi.org/10.1016/j.idc.2012.02.005>.

905 Komlan, Kossi, Patrick S. Vossberg, Richard G. Gantim, Tchalim Solim, Francois Korbmacher,
906 Méba Banla, Koffi Padjoudoum, Potchoziou Karabou, Carsten Köhler, et Peter T.
907 Soboslay. 2018. « *Onchocerca Volvulus* Infection and Serological Prevalence, Ocular
908 *Onchocerciasis* and Parasite Transmission in Northern and Central Togo after Decades
909 of *Simulium Damnosum* s.l. Vector Control and Mass Drug Administration of
910 Ivermectin ». Édité par Matthew C. Freeman. *PLOS Neglected Tropical Diseases* 12
911 (3): e0006312. <https://doi.org/10.1371/journal.pntd.0006312>.

912 Kwarteng, Alexander, Samuel Terkper Ahuno, et Freda Osei Akoto. 2016. « Killing Filarial
 913 Nematode Parasites: Role of Treatment Options and Host Immune Response ». *Infectious Diseases of Poverty* 5 (1): 86. <https://doi.org/10.1186/s40249-016-0183-0>.
 914
 915 Laidoudi, Younes, Jean-Lou Marie, Djamel Tahir, Stéphanie Watier-Grillot, Oleg
 916 Mediannikov, et Bernard Davoust. 2020. « Detection of Canine Vector-Borne Filariasis
 917 and Their Wolbachia Endosymbionts in French Guiana ». *Microorganisms* 8 (5): 770.
 918 <https://doi.org/10.3390/microorganisms8050770>.
 919 Lima, Nathália F., Cecilia A. Veggiani Aybar, María J. Dantur Juri, et Marcelo U. Ferreira.
 920 2016. « *Mansonella Ozzardi* : A Neglected New World Filarial Nematode ». *Pathogens*
 921 *and Global Health* 110 (3): 97-107. <https://doi.org/10.1080/20477724.2016.1190544>.
 922 Lustigman, Sara, Alexandra Grote, et Elodie Ghedin. 2017. « The Role of “omics” in the Quest
 923 to Eliminate Human Filariasis ». Édité par Judd L. Walson. *PLOS Neglected Tropical*
 924 *Diseases* 11 (4): e0005464. <https://doi.org/10.1371/journal.pntd.0005464>.
 925 M. Boussinesq. 2007. « Actualités thérapeutiques dans les filarioses – Recent advances in the
 926 treatment of filariases ». *mise au point*, n° 2, 55.
 927 Mandal, Debasish, Dipankar Roy, D. K. Bera, et Buddhadeb Manna. 2012. « Occurrence of
 928 Gravid Loa Loa in Subconjunctival Space of Man: A Case Report from West Bengal,
 929 India ». *Journal of Parasitic Diseases*, juillet. [https://doi.org/10.1007/s12639-012-](https://doi.org/10.1007/s12639-012-0130-3)
 930 [0130-3](https://doi.org/10.1007/s12639-012-0130-3).
 931 Manguin, S., M.J. Bangs, J. Pothikasikorn, et T. Chareonviriyaphap. 2010. « Review on Global
 932 Co-Transmission of Human Plasmodium Species and Wuchereria Bancrofti by
 933 Anopheles Mosquitoes ». *Infection, Genetics and Evolution* 10 (2): 159-77.
 934 <https://doi.org/10.1016/j.meegid.2009.11.014>.
 935 McGarry, Helen F, Ken Pfarr, Gill Egerton, Achim Hoerauf, Jean-Paul Akue, Peter Enyong,
 936 Samuel Wanji, et al. 2003. « [No title found] ». *Filaria Journal* 2 (1): 9.
 937 <https://doi.org/10.1186/1475-2883-2-9>.
 938 Meda, N., M.A. Daboue, W.P. Djiguimde, C. Meda, et S. Konate/Ouedraogo. 2009. « 571
 939 Manifestations oculaires de la loase ». *Journal Français d’Ophtalmologie* 32
 940 (avril):1S173. [https://doi.org/10.1016/S0181-5512\(09\)73695-2](https://doi.org/10.1016/S0181-5512(09)73695-2).
 941 Medeiros, Jansen Fernandes, James Lee Crainey, Felipe Arley Costa Pessoa, et Sérgio Luis
 942 Bessa Luz. 2017. « Mansonelliasis ». In *Arthropod Borne Diseases*, édité par Carlos
 943 Brisola Marcondes, 405-26. Cham: Springer International Publishing.
 944 https://doi.org/10.1007/978-3-319-13884-8_26.

- Mediannikov, O., et S. Ranque. 2018a. « Mansonellosis, the Most Neglected Human Filariasis ». *New Microbes and New Infections* 26 (novembre):S19-22. <https://doi.org/10.1016/j.nmni.2018.08.016>.
- . 2018b. « Mansonellosis, the Most Neglected Human Filariasis ». *New Microbes and New Infections* 26 (novembre):S19-22. <https://doi.org/10.1016/j.nmni.2018.08.016>.
- Melrose, Wayne D. 2002. « Lymphatic Filariasis: New Insights into an Old Disease ». *International Journal for Parasitology* 32 (8): 947-60. [https://doi.org/10.1016/S0020-7519\(02\)00062-0](https://doi.org/10.1016/S0020-7519(02)00062-0).
- Mendoza, Natalia, Alexa Li, Amber Gill, et Stephen Tyring. 2009. « Filariasis: Diagnosis and Treatment ». *Dermatologic Therapy* 22 (6): 475-90. <https://doi.org/10.1111/j.1529-8019.2009.01271.x>.
- Métais, Alice, Sophie Michalak, et Audrey Rousseau. 2021. « Albendazole-Related Loa Loa Encephalopathy ». *IDCases* 23:e01033. <https://doi.org/10.1016/j.idcr.2020.e01033>.
- Metzger, Wolfram Gottfried, et Benjamin Mordmüller. 2014. « Loa Loa—Does It Deserve to Be Neglected? » *The Lancet Infectious Diseases* 14 (4): 353-57. [https://doi.org/10.1016/S1473-3099\(13\)70263-9](https://doi.org/10.1016/S1473-3099(13)70263-9).
- Molyneux, David H., et Mark J. Taylor. 2001. « Current Status and Future Prospects of the Global Lymphatic Filariasis Programme ». *Current Opinion in Infectious Diseases* 14 (2): 155-59. <https://doi.org/10.1097/00001432-200104000-00008>.
- Mourembou, Gaël, Florence Fenollar, Jean Bernard Lekana-Douki, Angelique Ndjoi Mbiguino, Sydney Maghendji Nzondo, Pierre Blaise Matsiegui, Rella Zoleko Manego, et al. 2015a. « Mansonella, Including a Potential New Species, as Common Parasites in Children in Gabon ». Édité par Dunstan Mukoko. *PLOS Neglected Tropical Diseases* 9 (10): e0004155. <https://doi.org/10.1371/journal.pntd.0004155>.
- . 2015b. « Mansonella, including a Potential New Species, as Common Parasites in Children in Gabon ». *PLoS Neglected Tropical Diseases* 9 (10): e0004155. <https://doi.org/10.1371/journal.pntd.0004155>.
- Ngwewondo, Adela, Ivan Scandale, et Sabine Specht. 2021. « Onchocerciasis Drug Development: From Preclinical Models to Humans ». *Parasitology Research* 120 (12): 3939-64. <https://doi.org/10.1007/s00436-021-07307-4>.
- Nizeyimana, Jean Baptiste. 2021. « ÉTUDE DES CONDITIONS DE DEPISTAGE CLINIQUE ET LES EFFETS PSYCHO SOCIOECONOMIQUE DES PERSONNES PORTEUSES DE L'ELEPHANTIASIS DES JAMBES. CAS DE

L'ARRONDISSEMENT DE GUIDIGUIS CAS DE LA REGION DE L'EXTREME-NORD CAMEROUN DE 2017-2020 Par » 26 (03).

Njim, Tsi, Joel Mbigha Ngum, et Leopold Ndemnge Aminde. 2015. « Cutaneous Onchocerciasis in Dumbu, a Pastoral Area in the North-West Region of Cameroon: Diagnostic Challenge and Socio-Economic Implications ». *Pan African Medical Journal* 22. <https://doi.org/10.11604/pamj.2015.22.298.7707>.

Njoku, Chidiebere, Lawrence Nwankwo, Masayo Ozaki, Andrew Obasi, Chinyere N. Ukaga, Emmanuel Emukah, Kenrick Nwodu, et al. 2013. « Community-Wide Distribution of Long-Lasting Insecticidal Nets Can Halt Transmission of Lymphatic Filariasis in Southeastern Nigeria ». *The American Journal of Tropical Medicine and Hygiene* 89 (3): 578-87. <https://doi.org/10.4269/ajtmh.12-0775>.

Noroës, Joaquim, David Addiss, Agnaldo Cedenho, Jose Figueredo-Silvan, Guilherme Lima, et Gerusa Dreyer. 2003. « Pathogenesis of Filarial Hydrocele: Risk Associated with Intrascrotal Nodules caused by Death of Adult Wuchereria Bancrofti ». *Transactions of the Royal Society of Tropical Medicine and Hygiene* 97 (5): 561-66. [https://doi.org/10.1016/S0035-9203\(03\)80029-9](https://doi.org/10.1016/S0035-9203(03)80029-9).

Nutman, Thomas B. 2013. « Insights into the Pathogenesis of Disease in Human Lymphatic Filariasis ». *Lymphatic Research and Biology* 11 (3): 144-48. <https://doi.org/10.1089/lrb.2013.0021>.

O'Hanlon, Simon J., Hannah C. Slater, Robert A. Cheke, Boakye A. Boatın, Luc E. Coffeng, Sébastien D. S. Pion, Michel Boussinesq, Honorat G. M. Zouré, Wilma A. Stolk, et María-Gloria Basáñez. 2016. « Model-Based Geostatistical Mapping of the Prevalence of Onchocerca Volvulus in West Africa ». Édité par Ricardo J. Soares Magalhaes. *PLOS Neglected Tropical Diseases* 10 (1): e0004328. <https://doi.org/10.1371/journal.pntd.0004328>.

Ojurongbe, Olusola, Akeem Abiodun Akindele, Monsuru Adebayo Adeleke, Matthew Oyebo Oyebo, Samuel Adeyinka Adedokun, Josephine Folashade Ojo, Callistus Adewale Akinleye, Oloyede Samuel Bolaji, Olusegun Adelowo Adefioye, et Oluwaseyi Adegboyega Adeyeba. 2015. « Co-endemicity of Loiasis and Onchocerciasis in Rain Forest Communities in Southwestern Nigeria ». *PLOS Neglected Tropical Diseases* 9 (3): e0003633. <https://doi.org/10.1371/journal.pntd.0003633>.

« OMS MALI Rapport annuel 2018.pdf ». s. d. Consulté le 24 juillet 2024. <https://www.afro.who.int/sites/default/files/2019-01/OMS%20MALI%20Rapport%20annuel%202018.pdf>.

Opoku, Millicent, Corrado Minetti, Worlasi D. Kartey-Attipoe, Sampson Otoo, Joseph Otchere, Bruno Gomes, Dziedzom K. De Souza, et Lisa J. Reimer. 2018. « An Assessment of Mosquito Collection Techniques for Xenomonitoring of Anopheline-Transmitted Lymphatic Filariasis in Ghana ». *Parasitology* 145 (13): 1783-91. <https://doi.org/10.1017/S0031182018000938>.

Ottesen, Eric A. 2000. « Editorial: The Global Programme to Eliminate Lymphatic Filariasis ». *Tropical Medicine & International Health* 5 (9): 591-94. <https://doi.org/10.1046/j.1365-3156.2000.00620.x>.

———. 2006. « Lymphatic Filariasis: Treatment, Control and Elimination ». In *Advances in Parasitology*, 61:395-441. Elsevier. [https://doi.org/10.1016/S0065-308X\(05\)61010-X](https://doi.org/10.1016/S0065-308X(05)61010-X).

Ouedraogo, Amina Nomtondo, Emile Bégounriti Somda, Fagnima Traoré, Muriel Sidnoma Ouédraogo, Gilbert Patrice Tapsoba, Angèle Ouangre/Ouédraogo, Salamata Bara, et al. 2016. « Impact du traitement de masse de la filariose lymphatique par l'albendazole-ivermectine en zone de savane : cas de la région de l'Est du Burkina ». *HEALTH SCIENCES AND DISEASE* 17 (4). <http://hsd-fmsb.org/index.php/hsd/article/view/731>.

Pampiglione, S., G. Canestri Trotti, et F. Rivasi. 1995. « Human Dirofilaria Due to Dirofilaria (Noctiella) Repens: A Review of World Literature ». *Parassitologia* 37 (2-3): 149-93.

Pérez-Molina, José Antonio, Marta Díaz-Menéndez, Ana Pérez-Ayala, Federico Ferrere, Begoña Monje, Francesca Norman, et Rogelio López-Vélez. 2010. « Tratamiento de las enfermedades causadas por parásitos ». *Enfermedades Infecciosas y Microbiología Clínica* 28 (1): 44-59. <https://doi.org/10.1016/j.eimc.2009.11.003>.

Phillips, Richard O., Michael Frimpong, Fred S. Sarfo, Birte Kretschmer, Marcus Beissner, Alexander Debrah, Yaw Ampem-Amoako, et al. 2014. « Infection with *Mansonella perstans* Nematodes in Buruli Ulcer Patients, Ghana ». *Emerging Infectious Diseases* 20 (6): 1000-1003. <https://doi.org/10.3201/eid2006.131501>.

Pion, S. D. S., J. A. N. Filipe, J. Kamgno, J. Gardon, M.-G. Basáñez, et M. Boussinesq. 2006. « Microfilarial Distribution of Loa Loa in the Human Host: Population Dynamics and Epidemiological Implications ». *Parasitology* 133 (01): 101. <https://doi.org/10.1017/S0031182006000035>.

Pion, Sébastien Ds, Hugues Nana-Djeunga, Yannick Niamsi-Emalio, Cédric B Chesnais, Hugo Deléglise, Charles Mackenzie, Wilma Stolk, et al. 2020. « Implications for Annual Retesting after a Test-and-Not-Treat Strategy for Onchocerciasis Elimination in Areas

- Co-Endemic with Loa Loa Infection: An Observational Cohort Study ». *The Lancet Infectious Diseases* 20 (1): 102-9. [https://doi.org/10.1016/S1473-3099\(19\)30554-7](https://doi.org/10.1016/S1473-3099(19)30554-7).
- Poole, Catherine B., Zhiru Li, Andy Alhassan, Dylan Guelig, Steven Diesburg, Nathan A. Tanner, Yinhua Zhang, et al. 2017. « Colorimetric Tests for Diagnosis of Filarial Infection and Vector Surveillance Using Non-Instrumented Nucleic Acid Loop-Mediated Isothermal Amplification (NINA-LAMP) ». Édité par Matty Knight. *PLOS ONE* 12 (2): e0169011. <https://doi.org/10.1371/journal.pone.0169011>.
- Porter, Suzanne, et Adrian Hopkins. 2015. « Élimination de l'onchocercose et de la filariose lymphatique » 12.
- Puente, Sabino, Mar Lago, Mercedes Subirats, Ismael Sanz-Esteban, Marta Arsuaga, Belén Vicente, Montserrat Alonso-Sardon, Moncef Belhassen-Garcia, et Antonio Muro. 2020. « Imported Mansonella perstans infection in Spain ». *Infectious Diseases of Poverty* 9 (1): 105. <https://doi.org/10.1186/s40249-020-00729-9>.
- Ramaiah, K. D., et Eric A. Ottesen. 2014. « Progress and Impact of 13 Years of the Global Programme to Eliminate Lymphatic Filariasis on Reducing the Burden of Filarial Disease ». Édité par Moses Bockarie. *PLoS Neglected Tropical Diseases* 8 (11): e3319. <https://doi.org/10.1371/journal.pntd.0003319>.
- Ramharter, Michael, Joanna Butler, Ghyslain Mombo-Ngoma, Tamara Nordmann, Saskia Dede Davi, et Rella Zoleko Manego. 2023. « The African Eye Worm: Current Understanding of the Epidemiology, Clinical Disease, and Treatment of Loiasis ». *The Lancet Infectious Diseases*, octobre, S1473309923004383. [https://doi.org/10.1016/S1473-3099\(23\)00438-3](https://doi.org/10.1016/S1473-3099(23)00438-3).
- Rebollo, Maria P., et Moses J. Bockarie. 2017. « Can Lymphatic Filariasis Be Eliminated by 2020? » *Trends in Parasitology* 33 (2): 83-92. <https://doi.org/10.1016/j.pt.2016.09.009>.
- Richardson, Eugene T., Robert Luo, Doran L. Fink, Thomas B. Nutman, John K. Geisse, et Michele Barry. 2012. « Transient Facial Swellings in a Patient With a Remote African Travel History ». *Journal of Travel Medicine* 19 (3): 183-85. <https://doi.org/10.1111/j.1708-8305.2012.00612.x>.
- Risch, Frederic, Manuel Ritter, Achim Hoerauf, et Marc P. Hübner. 2021. « Human Filariasis—Contributions of the Litomosoides Sigmodontis and Acanthocheilonema Viteae Animal Model ». *Parasitology Research* 120 (12): 4125-43. <https://doi.org/10.1007/s00436-020-07026-2>.
- Robert, Vincent, E.H. Ndiaye, Nil Rahola, Gilbert Le Goff, Philippe Boussès, D. Diallo, Voahirana Le Goff, Lauriane Mariamé, et M. Diallo. 2022. « Clés dichotomiques

- illustrées d'identification des femelles et des larves de moustiques (Diptera : Culicidae) du Burkina Faso, Cap-Vert, Gambie, Mali, Mauritanie, Niger, Sénégal et Tchad », 181 p. <https://doi.org/10.23708/FDI:010084866>.
- Rodi, Miriam, Caspar Gross, Thaisa Lucas Sandri, Lilith Berner, Marina Marcet-Houben, Ersoy Kocak, Michaela Pogoda, et al. 2023a. « Whole genome analysis of two sympatric human *Mansonella*: *Mansonella perstans* and *Mansonella* sp “DEUX” ». *Frontiers in Cellular and Infection Microbiology* 13 (avril):1159814. <https://doi.org/10.3389/fcimb.2023.1159814>.
- . 2023b. « Whole Genome Analysis of Two Sympatric Human *Mansonella*: *Mansonella* *Perstans* and *Mansonella* Sp “DEUX” ». *Frontiers in Cellular and Infection Microbiology* 13:1159814. <https://doi.org/10.3389/fcimb.2023.1159814>.
- Sandri, Thaisa Lucas, Andrea Kreidenweiss, Simon Cavallo, David Weber, Sascha Juhas, Miriam Rodi, Tamirat Gebru Woldearegai, et al. 2021. « Molecular Epidemiology of *Mansonella* Species in Gabon ». *The Journal of Infectious Diseases* 223 (2): 287-96. <https://doi.org/10.1093/infdis/jiaa670>.
- Sangare, Moussa Brema, Yaya Ibrahim Coulibaly, Siaka Yamoussa Coulibaly, Michel Emmanuel Coulibaly, Bourama Traore, Ilo Dicko, Ibrahim Moussa Sissoko, et al. 2018. « A cross-sectional study of the filarial and *Leishmania* co-endemicity in two ecologically distinct settings in Mali ». *Parasites & Vectors* 11 (1): 18. <https://doi.org/10.1186/s13071-017-2531-8>.
- Sawadogo, Pm. 2019. « >Onchocercose au Burkina Faso : une endémie à transmission vectorielle en voie de réémergence ». *Bulletin de la société de pathologie exotique* 112 (5): 260-74. <https://doi.org/10.3166/bspe-2020-0106>.
- Shelley, Aj, et S Coscarón. 2001. « Simuliid blackflies (Diptera: Simuliidae) and ceratopogonid midges (Diptera: Ceratopogonidae) as vectors of *Mansonella ozzardi* (Nematoda: Onchocercidae) in northern Argentina ». *Memórias do Instituto Oswaldo Cruz* 96 (4): 451-58. <https://doi.org/10.1590/S0074-02762001000400003>.
- Shenoy, R. K. 2008. « Clinical and Pathological Aspects of Filarial Lymphedema and Its Management ». *The Korean Journal of Parasitology* 46 (3): 119. <https://doi.org/10.3347/kjp.2008.46.3.119>.
- Simón, Fernando, Mar Siles-Lucas, Rodrigo Morchón, Javier González-Miguel, Isabel Mellado, Elena Carretón, et Jose Alberto Montoya-Alonso. 2012. « Human and Animal Dirofilariasis: The Emergence of a Zoonotic Mosaic ». *Clinical Microbiology Reviews* 25 (3): 507-44. <https://doi.org/10.1128/CMR.00012-12>.

1113 Simonsen, Paul E., Yahya A. Derua, Stephen M. Magesa, Erling M. Pedersen, Anna-Sofie
 1114 Stensgaard, Mwelecele N. Malecela, et William N. Kisinza. 2014. « Lymphatic
 1115 filariasis control in Tanga Region, Tanzania: status after eight rounds of mass drug
 1116 administration ». *Parasites & Vectors* 7 (1): 507. [https://doi.org/10.1186/s13071-014-](https://doi.org/10.1186/s13071-014-0507-5)
 1117 0507-5.

1118 Simonsen, Paul E., Ambrose W. Onapa, et Santa Maria Asio. 2011a. « Mansonella Perstans
 1119 Filariasis in Africa ». *Acta Tropica*, The Diagnostics and Control of Neglected Tropical
 1120 Helminth Diseases, 120 (septembre):S109-20.
 1121 <https://doi.org/10.1016/j.actatropica.2010.01.014>.

1122 ———. 2011b. « Mansonella Perstans Filariasis in Africa ». *Acta Tropica* 120 Suppl 1
 1123 (septembre):S109-120. <https://doi.org/10.1016/j.actatropica.2010.01.014>.

1124 Small, Scott T., Daniel J. Tisch, et Peter A. Zimmerman. 2014. « Molecular Epidemiology,
 1125 Phylogeny and Evolution of the Filarial Nematode Wuchereria Bancrofti ». *Infection,
 1126 Genetics and Evolution* 28 (décembre):33-43.
 1127 <https://doi.org/10.1016/j.meegid.2014.08.018>.

1128 Supali, Taniawati, Is S. Ismid, Heri Wibowo, Yenny Djuardi, Esther Majawati, Praba Ginanjar,
 1129 et Peter Fischer. 2006. « Estimation of the Prevalence of Lymphatic Filariasis by a Pool
 1130 Screen PCR Assay Using Blood Spots Collected on Filter Paper ». *Transactions of the
 1131 Royal Society of Tropical Medicine and Hygiene* 100 (8): 753-59.
 1132 <https://doi.org/10.1016/j.trstmh.2005.10.005>.

1133 Tahita, Marc Christian, Thuy-Huong Ta-Tang, Bérénger Kaboré, Raquel Capote-Morales,
 1134 Irene Molina De La Fuente, Raquel Cruces, Hamidou Ilboudo, et al. 2024. « Molecular
 1135 Detection of Wuchereria Bancrofti, Loa Loa and Mansonella Perstans from Dried Blood
 1136 Spots Taken from Pregnant Women in Rural Burkina Faso ». *African Journal of
 1137 Parasitology, Mycology and Entomology* 1 (2): 1.
 1138 <https://doi.org/10.35995/ajpme2010002>.

1139 Tamarozzi, Francesca, Paola Rodari, Joaquín Salas-Coronas, Emmanuel Bottieau, Fernando
 1140 Salvador, Manuel Jesús Soriano-Pérez, María Isabel Cabeza-Barrera, et al. 2022. « A
 1141 Large Case Series of Travel-Related *Mansonella Perstans* (Vector-Borne Filarial
 1142 Nematode): A TropNet Study in Europe ». *Journal of Travel Medicine*, avril, taac048.
 1143 <https://doi.org/10.1093/jtm/taac048>.

1144 Ta-Tang, Thuy-Huong, James Crainey, Rory J. Post, Sergio LB. Luz, et José Rubio. 2018.
 1145 « Mansonellosis: Current Perspectives ». *Research and Reports in Tropical Medicine*
 1146 Volume 9 (janvier):9-24. <https://doi.org/10.2147/RRTM.S125750>.

1147 Ta-Tang, Thuy-Huong, Sergio L B Luz, James L Crainey, et José M Rubio. 2021a. « An
1148 Overview of the Management of Mansonellosis ». *Research and Reports in Tropical*
1149 *Medicine* 12 (mai):93-105. <https://doi.org/10.2147/RRTM.S274684>.

1150 Ta-Tang, Thuy-Huong, Sergio Lb Luz, James L Crainey, et José M Rubio. 2021b. « An
1151 Overview of the Management of Mansonellosis ». *Research and Reports in Tropical*
1152 *Medicine* Volume 12 (mai):93-105. <https://doi.org/10.2147/RRTM.S274684>.

1153 Tatuene, J. Kamtchum, R. Gounoue Fotsing, T. Nkoa, JB Tchateng Mbougua, HC Nana
1154 Djeunga, J. Bopda, et Joseph Kamgno. 2014a. « Epidemiology of Loa Loa and
1155 Mansonella Perstans Filariasis in the Akonolinga Health District, Centre Region,
1156 Cameroon. » *HEALTH SCIENCES AND DISEASE* 15 (1). [https://www.hsd-](https://www.hsd-fmsb.org/index.php/hsd/article/view/297)
1157 [fmsb.org/index.php/hsd/article/view/297](https://www.hsd-fmsb.org/index.php/hsd/article/view/297).

1158 ———. 2014b. « Epidemiology of Loa Loa and Mansonella Perstans Filariasis in the
1159 Akonolinga Health District, Centre Region, Cameroon. » *HEALTH SCIENCES AND*
1160 *DISEASE* 15 (1). <https://doi.org/10.5281/hsd.v15i1.297>.

1161 Tavares da Silva, Lucyane Bastos, James Lee Crainey, Túllio Romão Ribeiro da Silva, Uziel
1162 Ferreira Suwa, Ana Carolina Paulo Vicente, Jansen Fernandes de Medeiros, Felipe
1163 Arley Costa Pessoa, et Sérgio Luiz Bessa Luz. 2017. « Molecular Verification of New
1164 World Mansonella perstans Parasitemias ». *Emerging Infectious Diseases* 23 (3):
1165 545-47. <https://doi.org/10.3201/eid2303.161159>.

1166 Taylor, Mark J, Achim Hoerauf, et Moses Bockarie. 2010. « Lymphatic Filariasis and
1167 Onchocerciasis ». *The Lancet* 376 (9747): 1175-85. [https://doi.org/10.1016/S0140-](https://doi.org/10.1016/S0140-6736(10)60586-7)
1168 [6736\(10\)60586-7](https://doi.org/10.1016/S0140-6736(10)60586-7).

1169 Thylefors, B. 2004. « Eliminating Onchocerciasis as a Public Health Problem ». *Tropical*
1170 *Medicine & International Health* 9 (4). [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-3156.2004.01226.x)
1171 [3156.2004.01226.x](https://doi.org/10.1111/j.1365-3156.2004.01226.x).

1172 Unnasch, Thomas R, Allison Golden, Vitaliano Cama, et Paul T Cantey. 2018. « Diagnostics
1173 for Onchocerciasis in the Era of Elimination ». *International Health* 10 (suppl_1):
1174 i20-26. <https://doi.org/10.1093/inthealth/ihx047>.

1175 Varenne, F., J. Fillaux, M. Porterie, J. Soler, M. Cassagne, et V. Soler. 2016. « Loase sous-
1176 conjonctivale : à propos d'un cas ». *Journal Français d'Ophtalmologie* 39 (8): e193-94.
1177 <https://doi.org/10.1016/j.jfo.2015.08.018>.

1178 Vinkes Melchers, Natalie V. S., Sarah Mollenkopf, Robert Colebunders, Michael Edlinger,
1179 Luc E. Coffeng, Julia Irani, Trésor Zola, et al. 2018. « Burden of Onchocerciasis-

- Associated Epilepsy: First Estimates and Research Priorities ». *Infectious Diseases of Poverty* 7 (1). <https://doi.org/10.1186/s40249-018-0481-9>.
- Vinkeles Melchers, Natalie V. S., Wilma A. Stolk, Welmoed Van Loon, Belén Pedrique, Roel Bakker, Michele E. Murdoch, Sake J. De Vlas, et Luc E. Coffeng. 2021. « The Burden of Skin Disease and Eye Disease Due to Onchocerciasis in Countries Formerly under the African Programme for Onchocerciasis Control Mandate for 1990, 2020, and 2030 ». Édité par Alberto Novaes Ramos. *PLOS Neglected Tropical Diseases* 15 (7): e0009604. <https://doi.org/10.1371/journal.pntd.0009604>.
- Vlaminck, Johnny, Peter U. Fischer, et Gary J. Weil. 2015. « Diagnostic Tools for Onchocerciasis Elimination Programs ». *Trends in Parasitology* 31 (11): 571-82. <https://doi.org/10.1016/j.pt.2015.06.007>.
- Volpicelli, Lorenzo, Maurizio De Angelis, Alessandra Morano, Elisa Biliotti, Cristiana Franchi, Simona Gabrielli, Simonetta Mattiucci, Carlo Di Bonaventura, et Gloria Taliani. 2020. « Encephalopathy in a Patient with Loiasis Treated with Albendazole: A Case Report ». *Parasitology International* 75 (avril):102006. <https://doi.org/10.1016/j.parint.2019.102006>.
- Walther, Michael, et Ralph Muller. 2003. « Diagnosis of Human Filariases (except Onchocerciasis) ». In *Advances in Parasitology*, 53:149-93. Elsevier. [https://doi.org/10.1016/S0065-308X\(03\)53004-4](https://doi.org/10.1016/S0065-308X(03)53004-4).
- Wanji, S., N. Tendongfor, M. Esum, S. Ndindeng, et P. Enyong. 2003. « Epidemiology of Concomitant Infections Due to Loa Loa, Mansonella Perstans, and Onchocerca Volvulus in Rain Forest Villages of Cameroon ». *Medical Microbiology and Immunology* 192 (1): 15-21. <https://doi.org/10.1007/s00430-002-0154-x>.
- Whittaker, Charles, Martin Walker, Sébastien D.S. Pion, Cédric B. Chesnais, Michel Boussinesq, et María-Gloria Basáñez. 2018. « The Population Biology and Transmission Dynamics of Loa Loa ». *Trends in Parasitology* 34 (4): 335-50. <https://doi.org/10.1016/j.pt.2017.12.003>.
- World Health Organization (WHO). 2020. « Onchocerciasis factsheet. Geneva: World Health Organization », 16. <https://www.who.int/news-room/factsheets/detail/onchocerciasis>.
- World Health Organization. 2011. « Agir pour réduire l'impact mondial des maladies tropicales négligées : premier rapport de l'OMS sur les maladies tropicales négligées ». *Working to overcome the global impact of neglected tropical diseases: first WHO report on neglected tropical diseases*. <https://iris.who.int/handle/10665/44694>.

- . 2022. « Global programme to eliminate lymphatic filariasis: progress report, 2021 », 2022. <http://www.who.int/wer>.
- World Health Organization. *Bench Aids for the Diagnosis of Filarial Infection*. Geneves: 1997. <https://iris.who.int/handle/10665/37156> Consulté Le 2024-07-11. s. d.
- World Health Organization (WHO). 2021. « Global programme to eliminate lymphatic filariasis: progress report, 2020 ». *Weekly Epidemiological Record = Relevé épidémiologique hebdomadaire* 96 (41): 497-508. CORREA.
- World Health Organization (WHO). 2022. « Plan stratégique national de lutte contre les maladies tropicales négligées 2022 - 2025 Sénégal », 144. <https://espen.afro.who.int/system/files/content/resources/Senegal%20Plan%20National%20stratégique%20MTN%202022-2025.pdf>.
- World Health Organization (WHO). 2023. « Lymphatic Filariasis ». 1 juin 2023. <https://www.who.int/fr/news-room/fact-sheets/detail/lymphatic-filariasis>.
- Wynd, Shona. 2007. « Understanding the community impact of lymphatic filariasis: a review of the sociocultural literature ». *Bulletin of the World Health Organization* 85 (6): 493-98. <https://doi.org/10.2471/BLT.06.031047>.
- Yokoly, Firmain N., Julien B. Z. Zahouli, Aboulaye Méite, Millicent Opoku, Bernard L. Kouassi, Dziedzom K. De Souza, Moses Bockarie, et Benjamin G. Koudou. 2020. « Low Transmission of *Wuchereria Bancrofti* in Cross-Border Districts of Côte d'Ivoire: A Great Step towards Lymphatic Filariasis Elimination in West Africa ». Édité par Justin V. Remais. *PLOS ONE* 15 (4): e0231541. <https://doi.org/10.1371/journal.pone.0231541>.
- Zouré, Honorat Gustave Marie, Samuel Wanji, Mounkaïla Noma, Uche Veronica Amazigo, Peter J. Diggle, Afework Hailemariam Tekle, et Jan H. F. Remme. 2011. « The Geographic Distribution of *Loa Loa* in Africa: Results of Large-Scale Implementation of the Rapid Assessment Procedure for Loiasis (RAPLOA) ». Édité par Giovanna Raso. *PLoS Neglected Tropical Diseases* 5 (6): e1210. <https://doi.org/10.1371/journal.pntd.0001210>.

ANNEXES

Publication en collaboration

ARTICLE 3

Epidemiology of *Mansonella perstans* in Mali

(Soumis dans **Parasitology & Vector-Borne Diseases**)

Pascal Dembélé, **Manthia Diallo**, Papa Mouhamadou Gaye, Mady Cissoko, Mahamadou A
Thera and Stéphane Ranque

Epidemiology of *Mansonella perstans* in Mali

Pascal Dembélé^{1, 2, 3}, Manthia Diallo^{1, 4}, Papa Mouhamadou Gaye^{1, 2}, Mady Cissoko^{3, 5},
Mahamadou A Thera⁵ and Stéphane Ranque^{1, 2} *

¹Institut Hospitalo-Universitaire (IHU) Méditerranée Infection, 13005 Marseille, France

²Aix-Marseille University, AP-HM, SSA, RITMES, 13005 Marseille, France

³Programme National de Lutte Contre le Paludisme (PNLP), Bamako BP 233, Mali

⁴Aix-Marseille University, AP-HM, SSA, MEPHI, 13005 Marseille, France

⁵Malaria Research and Training Center (MRTC), FMOS-FAPH, Mali-NIAID-ICER,
Université des Sciences, des Techniques et des Technologies de Bamako, BP 423, Mali

*Correspondence: stephane.ranque@univ-amu.fr

Abstract

Introduction: Mansonellosis remains one of the most neglected tropical diseases caused by filarial nematodes of the genus *Mansonella*, whose different species are: *Mansonella perstans*, *Mansonella ozzardi* and *Mansonella streptocerca*. *Mansonella perstans* is the most common species in Mali. The aim of our study was to use real-time PCR (RT-PCR) to detect *Mansonella perstans* parasites in febrile patients in Mali, using malaria rapid diagnostic tests (RDTs).

Methods: From June to December 2021, negative and positive RDTs were collected from 41 sites in the nine regions of Mali. Molecular tools were used to detect and identify the presence of pan-filaria at the base of 28S in DNA samples extracted from the RDT cassette tape. Positive samples were tested with a (RT-PCR) specific for *Mansonella perstans*. Standard polymerase chain reaction (PCR) molecular analysis of targeting the DNA internal transcribed spacer 1 (ITS1) gene and phylogenetic analysis were performed.

Results: *Mansonella Perstans* DNA was detected in 1.33% of malaria-negative RDTs (2 out of 150 randomly selected), collected in the region of Koulikoro.

Conclusion: Result show the presence of *M. perstans*, in febrile patients using negative malaria RDTs although this infection due to a vector-borne filarial nematode is sufficiently neglected in Mali, and the need to further strengthen surveillance of mansonellosis caused by *M. perstans*.

Keywords: *Mansonella Perstan*; RDT; Mansonellosis; qPCR; Mali

Introduction

Mansonella perstans infection is a little-known parasitic disease caused by a filarial nematode transmitted to humans by the bites of tiny blood-sucking flies of the *Culicoides* genus. It commonly known as biting midges. These insects are tiny flies that feed on the blood of their hosts, enabling the transmission of filarial parasites. *Mansonella perstans* is widespread in several regions of sub-Saharan Africa, where it poses an often-underestimated public health issue. It is also found in some parts of Central and South America, although its geographical distribution there is more limited (Simonsen, Onapa, et Asio 2011b).

Three main species are responsible for mansonellosis in humans: *M. perstans*, *M. streptocerca* and *M. ozzardi* (E. R. Bregani, Tantardini, et Rovellini 2007; B. L. Downes et Jacobsen 2010). *M. perstans* and *M. streptocerca* are mainly found in west, east and central Africa, where they often coexist with other parasitic infections. These regions are characterized by environmental conditions that favor the proliferation of biting midges, thereby increasing the risk of transmission. when, *M. ozzardi* is mainly found in Latin America, with an uneven geographical distribution. It is endemic in certain regions Central and South America, as on several Caribbean islands [3]. In addition to *Mansonella perstans*, a potential new species of *Mansonella* sp "TWO" was recently identified in febrile children in Gabon (Rodi et al. 2023b). A large part of sub-Saharan Africa is affected by *M. perstans*, it extends from Senegal to Angola, through central Africa, north to southwest Sudan, east to Uganda, Kenya, Tanzania, and south to Zimbabwe (Agbolade et al. 2006). In the most affected countries, the species *M. perstans* mainly prevails in poor rural populations (Agbolade et al. 2006; Wanji et al. 2003). It is important to note that more than 100 million people are currently infected in Africa alone and approximately 600 million people live in 33 countries at high risk of *M. perstans* infection (Simonsen, Onapa, et Asio 2011b).

The essential data relating to mansonellosis and geographical distribution are based on the morphological characterization of microfilariae collected from blood samples by thick film or blood smear (Simonsen, Onapa, et Asio 2011b). Adult worms appear to live in various cavities of the body (pleural, peritoneal or pericardial, etc.) and the mesentery (El Setouhy et al. 2004), while the microfilariae produced by females circulate in the peripheral blood or the skin without any periodicity (*World Health Organization. Bench Aids for the Diagnosis of Filarial Infection. Geneves: 1997.*<https://iris.who.int/handle/10665/37156> Consulté Le 2024-07-11, s. d.). How long mature *M. perstans* worms survive in the human body remains unclear,

while their larval form, microfilariae, persist for approximately 4 months (Santa Maria Asio, Simonsen, et Onapa 2009a).

There are similarities in the life cycle of the three species of *Mansonella* parasites responsible for mansonellosis in humans and they vary in terms of biological, clinical and epidemiological characteristics (Simonsen, Onapa, et Asio 2011b; B. L. Downes et Jacobsen 2010; Mediannikov et Ranque 2018b). S1 Table summarizes the biological characteristics of these three filariasis (Simonsen, Onapa, et Asio 2011b; Mediannikov et Ranque 2018b).

Currently, it is recognized that *M. perstans* has mild pathogenic effects and mainly manifests asymptotically. Eosinophilia is a common feature (Mediannikov et Ranque 2018b). However, in *M. perstans* certain clinical manifestations could be associated with the migration of worms and in particular allergic reactions such as “Calabar edema”, pruritus, fever, headache, pericarditis and pleurisy (Simonsen, Onapa, et Asio 2011b). No duration between infection and the manifestation of microfilariae in peripheral blood has yet been proven [5]. Detection of microfilariae in blood smears helps confirm the diagnosis of *M. perstans* (Adolph, Kagan, et McQuay 1962; El Setouhy et al. 2004). One of the difficulties in the management of mansonellosis is the absence of a known standard treatment [5].

In a study carried out in a rural area of Senegal, the overall prevalence of *M. perstans* was 14.5% in febrile and non-febrile patients. It increases with age moreover this figure could reach up to 40% to 50% in rural areas (Bassene et al. 2015b; Santa Maria Asio, Simonsen, et Onapa 2009b). *Mansonella perstans* is the species involved in Mali, well known since 1912 (Keita 1979).

Several studies on mansonellosis caused by *M. perstans* in Mali attract our attention in particular: Thiroux had found a prevalence rate of 46% of *M. perstans* among soldiers recruited in French Sudan (present-day Mali) and that of 12.4% of *M. perstans* in the capital Bamako (Keita 1979). Similarly, Rougement in 1979 found in Masantola in the circle of Kolokani a high prevalence of (65%) then in Kadiolo in February 1978, (central district), a team from the epidemiology laboratory of parasitic infections of the faculty of medicine and dentistry stomatology (FMOS) estimated that the prevalence of *M. perstans* among schoolchildren aged 6 to 17 was 6.7% (Keita 1979). In the village of Lofigué in Kadiolo *M. perstans* was 28.45%. In the Kolokani circle (Central District) in April 1978, out of 565 daytime thick blood films carried out among the populations of 10 villages, the prevalence of *M. perstans* was between 5.5% and 78% (Yaya I. Coulibaly et al. 2009b; Keiser et al. 2003; 2008). Since then, no recent information has been available on this disease in Mali. Cases of non-malarial fever may be

attributed to other parasitoses. It is necessary to estimate the prevalence in different regions and transmission zones of Masantola.

The objective of this study was to assess the prevalence of this neglected parasite in febrile patients using nucleic acids (DNA) extracted from positive and negative malaria rapid diagnostic tests collected in different regions of Mali.

Materials and Methods

Study sites

Mali is a landlocked country in West Africa located between the 10- and 25-degree north latitudes and between the 4- and 12-degree west longitudes. It covers a 1,241,238 km² area. There are 3 climatic zones in Mali that extend from south to north: the Sudano-Guinean zone, which covers 25% of the territory and has a rainfall of approximately 1300 to 1500 mm per year; the Sahelian zone, which covers 50% of the territory and receives rainfall of 200 to 800 mm per year; and the Saharan desert zone, which represents 25% of the territory. This zone is marked by irregular rainfall, often less than 200 mm per year. For this study, we collected RDT cassettes from 41 sites in the following regions of Mali: Kayes, Koulikoro, Sikasso, Ségou, Mopti, Timbuktu, Gao, Kidal and Menaka, from June to December 2021. No sample was collected in the Taoudeni region or the capital city Bamako.

Ethical Considerations

Authorization to use malaria RDTs cassettes for research purposes was obtained from the competent authority of the National Malaria Control Program (NMCP) under number 064/MSDS-SG/NMCP, February 8, 2023), in accordance with Malian regulations on ethics and medical research.

Molecular analysis

DNA extraction from RDTs cassettes

A total of 1496 RDTs cassettes was used for genomic DNA extraction as a substrate to detect the filaria parasites including *M. perstans*. The RDTs cassettes were previously used to detect *Plasmodium falciparum* (*P.f*) (Dembélé et al. 2024). DNA was extracted using the EZ1 DNA Blood Kit (Qiagen) on an EZ1 BioRobot extraction device (Qiagen, Hilden, Germany) according the manufacturer's instructions and as described by Dembélé et al. (Dembélé et al. 2024).

Real-time PCR (RT-PCR) for detection of *M. perstans*

Real-time PCR was used to detect all filaria species and *M. perstans* respectively, in all DNA extracts from RDTs cassettes (n= 1496). The RT-PCR reaction targets the 28S region for all filaria species and another specific region to detect *M. perstans*. The primers used pan-filaria 28S (qFil_F and qFil_R and probe) (Table 1), targeting all filaria species were identical to those initially described by Laidoudi et al. (Laidoudi et al. 2020). All Pan-filaria 28S positive samples were then tested with a specific *M. perstans* system using the primers and probes previously described by Bassene et al (Bassene et al. 2015b) (**Table 1**).

The RT-PCR reactions were performed using a CFX96 thermal cycler (Bio-Rad, Marnes-la-Coquette, France) in a 20 µL reaction mixture containing 5 µL of DNA, 10 µL of the Master Mix, 3.5 µL of sterile distilled water, and 0.5 µL of each of the primers and 0.5 µL of the TaqMan probe (Applied Biosystems, Foster City, CA, USA). Our amplification program consisted of an initial 2-min denaturation step at 50 °C followed by a 3-min denaturation at 95 °C, then 40 cycles of 95 °C for 30 s and 60 °C for 1 min before holding the sample at 4 °C. For each RT-PCR run, negative (sterile distilled water) and positive controls (*Dirofilaria immitis* DNA was used for pan-filarial RT-PCR and a plasmid developed in our laboratory was used for *M. perstans* RT-PCR) were used.

Table 1. Primers used in the real-time PCR, Pan-filaria and *Mansonella perstans* parasite and *Mansonella* spp. sequencing protocols.

Primer names	Marker	Target organism	Primer sequences (5'-3')	Authors
Real-time PCR				
qFil_F	28S	Pan-filariae	TTGTTTGAGATTGCAGCCCA	Laidoudi et al (Laidoudi et al. 2020)Laidoudi et al (Laidoudi et al. 2020)
qFil_R			GTTTCCATCTCAGCGGTTTC	
Probe (FAM 5'–3'TAMRA)			CAAGTACCGTGAGGGAAAGT	
Mp_Hub_F	ITS	<i>M. perstans</i>	AGGATCATTAACGAGCTTCC	Bassene et al (Bassene et al. 2015b)Bassene et al (Bassene et al. 2015b)
Mp_Hub_R			CGA ATATCACCGTTAATTCAGT	
Probe (FAM 5'–3'TAMRA)			TTCACTTTTATTTAGCAACATGCA	
Standard PCR and sequencing				
ITS1_F	ITS1	<i>Mansonella</i> spp.	GGTGAACCTGCGGAAGGATC	Mourembou et al (Mourembou et al. 2015b)Mourembou et al (Mourembou et al. 2015b)
ITS1_R			CTCAATGCGTCTGCAATTCGC	

DNA-based sequencing

Only RT-PCR positive samples for both genes were subjected to PCR sequencing using Internal transcribed spacer 1 (ITS 1) region to identify *Mansonella* parasites. PCR reaction targeting a 240 bp region of ITS1 was performed in a thermal cycler (Applied Biosystems, 2720, Foster City, United States) with AmpliTaq Gold 360 PCR master mix (Applied Biosystems, Waltham, United States). The ITS region was amplified using primers forward ITS1_F and reverse ITS1_R Mourembou et al (Mourembou et al. 2015b) (**Table 1**). PCR was performed in a reaction mixture consisting of 5 µL of extracted DNA, 12.5 µL of AmpliTaq Gold 360 PCR master mix (Applied Biosystems, Waltham, MA, USA), 0.75 µL of each primer, and 6 µL of sterile distilled water, in a final reaction volume of 25 µL. The amplification program consisted in an initial denaturation at 95°C for 15 min, followed by 40 cycles at 95°C for 30 s, at 58°C for 30 s, 72°C for 1 min and a final step at 72°C for 5 min.

The amplified products were purified using a Macherey Nagel plate (NucleoFast 96 PCR, Düren, Germany). Sanger sequencing was performed using the same primers as for PCR with BigDye Terminator v1.1, v3.15× sequencing buffer (Applied Biosystems, Warrington, UK) and run on an ABI 3100 automated sequencer (Applied Biosystems). The sequences obtained were assembled with ChromasPro version 1.7.7 software (Technelyium Pty. Ltd, Tewantin, Australia) and aligned with BioEdit software (<http://www.mbio.ncsu.edu/BioEdit/bioedit.html>) using the ClustalW algorithm to perform multiple alignments of our sequences. The corrected sequences were compared to existing sequences in the online database of the National Centre for Biotechnology Information (NCBI) (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) using basic local alignment search tool (BLAST) analysis (Chenna et al. 2003). Phylogenetic tree (**Figure 1**) was constructed using the maximum likelihood (ML) statistical method with substitution model selection determined by the Molecular Evolutionary Genetics Analysis (MEGA) v7.0.26 software (Tamura K, 1993). Statistical support for the inner branches of the trees was assessed by bootstrapping with 1,000 iterations.

Results

Molecular characterization of filarial species and *M. perstans*

A total of 1,496 malaria RDTs cassettes were collected in Mali, of which 260 were *P. falciparum* positive and 1,236 negatives in febrile patients. All RDT cassettes were tested with the Pan-filaria 28S and *M. perstans* systems specifically. Of these samples, only 2/1496 (0.1%)

were positive for both Pan-filaria 28S RT-PCR and *M. perstans* (S1 fig). *M. perstans* species were only detected on TDRs collected in the Koulikoro region.

S1 Figure. Amplification curves for real-time PCR on CFX96 targeting the qFil 28S gene of *M. perstans* species in Koulikoro (T+: positive control and the two positive samples detected in the Kangaba health district (Koulikoro region).

DNA sequence-based identification of *Mansonella* parasites

Sequencing identification was carried out on the RT-PCR positive samples of *M. perstans* (n= 2) using the ITS1 region. The sequences obtained were subjected to BLAST analysis against reference sequences from the GenBank database (GB). BLAST analysis indicated that our *M. perstans* sequences matched 100.00% of their homologous sequences (GB accession number: PQ041650 and MN821068) described in Senegal and Gabon respectively and available in GenBank.

Our sequences have been deposited in the GB database under accession numbers XX and XX.

Phylogenetic Tree based ITS was constructed using our *Mansonella perstans* sequences in addition to other sequences downloaded from GenBank. The tree revealed that our sequences are close to those already found in Senegal, Brazil and Gabon and that they form the same clade. The phylogenetic tree also showed interspecific discrimination (**Figure 1**).




Figure 1. The tree is based on the nucleotide sequence of the ITS region, constructed using the maximum likelihood method with MEGA 7. Branch values are bootstrap support values based on 1000 replicates.

Discussion

Studies aimed at detecting and identifying filarial parasites including *M. perstans* from DNA extracted from malaria RDT cassettes collected in 9 regions of Mali.

Our findings demonstrate for the first time the presence of *M. perstans* Mansonellosis in Mali by using innovative approaches to detect the presence of the parasite from *P.f.* malaria RDT cassettes in febrile patients attending health facilities in the Koulikoro region.

Several studies on mansonellosis caused by *M. perstans* in Mali have been carried out in the south and north of the country in particular, surveys carried out by the FMOS Laboratory of Epidemiology of Parasitic Affections in the cercle of Kadiolo (Central Arrondissement, in February 1978) showed that, out of 782 thick diurnal drops in schoolchildren aged 6 to 17, the prevalence of *M. perstans* was around 6.7%, including 3.9% in children aged 7 to 10 and 10.9% in children aged 11 to 17 (Keita 1979) *perstans* was around 6.7%, including 3.9% in children aged 7 to 10 and 10.9% in children aged 11 to 17. Also, out of 123 thick nocturnal drops taken from adults in the village of Lofigué, the prevalence of *M. perstans* was 28.45%, with *M. perstans* + *W. bancrofti* associations accounting for 8.13% (Keita 1979).

In the cercle de Kolokani (Arrondissement Central, April 1978), *M. perstans* prevalence ranged from 5.5% to 78% in 565 thick diurnal drops from 10 villages (Keita 1979). In Gourma (April - May 1979), out of 608 thick diurnal drops made among nomadic populations north of parallel 15°30'N on either side of the 300 mm isohyet, none showed *M. perstans* microfilaria (Keita 1979). These results were obtained by microscopy. Then into 2003, Paul B Keiser et al reported a prevalence of 75% *M. perstans* in Mali (Keiser et al. 2003).

Although *M. perstans* infection is one of the well-known human filariasis in Mali, it must be noted that very few studies are still carried out on its epidemiology and its negative effects on the health of endemic populations.

We identified by real-time PCR, *M. perstans* in two DNA samples extracted from negative and positive *plasmodium falciparum* malaria (RDTs) in febrile patients, which represents a prevalence of 0.1% (2/1496). The low prevalence obtained in our results could probably be explained by the positive effects of the molecules used (albendazole and ivermectin) during mass drug distribution campaigns against neglected tropical diseases, a strategy considered effective and which is recommended by the World Health Organization (WHO) to reduce the burden of neglected tropical diseases in the community, interrupt

transmission towards their elimination and control by 2030 (« OMS MALI Rapport annuel 2018.pdf », s. d.). For example, during the mass treatment campaign against neglected tropical diseases with preventive chemotherapy in 2018, that is to say 3 years before the collection of the malaria RDT cassettes, the therapeutic coverage (Ivermectin + Albendazole) and Ivermectin in a single dose were 86.34% and 96.84% respectively (« OMS MALI Rapport annuel 2018.pdf », s. d.). Ivermectin distributed during mass campaigns against lymphatic filariasis and onchocerciasis may affect prevalence of *Mansonella perstans* species (Ta-Tang et al. 2021a).

Contrary to our results, in Senegal the overall prevalence of *M. perstans* in a rural area of Senegal was 14.5% (29/201) in febrile and non-febrile patients (Bassene et al. 2015b). In a previous study on the comparison of *M. perstans* detection techniques, the authors showed that qPCR could be successfully used to identify microfilariae in the samples used (Bassene et al. 2015b). It was demonstrated in a study conducted in two endemic communities in Uganda that the prevalence of *M. perstans* filariasis increased rapidly with age, reaching 89.2% and 81.4% in the age group of 15 at 19 years old. The overall prevalence of microfilariae (mf) was significantly higher in Mukono (76.5%) than in Luwero (57.7%). From the age of 1 to 4 years, 40.6% and 20.5% of children were positive for microfilariae (Santa Maria Asio, Simonsen, et Onapa 2009b).

Several studies of co-infection have sometimes been reported in some endemic countries. Consequently, it is appropriate to put in place adequate diagnostic tools whose sensitivity and specificity are indisputable in order to remove all ambiguities in the treatment of neglected filariasis. The study by Tatuene et al in the Akonolinga health district located in central Cameroon corroborates that the prevalence of *M. perstans* infestation was relatively high 12.4% and that of Loa co-infection. loa – *M. perstans* was 5.1% (Tatuene et al. 2014b). Furthermore, a similar study carried out in 16 isolated villages in southern Cameroon revealed that onchocerciasis and mansonellosis were highly endemic in all villages surveyed with prevalence ranging from 28.44% to 87.17% for *O. volvulus* and from 52.48% to 100% for *M. perstans* [9].

The fight and eradication of filarial infections in humans requires the implementation of prevention strategies, appropriate treatment and reinforced surveillance of the disease.

Limitations

In view of the interesting results obtained in this filarial endemic region of Mali, it is important to note that the conservation of RDT cassettes, and the lack of knowledge of patient demographics and clinical symptoms could be potential limitations of our approach.

Conclusion

For the first time, the use the malaria RDT cassettes enabled molecular screening of filarial species including *M. perstans* and revealed a low infestation rate in Mali located in the south. Further molecular and genetic studies of this little-known human filariasis are needed to properly assess its importance in terms of public health, and to decipher the molecular interactions between the different species of filarial parasites with a view to strengthening mansonellosis control strategies in Mali.

Author contributions: P.D., MD., M.A.T. and S.R. designed the study and wrote the first version of the manuscript.

P.D., M.D., P.M.G., and M.C., analyzed and interpreted the results.

M.D., P.M.G., M.A.T. and S.R. Contributed to the critical revision of the manuscript. All authors read and approved the final version.

Acknowledgment


This study was made possible thanks to the academic and financial support of the Mediterranean Infection Foundation in Marseille, France, and the support of the National Malaria Control Program. We thank each of you

Conflicts of interest:

The authors declare no conflict of interest.

325 **S1 Table.** Main characteristics of human mansonellosis.

Characteristic	<i>Mansonella streptocerca</i>	<i>Mansonella ozzardi</i>	<i>Mansonella perstans</i>
Endemic areas	West and Central Africa	Central and South America, Caribbean	Sub-Saharan and Northern Africa, South America
Arthropod vectors	<i>Culicoïdes</i> spp.	<i>Culicoïdes</i> spp; Simulium amazonicum	<i>Culicoïdes</i> spp.
Location of adult worms	Dermis of the trunk and upper shoulder girdle	Lymphatic vessels, pericardium, pleura and peritoneum	Pericardium, pleura and peritoneum; mesentery and retroperitoneum
Location of microfilariae	Skin	Blood and skin	Blood



326

327

328

329

Références

- Adam, Nouhou Diori, Yacoubou Soumana, Ali Saley, et Idrissa Saley. 2020. « Une Loase Oculaire: À Propos D'un Cas ». *Pan African Medical Journal* 36. <https://doi.org/10.11604/pamj.2020.36.302.25124>.
- Adewole, S.O., et S.K. Ayeni. 2009. « Clinical Manifestation of Onchocerciasis in Ise - Orun Local Government, Ekiti State, Nigeria ». *Pakistan Journal of Nutrition* 8 (2): 122-24. <https://doi.org/10.3923/pjn.2009.122.124>.
- Adhikari, Ram Kumar, Jeevan Bahadur Sherchand, Shiva Raj Mishra, Kamal Ranabhat, Amrit Pokharel, Pramila Devkota, Durga Mishra, et al. 2015. « Health-Seeking Behaviors and Self-Care Practices of People with Filarial Lymphoedema in Nepal: A Qualitative Study ». *Journal of Tropical Medicine* 2015:1-6. <https://doi.org/10.1155/2015/260359>.
- Adolph, Paul E., Irving G. Kagan, et Russell M. McQuay. 1962. « Diagnosis and Treatment of Acanthocheilonema Perstans Filariasis », janvier. <https://doi.org/10.4269/ajtmh.1962.11.76>.
- Agbolade, Olufemi Moses, Dora Olufunmilola Akinboye, Taiwo Monroof Olateju, Oluwatoyin Adepeju Ayanbiyi, Olukayode Olakunle Kuloyo, et Oluwalolami Oluwaseun Fenuga. 2006. « Biting of Anthropophilic Culicoides Fulvithorax (Diptera: Ceratopogonidae), a Vector of Mansonella Perstans in Nigeria ». *The Korean Journal of Parasitology* 44 (1): 67-72. <https://doi.org/10.3347/kjp.2006.44.1.67>.
- Akue, Jean Paul, Dieudonné Nkoghe, Cindy Padilla, Ghislain Moussavou, Hubert Moukana, Roger Antoine Mbou, Benjamin Ollomo, et Eric Maurice Leroy. 2011. « Epidemiology of Concomitant Infection Due to Loa Loa and Mansonella Perstans in Gabon ». Édité par Rachel Lisa Pullan. *PLoS Neglected Tropical Diseases* 5 (10): e1329. <https://doi.org/10.1371/journal.pntd.0001329>.
- Alhassan, Andy, Zhiru Li, Catherine B. Poole, et Clotilde K.S. Carlow. 2015. « Expanding the MDx Toolbox for Filarial Diagnosis and Surveillance ». *Trends in Parasitology* 31 (8): 391-400. <https://doi.org/10.1016/j.pt.2015.04.006>.
- Allen, Judith E., Ohene Adjei, Odile Bain, Achim Hoerauf, Wolfgang H. Hoffmann, Benjamin L. Makepeace, Hartwig Schulz-Key, et al. 2008. « Of Mice, Cattle, and Humans: The Immunology and Treatment of River Blindness ». Édité par Sara Lustigman. *PLoS Neglected Tropical Diseases* 2 (4): e217. <https://doi.org/10.1371/journal.pntd.0000217>.
- Alsarraf, Mustafa, Viktoria Levytska, Ewa J. Mierzejewska, Vasyl Poliukhovych, Anna Rodo, Mohammed Alsarraf, Dziyana Kavalevich, Dorota Dwużnik-Szarek, Jerzy M. Behnke,

- et Anna Bajer. 2021. « Emerging Risk of *Dirofilaria* Spp. Infection in Northeastern Europe: High Prevalence of *Dirofilaria Repens* in Sled Dog Kennels from the Baltic Countries ». *Scientific Reports* 11 (1): 1068. <https://doi.org/10.1038/s41598-020-80208-1>.
- Amazigo, U. 2008. « The African Programme for Onchocerciasis Control (APOC) ». *Annals of Tropical Medicine & Parasitology* 102 (sup1): 19-22. <https://doi.org/10.1179/136485908X337436>.
- Anderson, Roy C. 2000. « R.C. Anderson: Nematode Parasites of Vertebrates. Their Development and Transmission. 2nd Edition. CABI Publishing, Wallingford, Oxon (UK) 2000. 650 Pp., 43 Figs. Price £ 99.50 . », 1.
- Anosike, Jude C, Ikechukwu N S Dozie, Celestine O E Onwuliri, Bertram E B Nwoke, et Viola A Onwuliri. 2005. « PREVALENCE OF MANSONELLA PERSTANS INFECTIONS AMONG THE NOMADIC FULANIS OF NORTHERN NIGERIA », p.35-39. <https://pubmed.ncbi.nlm.nih.gov/16028864/>.
- Antinori, Spinello, Carlo Parravicini, Laura Galimberti, Antonella Tosoni, Paolo Giunta, Massimo Galli, Mario Corbellino, et Anna Lisa Ridolfo. 2017. « Is Imported Onchocerciasis a Truly Rare Entity? Case Report and Review of the Literature ». *Travel Medicine and Infectious Disease* 16 (mars):11-17. <https://doi.org/10.1016/j.tmaid.2017.02.006>.
- Asio, S. M., P. E. Simonsen, et A. W. Onapa. 2009. « *Mansonella Perstans* : Safety and Efficacy of Ivermectin Alone, Albendazole Alone and the Two Drugs in Combination ». *Annals of Tropical Medicine & Parasitology* 103 (1): 31-37. <https://doi.org/10.1179/136485909X384929>.
- Asio, Santa Maria, Paul E. Simonsen, et Ambrose W. Onapa. 2009a. « Analysis of the 24-h Microfilarial Periodicity of *Mansonella Perstans* ». *Parasitology Research* 104 (4): 945-48. <https://doi.org/10.1007/s00436-008-1312-x>.
- . 2009b. « *Mansonella Perstans* Filariasis in Uganda: Patterns of Microfilaraemia and Clinical Manifestations in Two Endemic Communities ». *Transactions of the Royal Society of Tropical Medicine and Hygiene* 103 (3): 266-73. <https://doi.org/10.1016/j.trstmh.2008.08.007>.
- Attout, Tarik, Achim Hoerauf, Gaëlle Dénécé, Alexander Yaw Debrah, Yeboah Marfo-Debrekyei, Michel Boussinesq, Samuel Wanji, et al. 2009. « Lymphatic Vascularisation and Involvement of Lyve-1+ Macrophages in the Human *Onchocerca* Nodule ». *Édité*

396 par Derya Unutmaz. *PLoS ONE* 4 (12): e8234.
 397 <https://doi.org/10.1371/journal.pone.0008234>.

398 Aubry, Arthur, Charles Hervochon, Marie Chauveau, Marie Denis-Musquer, Fakhri Jeddi,
 399 Rose-Anne Lavergne, Patrice Le Pape, et Florent Morio. 2022. « Une expulsion bien
 400 singulière d'un ver adulte de *Loa loa* ». *Revue Francophone des Laboratoires* 2022
 401 (538): 72-75. [https://doi.org/10.1016/S1773-035X\(21\)00381-6](https://doi.org/10.1016/S1773-035X(21)00381-6).

402 Aubry, Professeur Pierre, et Docteur Bernard-Alex Gaüzère. 2023. « Filarioses lymphatiques
 403 Actualités 2023 », P8.

404 Babu, Subash, et Thomas B. Nutman. 2012. « Immunopathogenesis of Lymphatic Filarial
 405 Disease ». *Seminars in Immunopathology* 34 (6): 847-61.
 406 <https://doi.org/10.1007/s00281-012-0346-4>.

407 Badia-Rius, Xavier, Salissou Adamou, Mark J. Taylor, et Louise A. Kelly-Hope. 2023.
 408 « Morbidity Hotspot Surveillance: A Novel Approach to Detect Lymphatic Filariasis
 409 Transmission in Non-Endemic Areas of the Tillabéry Region of Niger ». *Parasite
 410 Epidemiology and Control* 21 (mai):e00300.
 411 <https://doi.org/10.1016/j.parepi.2023.e00300>.

412 Bassene, Hubert, Masse Sambou, Florence Fenollar, Siân Clarke, Sawdiatou Djiba, Gaël
 413 Mourembou, Alioune Badara L Y, Didier Raoult, et Oleg Mediannikov. 2015a. « High
 414 Prevalence of *Mansonella Perstans* Filariasis in Rural Senegal ». *The American Journal
 415 of Tropical Medicine and Hygiene* 93 (3): 601-6. <https://doi.org/10.4269/ajtmh.15-0051>.
 416 0051.

417 ———. 2015b. « High Prevalence of *Mansonella Perstans* Filariasis in Rural Senegal ». *The
 418 American Journal of Tropical Medicine and Hygiene* 93 (3): 601-6.
 419 <https://doi.org/10.4269/ajtmh.15-0051>.

420 Bhalla, Devender, Michel Dumas, et Pierre-Marie Preux. 2013. « Neurological Manifestations
 421 of Filarial Infections ». In *Handbook of Clinical Neurology*, 114:235-42. Elsevier.
 422 <https://doi.org/10.1016/B978-0-444-53490-3.00018-2>.

423 Boatin, B. 2008. « The Onchocerciasis Control Programme in West Africa (OCP) ». *Annals of
 424 Tropical Medicine & Parasitology* 102 (sup1): 13-17.
 425 <https://doi.org/10.1179/136485908X337427>.

426 Bockarie, Moses J, et Rinki M Deb. 2010. « Elimination of Lymphatic Filariasis: Do We Have
 427 the Drugs to Complete the Job? ». *Current Opinion in Infectious Diseases* 23 (6):
 428 617-20. <https://doi.org/10.1097/QCO.0b013e32833fdee5>.

- Bockarie, Moses J, Mark J Taylor, et John O Gyapong. 2009. « Current Practices in the Management of Lymphatic Filariasis ». *Expert Review of Anti-Infective Therapy* 7 (5): 595-605. <https://doi.org/10.1586/eri.09.36>.
- Bottieau, Emmanuel, Ralph Huits, Steven Van Den Broucke, Ula Maniewski, Steven Declercq, Isabel Brosius, Caroline Theunissen, et al. 2022. « Human Filariasis in Travelers and Migrants: A Retrospective 25-Year Analysis at the Institute of Tropical Medicine, Antwerp, Belgium ». *Clinical Infectious Diseases* 74 (11): 1972-78. <https://doi.org/10.1093/cid/ciab751>.
- Bouchery, T., E. Lefoulon, G. Karadjian, A. Nieguitsila, et C. Martin. 2013. « The Symbiotic Role of Wolbachia in Onchocercidae and Its Impact on Filariasis ». *Clinical Microbiology and Infection* 19 (2): 131-40. <https://doi.org/10.1111/1469-0691.12069>.
- Bourée, P. 2005. « Parasitoses urinaires ». *Annales d'Urologie* 39 (6): 232-46. <https://doi.org/10.1016/j.anuro.2005.09.005>.
- Boussinesq, M. 2003. « L'onchocercose humaine en Afrique ». In *Trente ans de lutte contre l'onchocercose en Afrique de l'Ouest. Traitements larvicides et protection de l'environnement : Thirty years of onchocerciasis control in West Africa. Blackfly larviciding and environmental protection*, édité par Laurent Yaméogo, Christian Levêque, et Jean-Marc Hougard, 389-400. Synthèses. Marseille: IRD Éditions. <https://doi.org/10.4000/books.irdeditions.28767>.
- . 2006. « Loiasis ». *Annals of Tropical Medicine & Parasitology* 100 (8): 715-31. <https://doi.org/10.1179/136485906X112194>.
- Boussinesq, Michel. 2012. « Loiasis: New Epidemiologic Insights and Proposed Treatment Strategy ». *Journal of Travel Medicine* 19 (3): 140-43. <https://doi.org/10.1111/j.1708-8305.2012.00605.x>.
- Brattig, Norbert W., Robert A. Cheke, et Rolf Garms. 2021. « Onchocerciasis (River Blindness) – More than a Century of Research and Control ». *Acta Tropica* 218 (juin):105677. <https://doi.org/10.1016/j.actatropica.2020.105677>.
- Bregani, E. R., F. Tantardini, et A. Rovellini. 2007. « [Mansonella perstans filariasis] ». *Parassitologia* 49 (1-2): 23-26.
- Bregani, Enrico Rino, Angelo Rovellini, Narasem Mbaïdoum, et Maria Giovanna Magnini. 2006. « Comparison of Different Anthelmintic Drug Regimens against Mansonella Perstans Filariasis ». *Transactions of the Royal Society of Tropical Medicine and Hygiene* 100 (5): 458-63. <https://doi.org/10.1016/j.trstmh.2005.07.009>.

- Brengues, J., R. Subra, J. Mouchet, et G. S. Nelson. 1968. « [Transmission of *Wuchereria bancrofti* Cobbold in West Africa. Preliminary study of a focus in the savanna of north Guinea] ». *Bulletin of the World Health Organization* 38 (4): 595-608.
- Brugman, Victor A., Marion E. England, Joanne Stoner, Laura Tugwell, Lara E. Harrup, Anthony J. Wilson, Jolyon M. Medlock, et al. 2017. « How Often Do Mosquitoes Bite Humans in Southern England? A Standardised Summer Trial at Four Sites Reveals Spatial, Temporal and Site-Related Variation in Biting Rates ». *Parasites & Vectors* 10 (1): 420. <https://doi.org/10.1186/s13071-017-2360-9>.
- Burbelo, Peter D., Roshan Ramanathan, Amy D. Klion, Michael J. Iadarola, et Thomas B. Nutman. 2008. « Rapid, Novel, Specific, High-Throughput Assay for Diagnosis of *Loa Loa* Infection ». *Journal of Clinical Microbiology* 46 (7): 2298-2304. <https://doi.org/10.1128/JCM.00490-08>.
- Büttner, Dietrich W., Samuel Wanji, Chiara Bazzocchi, Odile Bain, et Peter Fischer. 2003. « Obligatory Symbiotic *Wolbachia* Endobacteria Are Absent from *Loa Loa* ». *Filaria Journal* 2 (1): 10. <https://doi.org/10.1186/1475-2883-2-10>.
- CAMPILLO, Jérémy T., Frédéric LOUYA, Paul BIKITA, François MISSAMOU, Michel BOUSSINESQ, Sebastien D. S. PION, Sébastien BERTOUT, et Cédric B CHESNAIS. 2023. « Évaluation en milieu rural au Congo de la fiabilité de la méthode de la goutte épaisse calibrée pour évaluer le niveau de la microfilarémie à *Loa loa* ». *MTSI*, mars, Vol. 3 No 1 (2023): MTSIRevue. <https://doi.org/10.48327/MTSI.V3I1.2023.297>.
- Carne, B., et P. Esterre. 2012. « Filarioses ». *EMC - Maladies infectieuses* 9 (2): 1-19. [https://doi.org/10.1016/S1166-8598\(12\)41786-0](https://doi.org/10.1016/S1166-8598(12)41786-0).
- CDC. 2024. « Symptoms of Onchocerciasis ». *Filarial Worms*. 5 juin 2024. <https://www.cdc.gov/filarial-worms/signs-symptoms/onchocerciasis.html>.
- Chakraborty, Sanjukta, Manokaran Gurusamy, David C. Zawieja, et Mariappan Muthuchamy. 2013. « Lymphatic Filariasis: Perspectives on Lymphatic Remodeling and Contractile Dysfunction in Filarial Disease Pathogenesis ». *Microcirculation* 20 (5): 349-64. <https://doi.org/10.1111/micc.12031>.
- Chandy, Anish, Alok Singh Thakur, Mukesh Pratap Singh, et Ashish Manigauha. 2011. « A Review of Neglected Tropical Diseases: Filariasis ». *Asian Pacific Journal of Tropical Medicine* 4 (7): 581-86. [https://doi.org/10.1016/S1995-7645\(11\)60150-8](https://doi.org/10.1016/S1995-7645(11)60150-8).
- Chelsea, Marie, et A. Petri William. 2022. « Mansonellose-Maladies infectieuses ». <https://www.msdmanuals.com/fr/professional/maladies-infectieuses/n%C3%A9matodes-vers-ronds/mansonellose>.

- Chenna, Ramu, Hideaki Sugawara, Tadashi Koike, Rodrigo Lopez, Toby J. Gibson, Desmond G. Higgins, et Julie D. Thompson. 2003. « Multiple Sequence Alignment with the Clustal Series of Programs ». *Nucleic Acids Research* 31 (13): 3497-3500. <https://doi.org/10.1093/nar/gkg500>.
- Chesnais, Cédric B, Charlotte Bizet, Jérémy T Campillo, Wepnyu Y Njamnshi, Jean Bopda, Philippe Nwane, Sébastien D Pion, Alfred K Njamnshi, et Michel Boussinesq. 2020. « A Second Population-Based Cohort Study in Cameroon Confirms the Temporal Relationship Between Onchocerciasis and Epilepsy ». *Open Forum Infectious Diseases* 7 (6): ofaa206. <https://doi.org/10.1093/ofid/ofaa206>.
- Coffeng, Luc E., Wilma A. Stolk, Achim Hoerauf, Dik Habbema, Roel Bakker, Adrian D. Hopkins, et Sake J. De Vlas. 2014. « Elimination of African Onchocerciasis: Modeling the Impact of Increasing the Frequency of Ivermectin Mass Treatment ». Édité par James M. McCaw. *PLoS ONE* 9 (12): e115886. <https://doi.org/10.1371/journal.pone.0115886>.
- Colebunders, Robert, Jane Y Carter, Peter Claver Olore, Kai Puok, Samit Bhattacharyya, Sonia Menon, Gasim Abd-Elfarag, et al. 2018. « High Prevalence of Onchocerciasis-Associated Epilepsy in Villages in Maridi County, Republic of South Sudan: A Community-Based Survey ». *Seizure* 63 (décembre):93-101. <https://doi.org/10.1016/j.seizure.2018.11.004>.
- Coreil, Jeannine, et Gladys Mayard. 2009. « Indigenization of Illness Support Groups for Lymphatic Filariasis in Haiti ». In *Anthropology and Public Health*, édité par Robert A Hahn et Marcia Inborn, 2^e éd., 245-65. Oxford University Press New York. <https://doi.org/10.1093/acprof:oso/9780195374643.003.0010>.
- Coulibaly, Yaya I., Benoit Dembele, Abdallah A. Diallo, Ettie M. Lipner, Salif S. Doumbia, Siaka Y. Coulibaly, Siaka Konate, et al. 2009a. « A Randomized Trial of Doxycycline for *Mansonella Perstans* Infection ». *New England Journal of Medicine* 361 (15): 1448-58. <https://doi.org/10.1056/NEJMoa0900863>.
- . 2009b. « A Randomized Trial of Doxycycline for *Mansonella Perstans* Infection ». *The New England Journal of Medicine* 361 (15): 1448-58. <https://doi.org/10.1056/NEJMoa0900863>.
- Coulibaly, Yaya Ibrahim, Benoit Dembele, Abdallah Amadou Diallo, Sibylle Kristensen, Siaka Konate, Housseini Dolo, Ilo Dicko, et al. 2013. « *Wuchereria Bancrofti* Transmission Pattern in Southern Mali Prior to and Following the Institution of Mass Drug

- Administration ». *Parasites & Vectors* 6 (1): 247. <https://doi.org/10.1186/1756-3305-6-247>.
- Crawford, Katie E, Shannon M Hedtke, Stephen R Doyle, Annette C Kuesel, Samuel Armoo, Mike Y Osei-Atweneboana, et Warwick N Grant. 2024. « Genome-Based Tools for Onchocerciasis Elimination: Utility of the Mitochondrial Genome for Delineating *Onchocerca Volvulus* Transmission Zones ». *International Journal for Parasitology* 54 (3-4): 171-83. <https://doi.org/10.1016/j.ijpara.2023.11.002>.
- Cromwell, Elizabeth A, Chris A Schmidt, Kevin T Kwong, David M Pigott, Denise Mupfasoni, Gautam Biswas, Shreya Shirude, et al. 2020. « The Global Distribution of Lymphatic Filariasis, 2000–18: A Geospatial Analysis ». *The Lancet Global Health* 8 (9): e1186-94. [https://doi.org/10.1016/S2214-109X\(20\)30286-2](https://doi.org/10.1016/S2214-109X(20)30286-2).
- Davis, Emma L., Lisa J. Reimer, Lorenzo Pellis, et T. Deirdre Hollingsworth. 2019. « Evaluating the Evidence for Lymphatic Filariasis Elimination ». *Trends in Parasitology* 35 (11): 860-69. <https://doi.org/10.1016/j.pt.2019.08.003>.
- De Sole, G., J. Giese, F.M. Keita, et J. Remme. 1991. « Detailed Epidemiological Mapping of Three Onchocerciasis Foci in West Africa ». *Acta Tropica* 48 (3): 203-13. [https://doi.org/10.1016/0001-706X\(91\)90048-O](https://doi.org/10.1016/0001-706X(91)90048-O).
- Debrah, Alexander Yaw, Sabine Specht, Ute Klarmann-Schulz, Linda Batsa, Sabine Mand, Yeboah Marfo-Debrekyei, Rolf Fimmers, et al. 2015. « Doxycycline Leads to Sterility and Enhanced Killing of Female *Onchocerca Volvulus* Worms in an Area With Persistent Microfilaridermia After Repeated Ivermectin Treatment: A Randomized, Placebo-Controlled, Double-Blind Trial ». *Clinical Infectious Diseases* 61 (4): 517-26. <https://doi.org/10.1093/cid/civ363>.
- Debrah, Linda Batsa, Norman Nausch, Vera Serwaa Opoku, Wellington Owusu, Yusif Mubarik, Daniel Antwi Berko, Samuel Wanji, et al. 2017. « Epidemiology of *Mansonella perstans* in the middle belt of Ghana ». *Parasites & Vectors* 10 (janvier):15. <https://doi.org/10.1186/s13071-016-1960-0>.
- Demanou, M., S. D. Pion, et M. Boussinesq. 2001. « [Entomologic study of loaiasis transmission in the Lekie area (Cameroon)] ». *Bulletin De La Societe De Pathologie Exotique (1990)* 94 (4): 347-52.
- Dembélé, Pascal, Mady Cissoko, Adama Zan Diarra, Lassana Doumbia, Aïssata Koné, Mahamadou H. Magassa, Maissane Mehadji, Mahamadou A. Thera, et Stéphane Ranque. 2024. « Evaluation of the Performance of Rapid Diagnostic Tests for Malaria Diagnosis and Mapping of Different Plasmodium Species in Mali ». *International*

- Journal of Environmental Research and Public Health* 21 (2): 228.
<https://doi.org/10.3390/ijerph21020228>.
- Deng, Miao-Han, Lan-Yi Zhong, Okanurak Kamolnetr, Yanin Limpanont, et Zhi-Yue Lv. 2019. « Detection of Helminths by Loop-Mediated Isothermal Amplification Assay: A Review of Updated Technology and Future Outlook ». *Infectious Diseases of Poverty* 8 (1): 20. <https://doi.org/10.1186/s40249-019-0530-z>.
- Desjardins, Christopher A, Gustavo C Cerqueira, Jonathan M Goldberg, Julie C Dunning Hotopp, Brian J Haas, Jeremy Zucker, José M C Ribeiro, et al. 2013. « Genomics of Loa Loa, a Wolbachia-Free Filarial Parasite of Humans ». *Nature Genetics* 45 (5): 495-500. <https://doi.org/10.1038/ng.2585>.
- Desoubeaux, Guillaume, Adélaïde Chesnay, et Jacques Chandenier. 2020. « Approches diagnostiques des filarioses en France métropolitaine ». *Revue Francophone des Laboratoires* 2020 (519): 72-80. [https://doi.org/10.1016/S1773-035X\(20\)30066-6](https://doi.org/10.1016/S1773-035X(20)30066-6).
- Devi, R Renuka, et V Raju. 2018. « Information Technology in Morbidity Management of Human Lymphatic Filariasis—A Promising Tool in Global Programme for Elimination of Lymphatic Filariasis ». *Journal of Vector Borne Diseases* 55 (1): 20. <https://doi.org/10.4103/0972-9062.234622>.
- Diagne, Jean Pierre, Aly Mbara Ka, Joseph Matar Ndiaye, ABoubacry Sadikh Sow, Aissatou Maguatte Wane, et Paule Aida Ndoeye Roth. 2018. « MIGRATION SOUS CUTANEE PALPEBRALE D'UNE LOASE: A PROPOS D'UN CAS », n° 1, 4. https://www.revues-ufhb-ci.org/fichiers/FICHIR_ARTICLE_2511.pdf.
- Diallo, S., M. Sarr, S. Diagne, et L. Konate. 1977. « Filariasis in the Nianing région" Senegal. Prevalence, clinical aspects, entomological findings. » *Medecine d'Afrique Noire* 24 (4): 233-42. <https://www.cabdirect.org/cabdirect/abstract/19772902103>.
- Díaz-Menéndez, Marta, Francesca Norman, Begoña Monge-Maillo, José Antonio Pérez-Molina, et Rogelio López-Vélez. 2011. « Las filariasis en la práctica clínica ». *Enfermedades Infecciosas y Microbiología Clínica* 29 (décembre):27-37. [https://doi.org/10.1016/S0213-005X\(11\)70041-6](https://doi.org/10.1016/S0213-005X(11)70041-6).
- Diwakar, Deepak Kumar, Nikita Wadhwani, Nikhil Prasad, et Ashwani Gupta. 2018. « Primary Retroperitoneal Filariasis: A Common Disease of Tropics with Uncommon Presentation and Review of Literature ». *BMJ Case Reports*, septembre, bcr-2018-226217. <https://doi.org/10.1136/bcr-2018-226217>.
- Dolo, Housseini, Yaya Ibrahim Coulibaly, Louise Kelly-Hope, Siaka Konate, Benoit Dembele, Siaka Yamoussa Coulibaly, Dramane Sanogo, et al. 2018. « Factors Associated with

- Wuchereria Bancrofti Microfilaremia in an Endemic Area of Mali ». *The American Journal of Tropical Medicine and Hygiene* 98 (6): 1782-87.
<https://doi.org/10.4269/ajtmh.17-0902>.
- Downes, B, et K Jacobsen. 2010. « A systematic review of the epidemiology of mansonelliasis ». *African Journal of Infectious Diseases* 4 (1).
<https://doi.org/10.4314/ajid.v4i1.55085>.
- Downes, Barbara L., et Kathryn H. Jacobsen. 2010. « A Systematic Review of the Epidemiology of Mansonelliasis ». *African Journal of Infectious Diseases* 4 (1): 7-14.
<https://doi.org/10.4314/ajid.v4i1.55085>.
- Drame, Papa M., Céline Montavon, Sébastien D. Pion, Joseph Kubofcik, Michael P. Fay, et Thomas B. Nutman. 2016. « Molecular Epidemiology of Blood-Borne Human Parasites in a Loa Loa-, Mansonella Perstans-, and Plasmodium Falciparum-Endemic Region of Cameroon ». *The American Journal of Tropical Medicine and Hygiene* 94 (6): 1301-8.
<https://doi.org/10.4269/ajtmh.15-0746>.
- Duong, Thanh Hai, et Dominique Richard-Lenoble. 2008. « Diagnostic des parasitoses à parasites sanguicoles ». *Revue Francophone des Laboratoires* 2008 (399): 29-39.
[https://doi.org/10.1016/S1773-035X\(08\)70180-1](https://doi.org/10.1016/S1773-035X(08)70180-1).
- Ehrens, Alexandra, Achim Hoerauf, et Marc P. Hübner. 2022. « Eosinophils in filarial infections: Inducers of protection or pathology? » *Frontiers in Immunology* 13 (octobre):983812. <https://doi.org/10.3389/fimmu.2022.983812>.
- El Setouhy, Maged, Reda M. R. Ramzy, Ehab S. Ahmed, Amr M. Kandil, Omar Hussain, Hoda A. Farid, Hanan Helmy, et Gary J. Weil. 2004. « A Randomized Clinical Trial Comparing Single- and Multi-Dose Combination Therapy with Diethylcarbamazine and Albendazole for Treatment of Bancroftian Filariasis ». *The American Journal of Tropical Medicine and Hygiene* 70 (2): 191-96.
- Emukah, Emmanuel, Lindsay J. Rakers, Barminas Kahansim, Emmanuel S. Miri, Bertram E. B. Nwoke, Emily Griswold, Yisa Saka, et al. 2018. « In Southern Nigeria Loa loa Blood Microfilaria Density is Very Low Even in Areas with High Prevalence of Loiasis: Results of a Survey Using the New LoaScope Technology ». *The American Journal of Tropical Medicine and Hygiene* 99 (1): 116-23. <https://doi.org/10.4269/ajtmh.18-0163>.
- Erickson, Sara M., Edward K. Thomsen, John B. Keven, Naomi Vincent, Gussy Koimbu, Peter M. Siba, Bruce M. Christensen, et Lisa J. Reimer. 2013. « Mosquito-Parasite Interactions Can Shape Filariasis Transmission Dynamics and Impact Elimination

- Programs ». Édité par Jesus G. Valenzuela. *PLoS Neglected Tropical Diseases* 7 (9): e2433. <https://doi.org/10.1371/journal.pntd.0002433>.
- Famakinde, Damilare. 2018. « Mosquitoes and the Lymphatic Filarial Parasites: Research Trends and Budding Roadmaps to Future Disease Eradication ». *Tropical Medicine and Infectious Disease* 3 (1): 4. <https://doi.org/10.3390/tropicalmed3010004>.
- Faye, O., B. Faye, J. L. Ndiaye, R. C. K. Tine, D. Ndiaye, et A. Haydara Mbacké. 2009. « [Epidemiological study on lymphatic filariasis (*Wuchereria bancrofti*) in Sénégal] ». *Medecine Tropicale: Revue Du Corps De Sante Colonial* 69 (1): 101-2.
- Ferreira, Marcelo U., James Lee Crainey, et Federico G. Gobbi. 2023. « The Search for Better Treatment Strategies for Mansonellosis: An Expert Perspective ». *Expert Opinion on Pharmacotherapy* 24 (15): 1685-92. <https://doi.org/10.1080/14656566.2023.2240235>.
- Ferreira, Marcelo U., James Lee Crainey, et Sérgio L.B. Luz. 2021. « *Mansonella Ozzardi* ». *Trends in Parasitology* 37 (1): 90-91. <https://doi.org/10.1016/j.pt.2020.03.005>.
- Frédéric, NAHMIAS. 2018. « Haute Autorité de santé ».
- Gaillard, Colette Marie, Sebastien David Pion, Hadjira Hamou, Constant Sirima, Charlotte Bizet, Thomas Lemarcis, Jules Rodrigues, et al. 2020. « Detection of DNA of *Filariae* Closely Related to *Mansonella Perstans* in Faecal Samples from Wild Non-Human Primates from Cameroon and Gabon ». *Parasites & Vectors* 13 (1): 313. <https://doi.org/10.1186/s13071-020-04184-1>.
- Gardon, Jacques, Nathalie Gardon-Wendel, Demanga-Ngangue, Joseph Kamgno, Jean-Philippe Chippaux, et Michel Boussinesq. 1997. « Serious Reactions after Mass Treatment of Onchocerciasis with Ivermectin in an Area Endemic for Loa Loa Infection ». *The Lancet* 350 (9070): 18-22. [https://doi.org/10.1016/S0140-6736\(96\)11094-1](https://doi.org/10.1016/S0140-6736(96)11094-1).
- Geelhand De Merxem, D., J. N. Siewe Fodjo, S. Menon, A. Hotterbeekx, et R. Colebunders. 2019. « Nodding Syndrome Research, Lessons Learned from the NSETHIO Project ». *Global Mental Health* 6:e26. <https://doi.org/10.1017/gmh.2019.24>.
- Gentilini, M., J.M. Pinon, et M. Danis. 1973. « Diagnostic biologique des filarioses humaines ». *Médecine et Maladies Infectieuses* 3 (8-9): 351-53. [https://doi.org/10.1016/S0399-077X\(73\)80130-1](https://doi.org/10.1016/S0399-077X(73)80130-1).
- Gobbi, Federico, Anna Beltrame, Dora Buonfrate, Silvia Staffolani, Monica Degani, Maria Gobbo, Andrea Angheben, Stefania Marocco, et Zeno Bisoffi. 2017. « Imported Infections with *Mansonella perstans* Nematodes, Italy ». *Emerging Infectious Diseases* 23 (9): 1539-42. <https://doi.org/10.3201/eid2309.170263>.

- González, Mikel A., Erin Dilger, María M. Ronderos, Gustavo R. Spinelli, Orin Courtenay, et James G. C. Hamilton. 2020. « Significant Reduction in Abundance of Peridomestic Mosquitoes (Culicidae) and Culicoides Midges (Ceratopogonidae) after Chemical Intervention in Western São Paulo, Brazil ». *Parasites & Vectors* 13 (1): 549. <https://doi.org/10.1186/s13071-020-04427-1>.
- Gyasi, Michael E., Ogugua N. Okonkwo, et Koushik Tripathy. 2024. « Onchocerciasis ». In *StatPearls*. Treasure Island (FL): StatPearls Publishing. <http://www.ncbi.nlm.nih.gov/books/NBK559027/>.
- Haddix, Anne C., et Andrew Kestler. 2000. « Lymphatic Filariasis: Economic Aspects of the Disease and Programmes for Its Elimination ». *Transactions of the Royal Society of Tropical Medicine and Hygiene* 94 (6): 592-93. [https://doi.org/10.1016/S0035-9203\(00\)90199-8](https://doi.org/10.1016/S0035-9203(00)90199-8).
- Harrup, Fabio Macchioni, Luca Chelucci, et Beatrice Torracca. 2016. « Advances in Control Techniques for Culicoides and Future Prospects ». *Veterinaria Italiana* 52 (3): 247-64. <https://doi.org/10.12834/VetIt.741.3602.3>.
- Hartley, Marie. 2021. « Onchocerciasis (river blindness) », 2021. <https://dermnetnz.org/topics/onchocerciasis>.
- Herbeuval, H., R. Herbeuval, et J. Duheille. 1962. « [The passage, in the circulating blood, of hematopoietic masses, erythroblastic islands, megakaryoblastic islands, mixed islands] ». *Nouvelle Revue Francaise D'hematologie* 2:616-19.
- Herricks, Jennifer R., Peter J. Hotez, Valentine Wanga, Luc E. Coffeng, Juanita A. Haagsma, María-Gloria Basáñez, Geoffrey Buckle, et al. 2017. « The Global Burden of Disease Study 2013: What Does It Mean for the NTDs? » Édité par Xiao-Nong Zhou. *PLOS Neglected Tropical Diseases* 11 (8): e0005424. <https://doi.org/10.1371/journal.pntd.0005424>.
- Hotez, Peter J., et Aruna Kamath. 2009. « Neglected Tropical Diseases in Sub-Saharan Africa: Review of Their Prevalence, Distribution, and Disease Burden ». Édité par Michael Cappello. *PLoS Neglected Tropical Diseases* 3 (8): e412. <https://doi.org/10.1371/journal.pntd.0000412>.
- Hotez, Peter J., David H. Molyneux, Alan Fenwick, Jacob Kumaresan, Sonia Ehrlich Sachs, Jeffrey D. Sachs, et Lorenzo Savioli. 2007. « Control of Neglected Tropical Diseases ». *New England Journal of Medicine* 357 (10): 1018-27. <https://doi.org/10.1056/NEJMra064142>.

- Ibe, Ogochukwu, Obinna Onwujekwe, Benjamin Uzochukwu, Miriam Ajuba, et Paul Okonkwo. 2015. « Exploring Consumer Perceptions and Economic Burden of Onchocerciasis on Households in Enugu State, South-East Nigeria ». Édité par Samuel V. Scarpino. *PLOS Neglected Tropical Diseases* 9 (11): e0004231. <https://doi.org/10.1371/journal.pntd.0004231>.
- Jacobsen, Kathryn H, Bailey C Andress, Elina A Bhagwat, Ciera A Bryant, Vipusha R Chandrapu, Caroline G Desmonts, Tania M Matthews, Akeem Ogunkoya, Tristan J Wheeler, et A'kayla S Williams. 2022. « A Call for Loiasis to Be Added to the WHO List of Neglected Tropical Diseases ». *The Lancet Infectious Diseases* 22 (10): e299-302. [https://doi.org/10.1016/S1473-3099\(22\)00064-0](https://doi.org/10.1016/S1473-3099(22)00064-0).
- Jain, Rajeev, Jern Y. Chen, Andrew R. Butcher, Robert Casson, et Dinesh Selva. 2008. « Subconjunctival Loa Loa Worm ». *International Journal of Infectious Diseases* 12 (6): e133-35. <https://doi.org/10.1016/j.ijid.2008.03.001>.
- James, Spencer L, Degu Abate, Kalkidan Hassen Abate, Solomon M Abay, Cristiana Abbafati, Nooshin Abbasi, Hedayat Abbastabar, et al. 2018. « Global, Regional, and National Incidence, Prevalence, and Years Lived with Disability for 354 Diseases and Injuries for 195 Countries and Territories, 1990–2017: A Systematic Analysis for the Global Burden of Disease Study 2017 ». *The Lancet* 392 (10159): 1789-1858. [https://doi.org/10.1016/S0140-6736\(18\)32279-7](https://doi.org/10.1016/S0140-6736(18)32279-7).
- Kamgno, Joseph, Sébastien D. Pion, Cédric B. Chesnais, Matthew H. Bakalar, Michael V. D'Ambrosio, Charles D. Mackenzie, Hugues C. Nana-Djeunga, et al. 2017. « A Test-and-Not-Treat Strategy for Onchocerciasis in *Loa Loa* –Endemic Areas ». *New England Journal of Medicine* 377 (21): 2044-52. <https://doi.org/10.1056/NEJMoa1705026>.
- Katawa, Gnatoulma, Laura E. Layland, Alex Y. Debrah, Charlotte Von Horn, Linda Batsa, Alexander Kwarteng, Sandra Arriens, et al. 2015. « Hyperreactive Onchocerciasis Is Characterized by a Combination of Th17-Th2 Immune Responses and Reduced Regulatory T Cells ». Édité par Benjamin L. Makepeace. *PLoS Neglected Tropical Diseases* 9 (1): e3414. <https://doi.org/10.1371/journal.pntd.0003414>.
- Katiyar, D., et L. K. Singh. 2011. « Filariasis: Current Status, Treatment and Recent Advances in Drug Development ». *Current Medicinal Chemistry* 18 (14): 2174-85. <https://doi.org/10.2174/092986711795656234>.
- Keiser, Paul B., Yaya I. Coulibaly, Falaye Keita, Diakaridia Traore, Abdallah Diallo, Dapa A. Diallo, Roshanak T. Semnani, et al. 2003. « Clinical Characteristics of Post-Treatment Reactions to Ivermectin/Albendazole for *Wuchereria Bancrofti* in a Region Co-

- Endemic for *Mansonella Perstans* ». *The American Journal of Tropical Medicine and Hygiene* 69 (3): 331-35.
- Keiser, Paul B., Yaya Coulibaly, Joseph Kubofcik, Abdallah A. Diallo, Amy D. Klion, Sekou F. Traoré, et Thomas B. Nutman. 2008. « Molecular Identification of *Wolbachia* from the Filarial Nematode *Mansonella Perstans* ». *Molecular and Biochemical Parasitology* 160 (2): 123-28. <https://doi.org/10.1016/j.molbiopara.2008.04.012>.
- Keita, Mamadou Fodé. 1979. « Aspects épidémiologiques des filarioses à *onchocerca volvulus*, *wuchereria bancrofti* et *tetrapetatonema perstans* ». Thesis, Ecole Nationale de Médecine et de Pharmacie. <https://www.bibliosante.ml/handle/123456789/9605>.
- Kelly-Hope, Louise, Rossely Paulo, Brent Thomas, Miguel Brito, Thomas R. Unnasch, et David Molyneux. 2017. « *Loa Loa* Vectors *Chrysops* Spp.: Perspectives on Research, Distribution, Bionomics, and Implications for Elimination of Lymphatic Filariasis and Onchocerciasis ». *Parasites & Vectors* 10 (1): 172. <https://doi.org/10.1186/s13071-017-2103-y>.
- Khan, A. M., P. Dutta, S. Das, A. K. Pathak, P. Sarmah, M. E. Hussain, et J. Mahanta. 2015. « Microfilarial Periodicity of *Wuchereria Bancrofti* in Assam, Northeast India ». *Journal of Vector Borne Diseases* 52 (3): 208-12.
- Kima, A., K.T. Guiguemde, M. Serme, Z.C. Meda, R. Bougma, J.P. Djiatsa, C. Bougouma, et F. Drabo. 2021. « Évaluation de la Transmission de la Filariose Lymphatique au Burkina Faso: À Propos de 4 Districts Sanitaires ». *Meédecine Tropicale et Santéé Internationale* 1 (2): mtsibulletin.n1.2021.83. <https://doi.org/10.48327/mtsibulletin.n1.2021.83>.
- Knopp, Stefanie, Peter Steinmann, Christoph Hatz, Jennifer Keiser, et Jürg Utzinger. 2012. « Nematode Infections »: *Infectious Disease Clinics of North America* 26 (2): 359-81. <https://doi.org/10.1016/j.idc.2012.02.005>.
- Komlan, Kossi, Patrick S. Vossberg, Richard G. Gantim, Tchalim Solim, Francois Korbmacher, Méba Banla, Koffi Padjoudoum, Potchoziou Karabou, Carsten Köhler, et Peter T. Soboslay. 2018. « *Onchocerca Volvulus* Infection and Serological Prevalence, Ocular Onchocerciasis and Parasite Transmission in Northern and Central Togo after Decades of *Simulium Damnosum* s.l. Vector Control and Mass Drug Administration of Ivermectin ». Édité par Matthew C. Freeman. *PLOS Neglected Tropical Diseases* 12 (3): e0006312. <https://doi.org/10.1371/journal.pntd.0006312>.

763 Kwarteng, Alexander, Samuel Terkper Ahuno, et Freda Osei Akoto. 2016. « Killing Filarial
 764 Nematode Parasites: Role of Treatment Options and Host Immune Response ». *Infectious Diseases of Poverty* 5 (1): 86. <https://doi.org/10.1186/s40249-016-0183-0>.
 765
 766 Laidoudi, Younes, Jean-Lou Marie, Djamel Tahir, Stéphanie Watier-Grillot, Oleg
 767 Mediannikov, et Bernard Davoust. 2020. « Detection of Canine Vector-Borne Filariasis
 768 and Their Wolbachia Endosymbionts in French Guiana ». *Microorganisms* 8 (5): 770.
 769 <https://doi.org/10.3390/microorganisms8050770>.
 770 Lima, Nathália F., Cecilia A. Veggiani Aybar, María J. Dantur Juri, et Marcelo U. Ferreira.
 771 2016. « *Mansonella Ozzardi* : A Neglected New World Filarial Nematode ». *Pathogens*
 772 *and Global Health* 110 (3): 97-107. <https://doi.org/10.1080/20477724.2016.1190544>.
 773 Lustigman, Sara, Alexandra Grote, et Elodie Ghedin. 2017. « The Role of “omics” in the Quest
 774 to Eliminate Human Filariasis ». Édité par Judd L. Walson. *PLOS Neglected Tropical*
 775 *Diseases* 11 (4): e0005464. <https://doi.org/10.1371/journal.pntd.0005464>.
 776 M. Boussinesq. 2007. « Actualités thérapeutiques dans les filarioses – Recent advances in the
 777 treatment of filariases ». *mise au point*, n° 2, 55.
 778 Mandal, Debasish, Dipankar Roy, D. K. Bera, et Buddhadeb Manna. 2012. « Occurrence of
 779 Gravid Loa Loa in Subconjunctival Space of Man: A Case Report from West Bengal,
 780 India ». *Journal of Parasitic Diseases*, juillet. [https://doi.org/10.1007/s12639-012-](https://doi.org/10.1007/s12639-012-0130-3)
 781 [0130-3](https://doi.org/10.1007/s12639-012-0130-3).
 782 Manguin, S., M.J. Bangs, J. Pothikasikorn, et T. Chareonviriyaphap. 2010. « Review on Global
 783 Co-Transmission of Human Plasmodium Species and Wuchereria Bancrofti by
 784 Anopheles Mosquitoes ». *Infection, Genetics and Evolution* 10 (2): 159-77.
 785 <https://doi.org/10.1016/j.meegid.2009.11.014>.
 786 McGarry, Helen F, Ken Pfarr, Gill Egerton, Achim Hoerauf, Jean-Paul Akue, Peter Enyong,
 787 Samuel Wanji, et al. 2003. « [No title found] ». *Filaria Journal* 2 (1): 9.
 788 <https://doi.org/10.1186/1475-2883-2-9>.
 789 Meda, N., M.A. Daboue, W.P. Djiguimde, C. Meda, et S. Konate/Ouedraogo. 2009. « 571
 790 Manifestations oculaires de la loase ». *Journal Français d’Ophtalmologie* 32
 791 (avril):1S173. [https://doi.org/10.1016/S0181-5512\(09\)73695-2](https://doi.org/10.1016/S0181-5512(09)73695-2).
 792 Medeiros, Jansen Fernandes, James Lee Crainey, Felipe Arley Costa Pessoa, et Sérgio Luis
 793 Bessa Luz. 2017. « Mansonelliasis ». In *Arthropod Borne Diseases*, édité par Carlos
 794 Brisola Marcondes, 405-26. Cham: Springer International Publishing.
 795 https://doi.org/10.1007/978-3-319-13884-8_26.

796 Mediannikov, O., et S. Ranque. 2018a. « Mansonellosis, the Most Neglected Human
797 Filariasis ». *New Microbes and New Infections* 26 (novembre):S19-22.
798 <https://doi.org/10.1016/j.nmni.2018.08.016>.

799 ———. 2018b. « Mansonellosis, the Most Neglected Human Filariasis ». *New Microbes and*
800 *New Infections* 26 (novembre):S19-22. <https://doi.org/10.1016/j.nmni.2018.08.016>.

801 Melrose, Wayne D. 2002. « Lymphatic Filariasis: New Insights into an Old Disease ». *International Journal for Parasitology* 32 (8): 947-60. [https://doi.org/10.1016/S0020-](https://doi.org/10.1016/S0020-7519(02)00062-0)
802 [7519\(02\)00062-0](https://doi.org/10.1016/S0020-7519(02)00062-0).

804 Mendoza, Natalia, Alexa Li, Amber Gill, et Stephen Tying. 2009. « Filariasis: Diagnosis and
805 Treatment ». *Dermatologic Therapy* 22 (6): 475-90. [https://doi.org/10.1111/j.1529-](https://doi.org/10.1111/j.1529-8019.2009.01271.x)
806 [8019.2009.01271.x](https://doi.org/10.1111/j.1529-8019.2009.01271.x).

807 Métais, Alice, Sophie Michalak, et Audrey Rousseau. 2021. « Albendazole-Related Loa Loa
808 Encephalopathy ». *IDCases* 23:e01033. <https://doi.org/10.1016/j.idcr.2020.e01033>.

809 Metzger, Wolfram Gottfried, et Benjamin Mordmüller. 2014. « Loa Loa—Does It Deserve to
810 Be Neglected? » *The Lancet Infectious Diseases* 14 (4): 353-57.
811 [https://doi.org/10.1016/S1473-3099\(13\)70263-9](https://doi.org/10.1016/S1473-3099(13)70263-9).

812 Molyneux, David H., et Mark J. Taylor. 2001. « Current Status and Future Prospects of the
813 Global Lymphatic Filariasis Programme »: *Current Opinion in Infectious Diseases* 14
814 (2): 155-59. <https://doi.org/10.1097/00001432-200104000-00008>.

815 Mourembou, Gaël, Florence Fenollar, Jean Bernard Lekana-Douki, Angelique Ndjoi
816 Mbiguino, Sydney Maghendji Nzondo, Pierre Blaise Matsiegui, Rella Zoleko Manego,
817 et al. 2015a. « Mansonella, Including a Potential New Species, as Common Parasites in
818 Children in Gabon ». Édité par Dunstan Mukoko. *PLOS Neglected Tropical Diseases* 9
819 (10): e0004155. <https://doi.org/10.1371/journal.pntd.0004155>.

820 ———. 2015b. « Mansonella, including a Potential New Species, as Common Parasites in
821 Children in Gabon ». *PLoS Neglected Tropical Diseases* 9 (10): e0004155.
822 <https://doi.org/10.1371/journal.pntd.0004155>.

823 Ngwewondo, Adela, Ivan Scandale, et Sabine Specht. 2021. « Onchocerciasis Drug
824 Development: From Preclinical Models to Humans ». *Parasitology Research* 120 (12):
825 3939-64. <https://doi.org/10.1007/s00436-021-07307-4>.

826 Nizeyimana, Jean Baptiste. 2021. « ÉTUDE DES CONDITIONS DE DEPISTAGE
827 CLINIQUE ET LES EFFETS PSYCHO SOCIOECONOMIQUE DES PERSONNES
828 PORTEUSES DE L'ELEPHANTIASIS DES JAMBES. CAS DE

L'ARRONDISSEMENT DE GUIDIGUIS CAS DE LA REGION DE L'EXTREME-NORD CAMEROUN DE 2017-2020 Par » 26 (03).

Njim, Tsi, Joel Mbigha Ngum, et Leopold Ndemnge Aminde. 2015. « Cutaneous Onchocerciasis in Dumbu, a Pastoral Area in the North-West Region of Cameroon: Diagnostic Challenge and Socio-Economic Implications ». *Pan African Medical Journal* 22. <https://doi.org/10.11604/pamj.2015.22.298.7707>.

Njoku, Chidiebere, Lawrence Nwankwo, Masayo Ozaki, Andrew Obasi, Chinyere N. Ukaga, Emmanuel Emukah, Kenrick Nwodu, et al. 2013. « Community-Wide Distribution of Long-Lasting Insecticidal Nets Can Halt Transmission of Lymphatic Filariasis in Southeastern Nigeria ». *The American Journal of Tropical Medicine and Hygiene* 89 (3): 578-87. <https://doi.org/10.4269/ajtmh.12-0775>.

Noroës, Joaquim, David Addiss, Agnaldo Cedenho, Jose Figueredo-Silvan, Guilherme Lima, et Gerusa Dreyer. 2003. « Pathogenesis of Filarial Hydrocele: Risk Associated with Intrascrotal Nodules caused by Death of Adult Wuchereria Bancrofti ». *Transactions of the Royal Society of Tropical Medicine and Hygiene* 97 (5): 561-66. [https://doi.org/10.1016/S0035-9203\(03\)80029-9](https://doi.org/10.1016/S0035-9203(03)80029-9).

Nutman, Thomas B. 2013. « Insights into the Pathogenesis of Disease in Human Lymphatic Filariasis ». *Lymphatic Research and Biology* 11 (3): 144-48. <https://doi.org/10.1089/lrb.2013.0021>.

O'Hanlon, Simon J., Hannah C. Slater, Robert A. Cheke, Boakye A. Boatın, Luc E. Coffeng, Sébastien D. S. Pion, Michel Boussinesq, Honorat G. M. Zouré, Wilma A. Stolk, et María-Gloria Basáñez. 2016. « Model-Based Geostatistical Mapping of the Prevalence of Onchocerca Volvulus in West Africa ». Édité par Ricardo J. Soares Magalhaes. *PLOS Neglected Tropical Diseases* 10 (1): e0004328. <https://doi.org/10.1371/journal.pntd.0004328>.

Ojurongbe, Olusola, Akeem Abiodun Akindele, Monsuru Adebayo Adeleke, Matthew Oyebo Oyebo, Samuel Adeyinka Adedokun, Josephine Folashade Ojo, Callistus Adewale Akinleye, Oloyede Samuel Bolaji, Olusegun Adelowo Adefioye, et Oluwaseyi Adegboyega Adeyeba. 2015. « Co-endemicity of Loiasis and Onchocerciasis in Rain Forest Communities in Southwestern Nigeria ». *PLOS Neglected Tropical Diseases* 9 (3): e0003633. <https://doi.org/10.1371/journal.pntd.0003633>.

« OMS MALI Rapport annuel 2018.pdf ». s. d. Consulté le 24 juillet 2024. <https://www.afro.who.int/sites/default/files/2019-01/OMS%20MALI%20Rapport%20annuel%202018.pdf>.

- Opoku, Millicent, Corrado Minetti, Worlasi D. Kartey-Attipoe, Sampson Otoo, Joseph Otchere, Bruno Gomes, Dziedzom K. De Souza, et Lisa J. Reimer. 2018. « An Assessment of Mosquito Collection Techniques for Xenomonitoring of Anopheline-Transmitted Lymphatic Filariasis in Ghana ». *Parasitology* 145 (13): 1783-91. <https://doi.org/10.1017/S0031182018000938>.
- Ottesen, Eric A. 2000. « Editorial: The Global Programme to Eliminate Lymphatic Filariasis ». *Tropical Medicine & International Health* 5 (9): 591-94. <https://doi.org/10.1046/j.1365-3156.2000.00620.x>.
- . 2006. « Lymphatic Filariasis: Treatment, Control and Elimination ». In *Advances in Parasitology*, 61:395-441. Elsevier. [https://doi.org/10.1016/S0065-308X\(05\)61010-X](https://doi.org/10.1016/S0065-308X(05)61010-X).
- Ouedraogo, Amina Nomtondo, Emile Bégnouriti Somda, Fagnima Traoré, Muriel Sidnoma Ouédraogo, Gilbert Patrice Tapsoba, Angèle Ouangre/Ouédraogo, Salamata Bara, et al. 2016. « Impact du traitement de masse de la filariose lymphatique par l'albendazole-ivermectine en zone de savane : cas de la région de l'Est du Burkina ». *HEALTH SCIENCES AND DISEASE* 17 (4). <http://hsd-fmsb.org/index.php/hsd/article/view/731>.
- Pampiglione, S., G. Canestri Trotti, et F. Rivasi. 1995. « Human Dirofilariasis Due to *Dirofilaria (Noctiella) Repens*: A Review of World Literature ». *Parassitologia* 37 (2-3): 149-93.
- Pérez-Molina, José Antonio, Marta Díaz-Menéndez, Ana Pérez-Ayala, Federico Ferrere, Begoña Monje, Francesca Norman, et Rogelio López-Vélez. 2010. « Tratamiento de las enfermedades causadas por parásitos ». *Enfermedades Infecciosas y Microbiología Clínica* 28 (1): 44-59. <https://doi.org/10.1016/j.eimc.2009.11.003>.
- Phillips, Richard O., Michael Frimpong, Fred S. Sarfo, Birte Kretschmer, Marcus Beissner, Alexander Debrah, Yaw Ampem-Amoako, et al. 2014. « Infection with *Mansonella perstans* Nematodes in Buruli Ulcer Patients, Ghana ». *Emerging Infectious Diseases* 20 (6): 1000-1003. <https://doi.org/10.3201/eid2006.131501>.
- Pion, S. D. S., J. A. N. Filipe, J. Kamgno, J. Gardon, M.-G. Basáñez, et M. Boussinesq. 2006. « Microfilarial Distribution of *Loa Loa* in the Human Host: Population Dynamics and Epidemiological Implications ». *Parasitology* 133 (01): 101. <https://doi.org/10.1017/S0031182006000035>.
- Pion, Sébastien Ds, Hugues Nana-Djeunga, Yannick Niamsi-Emalio, Cédric B Chesnais, Hugo Deléglise, Charles Mackenzie, Wilma Stolk, et al. 2020. « Implications for Annual Retesting after a Test-and-Not-Treat Strategy for Onchocerciasis Elimination in Areas

- Co-Endemic with Loa Loa Infection: An Observational Cohort Study ». *The Lancet Infectious Diseases* 20 (1): 102-9. [https://doi.org/10.1016/S1473-3099\(19\)30554-7](https://doi.org/10.1016/S1473-3099(19)30554-7).
- Poole, Catherine B., Zhiru Li, Andy Alhassan, Dylan Guelig, Steven Diesburg, Nathan A. Tanner, Yinhua Zhang, et al. 2017. « Colorimetric Tests for Diagnosis of Filarial Infection and Vector Surveillance Using Non-Instrumented Nucleic Acid Loop-Mediated Isothermal Amplification (NINA-LAMP) ». Édité par Matty Knight. *PLOS ONE* 12 (2): e0169011. <https://doi.org/10.1371/journal.pone.0169011>.
- Porter, Suzanne, et Adrian Hopkins. 2015. « Élimination de l'onchocercose et de la filariose lymphatique » 12.
- Puente, Sabino, Mar Lago, Mercedes Subirats, Ismael Sanz-Esteban, Marta Arsuaga, Belén Vicente, Montserrat Alonso-Sardon, Moncef Belhassen-Garcia, et Antonio Muro. 2020. « Imported Mansonella perstans infection in Spain ». *Infectious Diseases of Poverty* 9 (1): 105. <https://doi.org/10.1186/s40249-020-00729-9>.
- Ramaiah, K. D., et Eric A. Ottesen. 2014. « Progress and Impact of 13 Years of the Global Programme to Eliminate Lymphatic Filariasis on Reducing the Burden of Filarial Disease ». Édité par Moses Bockarie. *PLoS Neglected Tropical Diseases* 8 (11): e3319. <https://doi.org/10.1371/journal.pntd.0003319>.
- Ramharter, Michael, Joanna Butler, Ghyslain Mombo-Ngoma, Tamara Nordmann, Saskia Dede Davi, et Rella Zoleko Manego. 2023. « The African Eye Worm: Current Understanding of the Epidemiology, Clinical Disease, and Treatment of Loiasis ». *The Lancet Infectious Diseases*, octobre, S1473309923004383. [https://doi.org/10.1016/S1473-3099\(23\)00438-3](https://doi.org/10.1016/S1473-3099(23)00438-3).
- Rebollo, Maria P., et Moses J. Bockarie. 2017. « Can Lymphatic Filariasis Be Eliminated by 2020? » *Trends in Parasitology* 33 (2): 83-92. <https://doi.org/10.1016/j.pt.2016.09.009>.
- Richardson, Eugene T., Robert Luo, Doran L. Fink, Thomas B. Nutman, John K. Geisse, et Michele Barry. 2012. « Transient Facial Swellings in a Patient With a Remote African Travel History ». *Journal of Travel Medicine* 19 (3): 183-85. <https://doi.org/10.1111/j.1708-8305.2012.00612.x>.
- Risch, Frederic, Manuel Ritter, Achim Hoerauf, et Marc P. Hübner. 2021. « Human Filariasis—Contributions of the Litomosoides Sigmodontis and Acanthocheilonema Viteae Animal Model ». *Parasitology Research* 120 (12): 4125-43. <https://doi.org/10.1007/s00436-020-07026-2>.
- Robert, Vincent, E.H. Ndiaye, Nil Rahola, Gilbert Le Goff, Philippe Boussès, D. Diallo, Voahirana Le Goff, Lauriane Mariamé, et M. Diallo. 2022. « Clés dichotomiques

illustrées d'identification des femelles et des larves de moustiques (Diptera : Culicidae) du Burkina Faso, Cap-Vert, Gambie, Mali, Mauritanie, Niger, Sénégal et Tchad », 181 p. <https://doi.org/10.23708/FDI:010084866>.

Rodi, Miriam, Caspar Gross, Thaisa Lucas Sandri, Lilith Berner, Marina Marcet-Houben, Ersoy Kocak, Michaela Pogoda, et al. 2023a. « Whole genome analysis of two sympatric human *Mansonella*: *Mansonella perstans* and *Mansonella* sp “DEUX” ». *Frontiers in Cellular and Infection Microbiology* 13 (avril):1159814. <https://doi.org/10.3389/fcimb.2023.1159814>.

———. 2023b. « Whole Genome Analysis of Two Sympatric Human *Mansonella*: *Mansonella Perstans* and *Mansonella* Sp “DEUX” ». *Frontiers in Cellular and Infection Microbiology* 13:1159814. <https://doi.org/10.3389/fcimb.2023.1159814>.

Sandri, Thaisa Lucas, Andrea Kreidenweiss, Simon Cavallo, David Weber, Sascha Juhas, Miriam Rodi, Tamirat Gebru Woldearegai, et al. 2021. « Molecular Epidemiology of *Mansonella* Species in Gabon ». *The Journal of Infectious Diseases* 223 (2): 287-96. <https://doi.org/10.1093/infdis/jiaa670>.

Sangare, Moussa Brema, Yaya Ibrahim Coulibaly, Siaka Yamoussa Coulibaly, Michel Emmanuel Coulibaly, Bourama Traore, Ilo Dicko, Ibrahim Moussa Sissoko, et al. 2018. « A cross-sectional study of the filarial and *Leishmania* co-endemicity in two ecologically distinct settings in Mali ». *Parasites & Vectors* 11 (1): 18. <https://doi.org/10.1186/s13071-017-2531-8>.

Sawadogo, Pm. 2019. « >Onchocercose au Burkina Faso : une endémie à transmission vectorielle en voie de réémergence ». *Bulletin de la société de pathologie exotique* 112 (5): 260-74. <https://doi.org/10.3166/bspe-2020-0106>.

Shelley, Aj, et S Coscarón. 2001. « Simuliid blackflies (Diptera: Simuliidae) and ceratopogonid midges (Diptera: Ceratopogonidae) as vectors of *Mansonella ozzardi* (Nematoda: Onchocercidae) in northern Argentina ». *Memórias do Instituto Oswaldo Cruz* 96 (4): 451-58. <https://doi.org/10.1590/S0074-02762001000400003>.

Shenoy, R. K. 2008. « Clinical and Pathological Aspects of Filarial Lymphedema and Its Management ». *The Korean Journal of Parasitology* 46 (3): 119. <https://doi.org/10.3347/kjp.2008.46.3.119>.

Simón, Fernando, Mar Siles-Lucas, Rodrigo Morchón, Javier González-Miguel, Isabel Mellado, Elena Carretón, et Jose Alberto Montoya-Alonso. 2012. « Human and Animal Dirofilariasis: The Emergence of a Zoonotic Mosaic ». *Clinical Microbiology Reviews* 25 (3): 507-44. <https://doi.org/10.1128/CMR.00012-12>.

- Simonsen, Paul E., Yahya A. Derua, Stephen M. Magesa, Erling M. Pedersen, Anna-Sofie Stensgaard, Mwelecele N. Malecela, et William N. Kisinza. 2014. « Lymphatic filariasis control in Tanga Region, Tanzania: status after eight rounds of mass drug administration ». *Parasites & Vectors* 7 (1): 507. <https://doi.org/10.1186/s13071-014-0507-5>.
- Simonsen, Paul E., Ambrose W. Onapa, et Santa Maria Asio. 2011a. « Mansonella Perstans Filariasis in Africa ». *Acta Tropica*, The Diagnostics and Control of Neglected Tropical Helminth Diseases, 120 (septembre):S109-20. <https://doi.org/10.1016/j.actatropica.2010.01.014>.
- . 2011b. « Mansonella Perstans Filariasis in Africa ». *Acta Tropica* 120 Suppl 1 (septembre):S109-120. <https://doi.org/10.1016/j.actatropica.2010.01.014>.
- Small, Scott T., Daniel J. Tisch, et Peter A. Zimmerman. 2014. « Molecular Epidemiology, Phylogeny and Evolution of the Filarial Nematode Wuchereria Bancrofti ». *Infection, Genetics and Evolution* 28 (décembre):33-43. <https://doi.org/10.1016/j.meegid.2014.08.018>.
- Supali, Taniawati, Is S. Ismid, Heri Wibowo, Yenny Djuardi, Esther Majawati, Praba Ginanjar, et Peter Fischer. 2006. « Estimation of the Prevalence of Lymphatic Filariasis by a Pool Screen PCR Assay Using Blood Spots Collected on Filter Paper ». *Transactions of the Royal Society of Tropical Medicine and Hygiene* 100 (8): 753-59. <https://doi.org/10.1016/j.trstmh.2005.10.005>.
- Tahita, Marc Christian, Thuy-Huong Ta-Tang, Bérénger Kaboré, Raquel Capote-Morales, Irene Molina De La Fuente, Raquel Cruces, Hamidou Ilboudo, et al. 2024. « Molecular Detection of Wuchereria Bancrofti, Loa Loa and Mansonella Perstans from Dried Blood Spots Taken from Pregnant Women in Rural Burkina Faso ». *African Journal of Parasitology, Mycology and Entomology* 1 (2): 1. <https://doi.org/10.35995/ajpme2010002>.
- Tamarozzi, Francesca, Paola Rodari, Joaquín Salas-Coronas, Emmanuel Bottieau, Fernando Salvador, Manuel Jesús Soriano-Pérez, María Isabel Cabeza-Barrera, et al. 2022. « A Large Case Series of Travel-Related *Mansonella Perstans* (Vector-Borne Filarial Nematode): A TropNet Study in Europe ». *Journal of Travel Medicine*, avril, taac048. <https://doi.org/10.1093/jtm/taac048>.
- Ta-Tang, Thuy-Huong, James Crainey, Rory J. Post, Sergio LB. Luz, et José Rubio. 2018. « Mansonellosis: Current Perspectives ». *Research and Reports in Tropical Medicine* Volume 9 (janvier):9-24. <https://doi.org/10.2147/RRTM.S125750>.

998 Ta-Tang, Thuy-Huong, Sergio L B Luz, James L Crainey, et José M Rubio. 2021a. « An
999 Overview of the Management of Mansonellosis ». *Research and Reports in Tropical*
1000 *Medicine* 12 (mai):93-105. <https://doi.org/10.2147/RRTM.S274684>.

1001 Ta-Tang, Thuy-Huong, Sergio Lb Luz, James L Crainey, et José M Rubio. 2021b. « An
1002 Overview of the Management of Mansonellosis ». *Research and Reports in Tropical*
1003 *Medicine* Volume 12 (mai):93-105. <https://doi.org/10.2147/RRTM.S274684>.

1004 Tatuene, J. Kamtchum, R. Gounoue Fotsing, T. Nkoa, JB Tchateng Mbougua, HC Nana
1005 Djeunga, J. Bopda, et Joseph Kamgno. 2014a. « Epidemiology of Loa Loa and
1006 Mansonella Perstans Filariasis in the Akonolinga Health District, Centre Region,
1007 Cameroon. » *HEALTH SCIENCES AND DISEASE* 15 (1). [https://www.hsd-](https://www.hsd-fmsb.org/index.php/hsd/article/view/297)
1008 [fmsb.org/index.php/hsd/article/view/297](https://www.hsd-fmsb.org/index.php/hsd/article/view/297).

1009 ———. 2014b. « Epidemiology of Loa Loa and Mansonella Perstans Filariasis in the
1010 Akonolinga Health District, Centre Region, Cameroon. » *HEALTH SCIENCES AND*
1011 *DISEASE* 15 (1). <https://doi.org/10.5281/hsd.v15i1.297>.

1012 Tavares da Silva, Lucyane Bastos, James Lee Crainey, Túllio Romão Ribeiro da Silva, Uziel
1013 Ferreira Suwa, Ana Carolina Paulo Vicente, Jansen Fernandes de Medeiros, Felipe
1014 Arley Costa Pessoa, et Sérgio Luiz Bessa Luz. 2017. « Molecular Verification of New
1015 World Mansonella perstans Parasitemias ». *Emerging Infectious Diseases* 23 (3):
1016 545-47. <https://doi.org/10.3201/eid2303.161159>.

1017 Taylor, Mark J, Achim Hoerauf, et Moses Bockarie. 2010. « Lymphatic Filariasis and
1018 Onchocerciasis ». *The Lancet* 376 (9747): 1175-85. [https://doi.org/10.1016/S0140-](https://doi.org/10.1016/S0140-6736(10)60586-7)
1019 [6736\(10\)60586-7](https://doi.org/10.1016/S0140-6736(10)60586-7).

1020 Thylefors, B. 2004. « Eliminating Onchocerciasis as a Public Health Problem ». *Tropical*
1021 *Medicine & International Health* 9 (4). [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-3156.2004.01226.x)
1022 [3156.2004.01226.x](https://doi.org/10.1111/j.1365-3156.2004.01226.x).

1023 Unnasch, Thomas R, Allison Golden, Vitaliano Cama, et Paul T Cantey. 2018. « Diagnostics
1024 for Onchocerciasis in the Era of Elimination ». *International Health* 10 (suppl_1):
1025 i20-26. <https://doi.org/10.1093/inthealth/ihx047>.

1026 Varenne, F., J. Fillaux, M. Porterie, J. Soler, M. Cassagne, et V. Soler. 2016. « Loase sous-
1027 conjonctivale : à propos d'un cas ». *Journal Français d'Ophtalmologie* 39 (8): e193-94.
1028 <https://doi.org/10.1016/j.jfo.2015.08.018>.

1029 Vinkes Melchers, Natalie V. S., Sarah Mollenkopf, Robert Colebunders, Michael Edlinger,
1030 Luc E. Coffeng, Julia Irani, Trésor Zola, et al. 2018. « Burden of Onchocerciasis-

- Associated Epilepsy: First Estimates and Research Priorities ». *Infectious Diseases of Poverty* 7 (1). <https://doi.org/10.1186/s40249-018-0481-9>.
- Vinkeles Melchers, Natalie V. S., Wilma A. Stolk, Welmoed Van Loon, Belén Pedrique, Roel Bakker, Michele E. Murdoch, Sake J. De Vlas, et Luc E. Coffeng. 2021. « The Burden of Skin Disease and Eye Disease Due to Onchocerciasis in Countries Formerly under the African Programme for Onchocerciasis Control Mandate for 1990, 2020, and 2030 ». Édité par Alberto Novaes Ramos. *PLOS Neglected Tropical Diseases* 15 (7): e0009604. <https://doi.org/10.1371/journal.pntd.0009604>.
- Vlaminck, Johnny, Peter U. Fischer, et Gary J. Weil. 2015. « Diagnostic Tools for Onchocerciasis Elimination Programs ». *Trends in Parasitology* 31 (11): 571-82. <https://doi.org/10.1016/j.pt.2015.06.007>.
- Volpicelli, Lorenzo, Maurizio De Angelis, Alessandra Morano, Elisa Biliotti, Cristiana Franchi, Simona Gabrielli, Simonetta Mattiucci, Carlo Di Bonaventura, et Gloria Taliani. 2020. « Encephalopathy in a Patient with Loiasis Treated with Albendazole: A Case Report ». *Parasitology International* 75 (avril):102006. <https://doi.org/10.1016/j.parint.2019.102006>.
- Walther, Michael, et Ralph Muller. 2003. « Diagnosis of Human Filariases (except Onchocerciasis) ». In *Advances in Parasitology*, 53:149-93. Elsevier. [https://doi.org/10.1016/S0065-308X\(03\)53004-4](https://doi.org/10.1016/S0065-308X(03)53004-4).
- Wanji, S., N. Tendongfor, M. Esum, S. Ndindeng, et P. Enyong. 2003. « Epidemiology of Concomitant Infections Due to Loa Loa, Mansonella Perstans, and Onchocerca Volvulus in Rain Forest Villages of Cameroon ». *Medical Microbiology and Immunology* 192 (1): 15-21. <https://doi.org/10.1007/s00430-002-0154-x>.
- Whittaker, Charles, Martin Walker, Sébastien D.S. Pion, Cédric B. Chesnais, Michel Boussinesq, et María-Gloria Basáñez. 2018. « The Population Biology and Transmission Dynamics of Loa Loa ». *Trends in Parasitology* 34 (4): 335-50. <https://doi.org/10.1016/j.pt.2017.12.003>.
- World Health Organization (WHO). 2020. « Onchocerciasis factsheet. Geneva: World Health Organization », 16. <https://www.who.int/news-room/factsheets/detail/onchocerciasis>.
- World Health Organization. 2011. « Agir pour réduire l'impact mondial des maladies tropicales négligées : premier rapport de l'OMS sur les maladies tropicales négligées ». *Working to overcome the global impact of neglected tropical diseases: first WHO report on neglected tropical diseases*. <https://iris.who.int/handle/10665/44694>.

- . 2022. « Global programme to eliminate lymphatic filariasis: progress report, 2021 », 2022. <http://www.who.int/wer>.
- World Health Organization. *Bench Aids for the Diagnosis of Filarial Infection*. Geneves: 1997. <https://iris.who.int/handle/10665/37156> Consulté Le 2024-07-11. s. d.
- World Health Organization (WHO). 2021. « Global programme to eliminate lymphatic filariasis: progress report, 2020 ». *Weekly Epidemiological Record = Relevé épidémiologique hebdomadaire* 96 (41): 497-508. CORREA.
- World Health Organization (WHO). 2022. « Plan stratégique national de lutte contre les maladies tropicales négligées 2022 - 2025 Sénégal », 144. <https://espen.afro.who.int/system/files/content/resources/Senegal%20Plan%20National%20stratégique%20MTN%202022-2025.pdf>.
- World Health Organization (WHO). 2023. « Lymphatic Filariasis ». 1 juin 2023. <https://www.who.int/fr/news-room/fact-sheets/detail/lymphatic-filariasis>.
- Wynd, Shona. 2007. « Understanding the community impact of lymphatic filariasis: a review of the sociocultural literature ». *Bulletin of the World Health Organization* 85 (6): 493-98. <https://doi.org/10.2471/BLT.06.031047>.
- Yokoly, Firmain N., Julien B. Z. Zahouli, Aboulaye Méite, Millicent Opoku, Bernard L. Kouassi, Dziedzom K. De Souza, Moses Bockarie, et Benjamin G. Koudou. 2020. « Low Transmission of Wuchereria Bancrofti in Cross-Border Districts of Côte d'Ivoire: A Great Step towards Lymphatic Filariasis Elimination in West Africa ». Édité par Justin V. Remais. *PLOS ONE* 15 (4): e0231541. <https://doi.org/10.1371/journal.pone.0231541>.
- Zouré, Honorat Gustave Marie, Samuel Wanji, Mounkaïla Noma, Uche Veronica Amazigo, Peter J. Diggle, Afework Hailemariam Tekle, et Jan H. F. Remme. 2011. « The Geographic Distribution of Loa Loa in Africa: Results of Large-Scale Implementation of the Rapid Assessment Procedure for Loiasis (RAPLOA) ». Édité par Giovanna Raso. *PLoS Neglected Tropical Diseases* 5 (6): e1210. <https://doi.org/10.1371/journal.pntd.0001210>.

CONCLUSIONS ET PERSPECTIVES

1096 Notre projet de thèse, dans son ensemble nous a permis d'approfondir nos connaissances
1097 sur l'épidémiologie, la transmission de la filariose au Sénégal et la recherche du vecteur
1098 impliqués dans la transmission de la maladie. Nous avons scindé nos travaux en trois parties en
1099 relation avec les aspects cités précédemment.

1100 Dans la première partie de cette thèse, nous avons réalisé une revue de la littérature qui
1101 se concentre sur les filarioses en Afrique de l'Ouest. Nous avons développé les différentes
1102 filarioses en Afrique de l'Ouest, leur épidémiologie, leurs pathogénicité, les méthodes de
1103 diagnostic le traitement et les moyens de prévention ciblant les arthropodes vecteurs. Ces
1104 données disponibles montrent que la filariose humaine est largement répandue en Afrique de
1105 l'Ouest. La mise en évidence des filaires dans leur rôle de transmission, ainsi que la détection
1106 du parasite adulte ou au stade de microfilaire, sont des éléments clés pour mieux comprendre
1107 l'infection. Cependant un diagnostic fiable est nécessaire pour discriminer les principales
1108 filaires permettant d'affirmer l'infection avec certitude. D'où, l'utilisation des différentes
1109 méthodes de détection des parasites chez les hôtes intermédiaires et définitifs qui revêtent une
1110 importance cruciale pour optimiser les stratégies de lutte contre les maladies parasitaires.

1111 Dans la seconde partie de ce travail, nous avons porté notre attention sur la détection des
1112 filaires chez les patients, à partir des cassettes TDR du paludisme négatif et chez les arthropodes
1113 vecteurs. Nous avons utilisé un outil moléculaire qui a mis en évidence l'espèce *M. perstans*
1114 dans les prélèvements sanguins chez les patients atteints d'éléphantiasis et les sujets fébriles.
1115 Cette même espèce a été détecté dans les cassettes TDR paludisme négatifs à *Plasmodium*
1116 *falciparum* qui constituent de bonnes sources d'ADN filarien pour la détection des infections
1117 filariennes.

1118 Nos résultats confirment la transmission de la mansonellose à *M. perstans* au Sénégal. Cette
1119 étude démontre, pour la première fois au Sénégal, la nécessité de réutiliser les TDR négatifs,
1120 pour une meilleure stratégie de surveillance de la transmission et définir son épidémiologie.
1121 Tous les moustiques et culicoïdes testés aucun n'est responsable de la transmission de la
1122 filariose.

1123 Cette étude suggère des recherches plus approfondies sur les vecteurs de la filariose au
1124 Sénégal. Bien que la PCR en temps réel et le séquençage permettent d'identifier la présence de
1125 *M. perstans*, il est recommandé de développer des tests plus rapides, accessibles et spécifiques

1126 pour une utilisation de routine dans les structures sanitaires en zone rurale. De plus, Étant donné
1127 que *Mansonella perstans* a montré une certaine insensibilité aux traitements antifilaricides
1128 courants tels que la diéthylcarbamazine et l'ivermectine, il est crucial de poursuivre des
1129 recherches pour identifier des alternatives thérapeutiques efficaces et adaptées.

1130

1131

1132

1133

1134

1135

1136

1137

1138

1139

1140

1141

1142

1143

1144

1145

1146

1147

1148

1149

1150

Références bibliographiques

- Adewole, S.O., et S.K. Ayeni. 2009. « Clinical Manifestation of Onchocerciasis in Ise - Orun Local Government, Ekiti State, Nigeria ». *Pakistan Journal of Nutrition* 8 (2): 122-24. <https://doi.org/10.3923/pjn.2009.122.124>.
- Alhassan, Andy, Zhiru Li, Catherine B. Poole, et Clotilde K.S. Carlow. 2015. « Expanding the MDx Toolbox for Filarial Diagnosis and Surveillance ». *Trends in Parasitology* 31 (8): 391-400. <https://doi.org/10.1016/j.pt.2015.04.006>.
- Anderson, Roy C. 2000. « R.C. Anderson: Nematode Parasites of Vertebrates. Their Development and Transmission. 2nd Edition. CABI Publishing, Wallingford, Oxon (UK) 2000. 650 Pp., 43 Figs. Price £ 99.50 . », 1.
- Asio, S. M., P. E. Simonsen, et A. W. Onapa. 2009. « *Mansonella Perstans* : Safety and Efficacy of Ivermectin Alone, Albendazole Alone and the Two Drugs in Combination ». *Annals of Tropical Medicine & Parasitology* 103 (1): 31-37. <https://doi.org/10.1179/136485909X384929>.
- Aubry, Arthur, Charles Hervochon, Marie Chauveau, Marie Denis-Musquer, Fakhri Jeddi, Rose-Anne Lavergne, Patrice Le Pape, et Florent Morio. 2022. « Une expulsion bien singulière d'un ver adulte de *Loa loa* ». *Revue Francophone des Laboratoires* 2022 (538): 72-75. [https://doi.org/10.1016/S1773-035X\(21\)00381-6](https://doi.org/10.1016/S1773-035X(21)00381-6).
- Bassene, Hubert, Masse Sambou, Florence Fenollar, Siân Clarke, Sawdiatou Djiba, Gaël Mourembou, Alioune Badara L Y, Didier Raoult, et Oleg Mediannikov. 2015. « High Prevalence of *Mansonella Perstans* Filariasis in Rural Senegal ». *The American Journal of Tropical Medicine and Hygiene* 93 (3): 601-6. <https://doi.org/10.4269/ajtmh.15-0051>.
- Bouchery, T., E. Lefoulon, G. Karadjian, A. Nieguitsila, et C. Martin. 2013. « The Symbiotic Role of *Wolbachia* in Onchocercidae and Its Impact on Filariasis ». *Clinical Microbiology and Infection* 19 (2): 131-40. <https://doi.org/10.1111/1469-0691.12069>.
- Chelsea, Marie, et A. Petri William. 2022. « Mansonellose-Maladies infectieuses ». <https://www.msdmanuals.com/fr/professional/maladies-infectieuses/n%C3%A9matodes-vers-ronds/mansonellose>.
- Colebunders, Robert, Jane Y Carter, Peter Claver Olore, Kai Puok, Samit Bhattacharyya, Sonia Menon, Gasim Abd-Elfarag, et al. 2018. « High Prevalence of Onchocerciasis-Associated Epilepsy in Villages in Maridi County, Republic of South Sudan: A

Community-Based Survey ». *Seizure* 63 (décembre):93-101.
<https://doi.org/10.1016/j.seizure.2018.11.004>.

Desoubeaux, Guillaume, Adélaïde Chesnay, et Jacques Chandenier. 2020. « Approches diagnostiques des filarioses en France métropolitaine ». *Revue Francophone des Laboratoires* 2020 (519): 72-80. [https://doi.org/10.1016/S1773-035X\(20\)30066-6](https://doi.org/10.1016/S1773-035X(20)30066-6).

Devi, R Renuka, et V Raju. 2018. « Information Technology in Morbidity Management of Human Lymphatic Filariasis—A Promising Tool in Global Programme for Elimination of Lymphatic Filariasis ». *Journal of Vector Borne Diseases* 55 (1): 20. <https://doi.org/10.4103/0972-9062.234622>.

Diagne, Jean Pierre, Aly Mbara Ka, Joseph Matar Ndiaye, ABoubacry Sadikh Sow, Aissatou Maguatte Wane, et Paule Aida Ndoeye Roth. 2018. « MIGRATION SOUS CUTANEE PALPEBRALE D'UNE LOASE: A PROPOS D'UN CAS », n° 1, 4. https://www.revues-ufhb-ci.org/fichiers/FICHIR_ARTICLE_2511.pdf.

Diwakar, Deepak Kumar, Nikita Wadhwani, Nikhil Prasad, et Ashwani Gupta. 2018. « Primary Retroperitoneal Filariasis: A Common Disease of Tropics with Uncommon Presentation and Review of Literature ». *BMJ Case Reports*, septembre, bcr-2018-226217. <https://doi.org/10.1136/bcr-2018-226217>.

Ehrens, Alexandra, Achim Hoerauf, et Marc P. Hübner. 2022. « Eosinophils in filarial infections: Inducers of protection or pathology? » *Frontiers in Immunology* 13 (octobre):983812. <https://doi.org/10.3389/fimmu.2022.983812>.

Faye, O., B. Faye, J. L. Ndiaye, R. C. K. Tine, D. Ndiaye, et A. Haydara Mbacké. 2009. « [Epidemiological study on lymphatic filariasis (*Wuchereria bancrofti*) in Sénégal] ». *Medecine Tropicale: Revue Du Corps De Sante Coloniale* 69 (1): 101-2.

Gardon, Jacques, Nathalie Gardon-Wendel, Demanga-Ngangue, Joseph Kamgno, Jean-Philippe Chippaux, et Michel Boussinesq. 1997. « Serious Reactions after Mass Treatment of Onchocerciasis with Ivermectin in an Area Endemic for Loa Loa Infection ». *The Lancet* 350 (9070): 18-22. [https://doi.org/10.1016/S0140-6736\(96\)11094-1](https://doi.org/10.1016/S0140-6736(96)11094-1).

Hotez, Peter J., et Aruna Kamath. 2009. « Neglected Tropical Diseases in Sub-Saharan Africa: Review of Their Prevalence, Distribution, and Disease Burden ». Édité par Michael Cappello. *PLoS Neglected Tropical Diseases* 3 (8): e412. <https://doi.org/10.1371/journal.pntd.0000412>.

Ibe, Ogochukwu, Obinna Onwujekwe, Benjamin Uzochukwu, Miriam Ajuba, et Paul Okonkwo. 2015. « Exploring Consumer Perceptions and Economic Burden of

1217 Onchocerciasis on Households in Enugu State, South-East Nigeria ». Édité par Samuel
1218 V. Scarpino. *PLOS Neglected Tropical Diseases* 9 (11): e0004231.
1219 <https://doi.org/10.1371/journal.pntd.0004231>.

1220 Katawa, Gnatoulma, Laura E. Layland, Alex Y. Debrah, Charlotte Von Horn, Linda Batsa,
1221 Alexander Kwarteng, Sandra Arriens, et al. 2015. « Hyperreactive Onchocerciasis Is
1222 Characterized by a Combination of Th17-Th2 Immune Responses and Reduced
1223 Regulatory T Cells ». Édité par Benjamin L. Makepeace. *PLoS Neglected Tropical*
1224 *Diseases* 9 (1): e3414. <https://doi.org/10.1371/journal.pntd.0003414>.

1225 Katiyar, D., et L. K. Singh. 2011. « Filariasis: Current Status, Treatment and Recent Advances
1226 in Drug Development ». *Current Medicinal Chemistry* 18 (14): 2174-85.
1227 <https://doi.org/10.2174/092986711795656234>.

1228 Lustigman, Sara, Alexandra Grote, et Elodie Ghedin. 2017. « The Role of “omics” in the Quest
1229 to Eliminate Human Filariasis ». Édité par Judd L. Walson. *PLOS Neglected Tropical*
1230 *Diseases* 11 (4): e0005464. <https://doi.org/10.1371/journal.pntd.0005464>.

1231 M. Boussinesq. 2007. « Actualités thérapeutiques dans les filarioses – Recent advances in the
1232 treatment of filariases ». *mise au point*, n° 2, 55.

1233 Manguin, S., M.J. Bangs, J. Pothikasikorn, et T. Chareonviriyaphap. 2010. « Review on Global
1234 Co-Transmission of Human Plasmodium Species and Wuchereria Bancrofti by
1235 Anopheles Mosquitoes ». *Infection, Genetics and Evolution* 10 (2): 159-77.
1236 <https://doi.org/10.1016/j.meegid.2009.11.014>.

1237 Medeiros, Jansen Fernandes, James Lee Crainey, Felipe Arley Costa Pessoa, et Sérgio Luis
1238 Bessa Luz. 2017. « Mansonelliasis ». In *Arthropod Borne Diseases*, édité par Carlos
1239 Brisola Marcondes, 405-26. Cham: Springer International Publishing.
1240 https://doi.org/10.1007/978-3-319-13884-8_26.

1241 Mendoza, Natalia, Alexa Li, Amber Gill, et Stephen Tying. 2009. « Filariasis: Diagnosis and
1242 Treatment ». *Dermatologic Therapy* 22 (6): 475-90. <https://doi.org/10.1111/j.1529-8019.2009.01271.x>.

1244 Mourembou, Gaël, Florence Fenollar, Jean Bernard Lekana-Douki, Angelique Ndjoi
1245 Mbiguino, Sydney Maghendji Nzondo, Pierre Blaise Matsiegui, Rella Zoleko Manego,
1246 et al. 2015. « Mansonella, Including a Potential New Species, as Common Parasites in
1247 Children in Gabon ». Édité par Dunstan Mukoko. *PLOS Neglected Tropical Diseases* 9
1248 (10): e0004155. <https://doi.org/10.1371/journal.pntd.0004155>.

1249 Njim, Tsi, Joel Mbigha Ngum, et Leopold Ndemnge Aminde. 2015. « Cutaneous
1250 Onchocerciasis in Dumbu, a Pastoral Area in the North-West Region of Cameroon:

Diagnostic Challenge and Socio-Economic Implications ». *Pan African Medical Journal* 22. <https://doi.org/10.11604/pamj.2015.22.298.7707>.

Opoku, Millicent, Corrado Minetti, Worlasi D. Kartey-Attipoe, Sampson Otoo, Joseph Otchere, Bruno Gomes, Dziedzom K. De Souza, et Lisa J. Reimer. 2018. « An Assessment of Mosquito Collection Techniques for Xenomonitoring of Anopheline-Transmitted Lymphatic Filariasis in Ghana ». *Parasitology* 145 (13): 1783-91. <https://doi.org/10.1017/S0031182018000938>.

Ottesen, Eric A. 2006. « Lymphatic Filariasis: Treatment, Control and Elimination ». In *Advances in Parasitology*, 61:395-441. Elsevier. [https://doi.org/10.1016/S0065-308X\(05\)61010-X](https://doi.org/10.1016/S0065-308X(05)61010-X).

Pérez-Molina, José Antonio, Marta Díaz-Menéndez, Ana Pérez-Ayala, Federico Ferrere, Begoña Monje, Francesca Norman, et Rogelio López-Vélez. 2010. « Tratamiento de las enfermedades causadas por parásitos ». *Enfermedades Infecciosas y Microbiología Clínica* 28 (1): 44-59. <https://doi.org/10.1016/j.eimc.2009.11.003>.

Ramaiah, K. D., et Eric A. Ottesen. 2014. « Progress and Impact of 13 Years of the Global Programme to Eliminate Lymphatic Filariasis on Reducing the Burden of Filarial Disease ». Édité par Moses Bockarie. *PLoS Neglected Tropical Diseases* 8 (11): e3319. <https://doi.org/10.1371/journal.pntd.0003319>.

Rodi, Miriam, Caspar Gross, Thaisa Lucas Sandri, Lilith Berner, Marina Marcet-Houben, Ersoy Kocak, Michaela Pogoda, et al. 2023. « Whole genome analysis of two sympatric human *Mansonella*: *Mansonella perstans* and *Mansonella* sp “DEUX” ». *Frontiers in Cellular and Infection Microbiology* 13 (avril):1159814. <https://doi.org/10.3389/fcimb.2023.1159814>.

Sandri, Thaisa Lucas, Andrea Kreidenweiss, Simon Cavallo, David Weber, Sascha Juhas, Miriam Rodi, Tamirat Gebru Woldearegai, et al. 2021. « Molecular Epidemiology of *Mansonella* Species in Gabon ». *The Journal of Infectious Diseases* 223 (2): 287-96. <https://doi.org/10.1093/infdis/jiaa670>.

Simonsen, Paul E., Ambrose W. Onapa, et Santa Maria Asio. 2011. « *Mansonella Perstans* Filariasis in Africa ». *Acta Tropica, The Diagnostics and Control of Neglected Tropical Helminth Diseases*, 120 (septembre):S109-20. <https://doi.org/10.1016/j.actatropica.2010.01.014>.

Tahita, Marc Christian, Thuy-Huong Ta-Tang, Bérénger Kaboré, Raquel Capote-Morales, Irene Molina De La Fuente, Raquel Cruces, Hamidou Ilboudo, et al. 2024. « Molecular Detection of *Wuchereria Bancrofti*, *Loa Loa* and *Mansonella Perstans* from Dried Blood

1285 Spots Taken from Pregnant Women in Rural Burkina Faso ». *African Journal of*
1286 *Parasitology, Mycology and Entomology* 1 (2): 1.
1287 <https://doi.org/10.35995/ajpme2010002>.

1288 Ta-Tang, Thuy-Huong, James Crainey, Rory J. Post, Sergio LB. Luz, et José Rubio. 2018.
1289 « Mansonellosis: Current Perspectives ». *Research and Reports in Tropical Medicine*
1290 Volume 9 (janvier):9-24. <https://doi.org/10.2147/RRTM.S125750>.

1291 Ta-Tang, Thuy-Huong, Sergio Lb Luz, James L Crainey, et José M Rubio. 2021. « An
1292 Overview of the Management of Mansonellosis ». *Research and Reports in Tropical*
1293 *Medicine* Volume 12 (mai):93-105. <https://doi.org/10.2147/RRTM.S274684>.

1294 Vinkeles Melchers, Natalie V. S., Sarah Mollenkopf, Robert Colebunders, Michael Edlinger,
1295 Luc E. Coffeng, Julia Irani, Trésor Zola, et al. 2018. « Burden of Onchocerciasis-
1296 Associated Epilepsy: First Estimates and Research Priorities ». *Infectious Diseases of*
1297 *Poverty* 7 (1). <https://doi.org/10.1186/s40249-018-0481-9>.

1298 Walther, Michael, et Ralph Muller. 2003. « Diagnosis of Human Filariases (except
1299 Onchocerciasis) ». In *Advances in Parasitology*, 53:149-93. Elsevier.
1300 [https://doi.org/10.1016/S0065-308X\(03\)53004-4](https://doi.org/10.1016/S0065-308X(03)53004-4).

1301 World Health Organization (WHO). 2020. « Onchocerciasis factsheet. Geneva: World Health
1302 Organization », 16. <https://www.who.int/news-room/factsheets/detail/onchocerciasis>.

1303 World Health Organization. 2022. « Global programme to eliminate lymphatic filariasis:
1304 progress report, 2021 », 2022. <http://www.who.int/wer>.

1305 World Health Organization (WHO). 2022. « Plan stratégique national de lutte contre les
1306 maladies tropicales négligées 2022 - 2025 Sénégal », 144.
1307 [https://espen.afro.who.int/system/files/content/resources/Senegal%20Plan%20Nationa](https://espen.afro.who.int/system/files/content/resources/Senegal%20Plan%20National%20strat%C3%A9gique%20MTN%202022-2025.pdf)
1308 [l%20strat%C3%A9gique%20MTN%202022-2025.pdf](https://espen.afro.who.int/system/files/content/resources/Senegal%20Plan%20National%20strat%C3%A9gique%20MTN%202022-2025.pdf).

1309 Zouré, Honorat Gustave Marie, Samuel Wanji, Mounkaïla Noma, Uche Veronica Amazigo,
1310 Peter J. Diggle, Afework Hailemariam Tekle, et Jan H. F. Remme. 2011. « The
1311 Geographic Distribution of Loa Loa in Africa: Results of Large-Scale Implementation
1312 of the Rapid Assessment Procedure for Loiasis (RAPLOA) ». Édité par Giovanna Raso.
1313 *PLoS Neglected Tropical Diseases* 5 (6): e1210.
1314 <https://doi.org/10.1371/journal.pntd.0001210>.

1315

1316